The dirac equation with light-cone data

Ottesen, Johnny T.

Publication date:
1988

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Ottesen, J. T. (1988). *The dirac equation with light-cone data*. Roskilde Universitet.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

TEKST NR 171

1988

The Dirac Equation with Light-Cone Data.

Johnny Tom Ottesen

TEKSTER fra

IMFUFA

ROSKILDE UNIVERSITETSCENTER
INSTITUT FOR STUDIET AF MATEMATIK OG FYSIK SAMT DERES
FUNKTIONER I UNDERVISNING, FORSKNING OG ANVENDELSER

IMFUFA, Roskilde Universitetscenter, Postbox 260,
4000 Roskilde.

The Dirac Equation with Light-Cone Data

af: Johnny Tom Ottesen

IMFUFA tekst nr. 171/88

29 sider

ISSN 0106-6242

ABSTRACT

The Dirac equation with light-cone data ;

$$(i\gamma^0 \frac{\partial}{\partial t} + i\gamma \cdot \nabla - m\mathbb{1})u(t,x) = 0.$$

$$u(-|x|,x) = f(x), \quad x \in \mathbb{R}^3$$

is considered and explicitly solved in terms of a "Light-cone

Fourier-transform" under appropriate conditions on f .

1. Introduction.

We shall consider the characteristic Cauchy problem for the free Dirac equation

$$(i\gamma^0 \frac{\partial}{\partial t} + i\gamma^j \nabla_j - m\mathbb{1})u(t,x) = 0 \quad (1.1)$$

$$u(-|x|,x) = f(x) \quad (1.2)$$

under suitable conditions on f , in four space-time dimensions.

The analogous problem for the Klein-Gordon equation was considered in [1]. The wave-equation has been considered by Riesz [2] and Strichartz [3], but in a very different way. For some general results on characteristic Cauchy-problems see Hörmander [4].

We shall show that there exist a Hilbert-space $\mathcal{H}$ of light-cone data such that (1.1) and (1.2) have a unique weak solution $u(t,x)$, with $u(t,\cdot) \in L^2(\mathbb{R}^3, \mathbb{C}^4)$ for f in $\mathcal{H}$ and so that all " L^2 -solutions" of (1.1) have the property that $u(t-|\cdot|,\cdot)$ is in $\mathcal{H}$.

The Dirac equation (1.1) will be rewritten, as an evolution

equation $i \frac{\partial u}{\partial t} = Hu$ in $\mathcal{H}$ and the generator H will be diagonalized

by a "light-cone Fourier transform", which then allows us to give an explicit formula for $u(t,x)$ in terms of f .

2. The Dirac Equation and Associated Evolution Equations.

The Dirac equation is a hyperbolic system of linear partial differential equations which can be written as

$$(i\gamma^0 \frac{\partial}{\partial t} + i\gamma \cdot \nabla - m\mathbb{1})u(t,x) = 0 \quad (2.1)$$

where $m \neq 0$ is the mass of the particle, $u: \mathbb{R} \times \mathbb{R}^3 \rightarrow \mathbb{C}^4$, γ^0 and the three components of the vector $\gamma = (\gamma^1, \gamma^2, \gamma^3)$ are 4×4 complex matrices fulfilling the anti-commutation relation

$$\gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu = 2g^{\mu\nu} \mathbb{1} \quad (2.2)$$

where $g^{\mu\nu} = 0$ for $\mu \neq \nu$, $g^{00} = -g^{ii} = 1$, for $i = 1, 2, 3$, and $\mathbb{1}$ is the 4×4 unit matrix. An explicit representation of the γ^μ 's is given by

$$\gamma^0 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad \text{and} \quad \gamma^j = \begin{bmatrix} 0 & -\sigma^j \\ \sigma^j & 0 \end{bmatrix}, \quad \text{where } 1 \text{ in } \gamma^0 \text{ is the}$$

2×2 unit matrix and the σ^j 's are the familiar Pauli-matrices given by

$$\sigma^1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \sigma^2 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \sigma^3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}. \quad \text{We shall use this representation}$$

in the following.

The Dirac equation (2.1) is often considered as an evolution equation with initial data on a space-like hyperplane, consider for example a $t = \text{constant}$ hyperplane and write the equation as

$$i \frac{\partial u}{\partial t} = \gamma^0(-i\gamma \cdot \nabla + m1)u, \quad \text{which can be considered as an evolution}$$

equation in the Hilbert-space $L^2(\mathbb{R}^3, \mathbb{C}^4)$. The generator

$D = \gamma^0(-i\gamma \cdot \nabla + m1)$ defines a self-adjoint operator D with domain

$\mathcal{D}(D) = H^1(\mathbb{R}^3, \mathbb{C}^4)$, the direct sum of the four identical Sobolev

spaces $H^1(\mathbb{R}^3)$. The associated spectral representation of a solution

$e^{-iDt}f$ can be written

$$u(t, x) = (2\pi)^{-3/2} \sum_{\pm} \sum_{s=1,2} \int_{\mathbb{R}^3} \frac{d^3k}{\omega_k} e^{\mp i\omega_k t + ik \cdot x} u_{\pm}(k, s) \hat{f}_{\pm}(k, s) \quad (2.3)$$

for $f(\cdot) \in \mathcal{S}(\mathbb{R}^3, \mathbb{C}^4)$, the direct sum of the four identical Schwartz-spaces $\mathcal{S}(\mathbb{R}^3)$, where $u_{\pm}(k, s)$, $s = 1, 2$, are four orthogonal eigenvectors of $\hat{D} = \gamma^0(\gamma \cdot k + m\mathbb{1})$ normalized such that $u_{\pm}^* u_{\pm} = \omega_k = (|k|^2 + m^2)^{1/2}$. The "Fourier-components" $\hat{f}_{\pm}(\cdot, s)$ is essentially given by the ordinary Fourier-Plancherel transformation,

since $L^2(\mathbb{R}^3, \mathbb{C}^4) = \bigoplus_{n=1}^4 L^2(\mathbb{R}^3)$ it follows that the ordinary Fourier-

Plancherel transform lifts in a canonical way to a unitary operator $F_0 : L^2(\mathbb{R}_x^3, \mathbb{C}^4) \rightarrow L^2(\mathbb{R}_k^3, \mathbb{C}^4)$ such that

$F_0 D F_0^* = \hat{D} = \gamma^0(\gamma \cdot k + m\mathbb{1})$ and $\hat{f} = F_0 f$. For $f(\cdot) \in \mathcal{S}(\mathbb{R}^3, \mathbb{C}^4)$ we have the following explicit formula

$$(F_0 f)_{\pm}(k, s) = \langle v_{\pm}(\cdot, k, s), f(\cdot) \rangle_{L^2(\mathbb{R}_x^3, \mathbb{C}^4)} \quad (2.4)$$

Where

$$v_{\pm}(x, k, s) = (2\pi)^{-3/2} e^{ik \cdot x} u_{\pm}(k, s) \quad (2.5)$$

The associated Parseval's formula reads

$$\int_{\mathbb{R}^3} d^3x \, u^*(t,x)u(t,x) = \sum_{\pm} \sum_{s=1,2} \int_{\mathbb{R}^3} \frac{d^3k}{\omega_k} \left| \hat{f}_{\pm}(k,s) \right|^2 \quad (2.6)$$

for every $t \in \mathbb{R}$. Consider a solution $u(t,x)$ such that $\hat{f}_{\pm} \in \mathcal{S}(\mathbb{R}^3)$ and construct the current $J^{\mu} = \bar{u} \gamma^{\mu} u$, where $\bar{u} = u^* \gamma^0$. It follows from the Dirac equation (2.1) that J^{μ} is conserved, i.e.

$$\nabla_{\mu} J^{\mu} \equiv \frac{\partial}{\partial t} J^0 + \nabla \cdot J = 0. \text{ Intergration of this conservation equation}$$

over the region in $\mathbb{R} \times \mathbb{R}^3$ between a $t=t_0$ hyperplane, $t_0 \in \mathbb{R}$, and the ligh-cone $\mathfrak{C} = \{ (t,x) : t=t'-|x|, t' < t_0 \}$ together with Gauss's theorem and a decay estimate, given in [1], gives

$$\begin{aligned} & \int_{\mathbb{R}^3} d^3x \, \bar{u}(t'-|x|,x) (\gamma^0 + \gamma \cdot n) u(t'-|x|,x) \\ &= \sum_{s=1,2} \int_{\mathbb{R}^3} \frac{d^3k}{\omega_k} \left(\left| \hat{f}_{+}(k,s) \right|^2 + \left| \hat{f}_{-}(k,s) \right|^2 \right) \end{aligned} \quad (2.7)$$

where $n = x|x|^{-1}$, equation (2.7) resembles a Parseval's formula.

This indicates that there might exist a Hilbert-space $\mathcal{H}$ of lighth-cone data $u(t-|\cdot|, \cdot)$ with the left-hand side of (2.7) as scalar product and such that the map $u(-|\cdot|, \cdot) \rightarrow u(t-|\cdot|, \cdot)$ defines a unitary strongly continuous one-parameter group $t \rightarrow U(t)$ on $\mathcal{H}$. This will be shown to be the case in the next section.

Let $u(0, x)$ have support in the region $|x| \leq R$ and consider $t \geq R$. It then follows from the finite propagation velocity that the support of $u(t-|x|, x)$ is in the region $|x| \leq (R+t)/2$. This means that there is a subspace $\mathcal{H}_0 \subset \mathcal{H}$ of lighth-cone data with compact support.

It follows again from the finite propagation velocity, that $U(t)\mathcal{H}_0 \subset \mathcal{H}_0$, for $t \geq 0$. This need not be the case for $t < 0$, which means that compact support of lighth-cone data, is in general, only a semi-group property.

3. The light-cone evolution equation.

Consider the Dirac equation (2.1) and make the coordinate exchange $t \rightarrow t' = t + |x|$ and $x \rightarrow x' = x$ such that a $t' = \text{constant}$ is the backward light-cone with its apex at $(t, 0)$. The new coordinates will be called light-cone (LC) coordinates. In the LC coordinates the Dirac

equation becomes $i(1 + \alpha \cdot n') \frac{\partial u'}{\partial t'} = (-i\alpha \cdot \nabla' + m\gamma^0)u'$ where

$u'(t', x') = u(t' - |x'|, x')$, $\alpha = \gamma^0 \gamma$ and $n' = x' |x'|^{-1}$, note that ∇' is associated with the light-cone $\mathfrak{C} = \{(t, x) : t = -|x|\}$ in the original coordinates and $\mathfrak{C} = \{x' \in \mathbb{R}^3\}$ in the new coordinates, i.e. the light-cone can be identified with $\mathbb{R}^3$ which will be done in the following. We shall only use the LC coordinates from now on and can therefore drop the primes. Hence the Dirac equation becomes

$$i(1 + \alpha \cdot n) \frac{\partial u}{\partial t} = (-i\alpha \cdot \nabla + m\gamma^0)u \quad (3.1)$$

where $n = x |x|^{-1}$ and $\alpha = \gamma^0 \gamma$.

Definition 3.1. Let $L^2 = L^2(\mathfrak{E}, \mathbb{C}^4)$ where the measure on $\mathfrak{E}$ is the Lebesgue measure on $\mathbb{R}^3$. Put

$$Q = \mathbb{1} + \alpha \cdot n \quad \text{and} \quad D = -i\alpha \cdot \nabla + \gamma^0 m \quad (3.2)$$

It then follows that $2^{-1}Q$ is an orthogonal projection in L^2 and that the Dirac operator D defines a self-adjoint operator in L^2 with spectrum $\sigma(D) = (-\infty, -m] \cup [m, \infty)$, $m > 0$.

The Dirac equation (3.1) can now be considered as an equation in L^2

$$iQ \frac{du}{dt}(t) = Du(t) \quad (3.3)$$

which can be written as

$$iD^{-1}Q \frac{du}{dt}(t) = u(t) \quad (3.4)$$

since $0 \notin \sigma(D)$. From equation (3.4) it follows that if $\frac{du}{dt} \in L^2$

then $u \in D^{-1}QL^2$.

Definition 3.2. Let $B = D^{-1}Q$ and $\mathcal{H}_0 = BL^2$. Furthermore put

$$\langle f, g \rangle_{\mathcal{H}_0} = \langle f, Qg \rangle_{L^2} \quad (3.5)$$

for $f, g \in \mathcal{H}_0$.

It will be proved that $\mathcal{H}_0$ is a pre-Hilbert space, but before doing that we have to mention two lemmas, both proved in [1].

Lemma 3.3. Let $S_0 = -i(n \cdot \nabla + \nabla \cdot n)$ with domain $\mathcal{D}(S_0) = C_0^\infty(\mathbb{R}^3)$

in $L^2(\mathbb{R}^3)$, then the following inequality holds

$$\|S_0 f\|_{L^2(\mathbb{R}^3)} \leq 4 \int_{\mathbb{R}^3} d^3x |\nabla f(x)|^2 \quad (3.6)$$

for all $f \in \mathcal{D}(S_0)$.

Lemma 3.4. The range $\mathcal{R}(S_0)$ of S_0 is dense in $L^2(\mathbb{R}^3)$.

Due to the inequality (3.6) S_0 can be extended continuously to the Sobolev space $H^1(\mathbb{R}^3)$, let this extension be denoted S , it also follows that S is symmetric in $L^2(\mathbb{R}^3)$, since S_0 is. Due to lemma 3.4. also S have dense range $\mathcal{R}(S_0)$ in $L^2(\mathbb{R}^3)$, and since S is symmetric in $L^2(\mathbb{R}^3)$ it follows that S is one-to-one, i.e. S has a densely defined inverse in $L^2(\mathbb{R}^3)$, in fact it is easy to show that

$$(S_0^{-1}g)(x) = \frac{i}{2r} \int_{-\infty}^r \rho g(\rho, x|x|^{-1}) d\rho \quad (3.7)$$

for $g \in \mathcal{R}(S_0)$.

Proposition 3.5. $\mathcal{H}_0$ is a pre-Hilbert space.

Proof. The non-trivial part is to prove that if $\langle f, f \rangle_{\mathcal{H}_0} = 0$ then $f = 0$

and since $\langle f, f \rangle_{\mathcal{H}_0} = \langle f, Qf \rangle_{L^2}$ this is to prove that $Qf = 0$ implies

$f = 0$ in L^2 . Let $f \in \mathcal{H}_0 \setminus \{0\}$. From the anti-commutation relation (2.2)

it follows by a direct computation that

$$(\alpha \cdot n)(\alpha \cdot \nabla) + (\alpha \cdot \nabla)(\alpha \cdot n) = (n \cdot \nabla + \nabla n) \mathbf{1} \quad (3.8)$$

on $C_0^\infty(\mathbb{R}^3, \mathbb{C}^4)$. In section 2 it was shown that the Dirac operator D in

L^2 is self-adjoint with domain $\mathcal{D}(D) = H^1(\mathbb{R}^3, \mathbb{C}^4)$. Now the above

defined operator S can be lifted in a canonical way to an operator $S \cdot \mathbf{1}$

on $\mathcal{D}(D)$, where $\mathbf{1}$ here denotes the 4×4 unit matrix. Hence

$$DQ = (2 - Q)D + S \cdot \mathbf{1} \quad (3.9)$$

on $\mathcal{D}(D)$, since the left-hand side and the right-hand side agree on the

dense subspace $C_0^\infty(\mathbb{R}^3, \mathbb{C}^4)$. Since $0 \notin \sigma(D)$ then D have a bound invers

D^{-1} defined on $\mathcal{R}(D)$, which is dense in L^2 , hence D^{-1} can be extended

to a continuous operator on L^2 also denoted by D^{-1} . Acting on (3.9) with D^{-1} from both left and right gives

$$QD^{-1} = D^{-1}(2-Q) + D^{-1}S \cdot 1D^{-1} \quad (3.10)$$

on $\mathcal{R}(D)$. By acting with Q from both left and right equation (3.10) gives

$$QB = D^{-1}S \cdot 1B \quad (3.11)$$

on $\{f \in L^2 : Qf \in \mathcal{R}(D)\}$, since $Q^2 = 2Q$. Now $f = Bg \in \mathcal{H}_0 \setminus \{0\}$ so it follows from equation (3.11) that $Qf = QBg = D^{-1}S \cdot 1Bg$ and since D^{-1} is a bounded operator and the kernel for S is trivial in $L^2(\mathbb{R}^3)$ then $D^{-1}S \cdot 1Bg \neq 0$, i.e. if $f \in \mathcal{H}_0 \setminus \{0\}$ then $Qf \in L^2 \setminus \{0\}$, or in other words if $Qf = 0$ in L^2 then $f = 0$ in $\mathcal{H}_0$.

Definition 3.6. Let $\mathcal{H}$ be the completion of $\mathcal{H}_0$.

It follows from the definition that $2^{-1/2}Q$ defines an isometry from $\mathcal{H}$ into L^2 , and that B defines a bounded self-adjoint operator on $\mathcal{H}$.

Proposition 3.7. The range $\mathcal{R}(B)$ of $B : \mathcal{H} \rightarrow \mathcal{H}$ is dense in $\mathcal{H}$.

Proof. Let $\psi \in (B\mathcal{H})^\perp$, i.e. ψ is in the orthogonal complement of $B\mathcal{H}$,

so $\langle \psi, Bf \rangle_{\mathcal{H}} = 0$ for all $f \in \mathcal{H}$. Consider $f \in \mathcal{H}_0$, then $f = Bg$ for some

$g \in L^2$. Then $\langle \psi, Bf \rangle = 0$ for all $f \in \mathcal{H}$ implies that

$\langle \psi, QD^{-1}QD^{-1}Qg \rangle_{L^2} = 0$ for all $g \in L^2$. Since $QD^{-1}QD^{-1}Q = (D^{-1}S \cdot 1D^{-1})^2Q$

and the commutator $[Q, D^{-1}S \cdot 1D^{-1}] = 0$ then

$$\begin{aligned} \langle (D^{-1}S \cdot 1D^{-1})^2Q\psi, g \rangle_{L^2} &= \langle QD^{-1}QD^{-1}Q\psi, g \rangle_{L^2} \\ &= \langle \psi, QD^{-1}QD^{-1}Qg \rangle_{L^2} \\ &= \langle \psi, D^{-1}QD^{-1}Qg \rangle_{L^2} = 0 \end{aligned} \quad (3.12)$$

for all $g \in L^2$, i.e. $(D^{-1}S \cdot 1D^{-1})^2Q\psi = 0$. Hence $Q\psi = 0$ in L^2 , since D^{-1}

is invertible and the kernel for S is trivial in $L^2(\mathbb{R}^3)$, i.e. $\psi = 0$ in $\mathcal{H}$

(which is proved in proposition 3.5.)

Proposition 3.7. shows that B is one-to-one since it is self-adjoint.

Definition 3.8. Put $H = B^{-1}$ with $\mathcal{D}(H) = \mathcal{R}(B)$ in $\mathcal{H}$.

Note that H is self-adjoint since B is. Equation (3.4) can then be written as

$$i \frac{du}{dt}(t) = Hu(t) \quad (3.13)$$

with the solution $u(t) = e^{-iHt}u(0)$, $u(0) \in \mathcal{D}(H)$, i.e. $U_t = e^{-iHt}$ is the time-evolution operator for light-cone data. From definition 3.8 it follows that

$$\langle f, Hf \rangle_{\mathcal{H}} = \langle f, Df \rangle_{L^2} \quad (3.14)$$

and

$$\langle Hf, Hf \rangle_{\mathcal{H}} = \langle Df, Df \rangle_{L^2} \quad (3.15)$$

for all $f \in \mathcal{D}(H)$, i.e. $\mathcal{D}(H) \subset \mathcal{D}(D) = H^1(\mathbb{R}^3, \mathbb{C}^4)$.

4. The light-cone Fourier transform.

The self-adjointness of H implies, due to the spectral theorem, that

there exist a spectral family $\{E(\lambda)\}_{\lambda \in \mathbb{R}}$ such that $H = \int_{\mathbb{R}} \lambda dE(\lambda)$.

The following well-known formula allows $E(\lambda)$ to be expressed in terms of the resolvent $R(z) = (H-z)^{-1}$, for z in the resolvent set $\rho(H)$

$$\begin{aligned} & \langle f, 1/2[(E(\beta)+E(\beta-0))-(E(\alpha)+E(\alpha-0))]f \rangle_{\mathcal{H}} \\ &= \lim_{\varepsilon \searrow 0} \frac{1}{2\pi i} \int_{\alpha}^{\beta} d\mu \langle f, [R(\mu+i\varepsilon)-R(\mu-i\varepsilon)]f \rangle_{\mathcal{H}} \end{aligned} \quad (4.1)$$

In the following the right-hand side of equation (4.1) will be evaluated.

Let f belong to $C_0^{\infty}(\mathbb{R}^3, \mathbb{C}^4)$ and put $g = R(z)f$ for $\text{Im}(z) \neq 0$.

It then follows that g belongs to $\mathcal{D}(H)$ and fulfills the equation

$$(H-z)g = f \quad (4.2)$$

which also can be written as $(B^{-1}-z)g = f$. Now applying B to this equation gives $(1-zB)g = Bf$. Consider this equation in L^2 , then the fact that $\mathcal{D}(H) \subset \mathcal{D}(D)$ allows us to act with D , which gives

$$(D-zQ)g = Qf \tag{4.3}$$

since $B = D^{-1}Q$.

Definition 4.1. Let V_z be defined in L^2 by

$$(V_z\varphi)(x) = e^{-iz|x|}\varphi(x) \tag{4.4}$$

on its maximal domain.

Proposition 4.2. If φ is in L^2 such that $V_z^{-1}\varphi$ is in $\mathcal{D}(D)$ then

$$(D-zQ)V_z^{-1}\varphi = V_z^{-1}(D-z)\varphi \tag{4.5}$$

Proof. Equation (4.5) follows from a straightforward computation.

Rewriting equation (4.3) as $(D-zQ)V_z^{-1}V_zg = Qf$ and using

proposition 4.2 gives

$$V_z^{-1}(D-z)V_z g = f \quad (4.6)$$

for $\text{Im}(z) < 0$, such that V_z is bounded. Equation (4.6) can be solved for g , which gives

$$g = V_z^{-1}(D-z)^{-1}V_z Qf \quad (4.7)$$

and then

$$\langle f, R(z)f \rangle_{\mathcal{H}} = \langle V_{\bar{z}} Qf, (D-z)^{-1}V_z Qf \rangle_{L^2} \quad (4.8)$$

for $\text{Im}(z) < 0$. The left-hand side of (4.8) is analytic for z not in the spectra $\sigma(H)$ of H , which then shows that the right-hand side has an analytic continuation to $\text{Im}(z) > 0$.

As described in section 2 the spectral theory for the Dirac operator D in L^2 , is given by a unitary operator F_0 on L^2 , such that

$F_0 D F_0^* = \hat{D} = \alpha \cdot k + \gamma^0 m$, $k \in \mathbb{R}^3$, with eigenvalues $\pm \omega_k$. For $f \in L^2$

put for short $\hat{f} = F_0 f$. Equation (4.8) can then be written as

$$\langle f, R(z)f \rangle_{\mathcal{H}} = \langle \widehat{V_{\bar{z}} Q f}, (\hat{D} - z)^{-1} \widehat{V_z Q f} \rangle_{L^2} \quad (4.9)$$

where $(\hat{D} - z)^{-1}$ has the following explicit representation

$$(\hat{D} - z)^{-1} = \omega_k^{-1} \sum_{\pm} \sum_{s=1,2} (\pm \omega_k - z)^{-1} u_{\pm}(k,s) u_{\pm}^*(k,s) \quad (4.10)$$

and $u(\cdot, s)$ are four orthogonal eigenvectors of $\hat{D}$ normalized such

that $u_{\pm}^*(k,s) u_{\pm}(k,s) = \omega_k$ also described in section 2.

Definition 4.3. If $f \in C_0^{\infty}(\mathbb{R}^3, \mathbb{C}^4)$ then $f(\cdot, s)$ is defined by

$$f_{\pm}(k,s) = u_{\pm}^*(k,s) (\widehat{V_{\pm \omega_k} Q f})(k) \quad (4.11)$$

for $s = 1, 2$.

With proper interpretation formula (4.11) also can be written as

$$\widehat{f}_{\pm}(k,s) = \langle u_{\pm}(\cdot, k, s), f(\cdot) \rangle_{\mathcal{H}} \quad (4.12)$$

where

$$u_{\pm}(x, k, s) = (2\pi)^{-3/2} e^{\pm i\omega_k |x| + ik \cdot x} u_{\pm}(k, s) \quad (4.13)$$

Now the right-hand side of equation (4.1) can be evaluated. Let

$f \in C_0^{\infty}(\mathbb{R}^3, \mathbb{C}^4)$ and put

$$g_{\pm}(k, s, z) = (u_{\pm}^*(k, s) \widehat{(V_{\bar{z}} Q f)}(k)) * (u_{\pm}^*(k, s) \widehat{(V_z Q f)}(k)) \quad (4.14)$$

then formula (4.9) becomes

$$\langle f, R(z)f \rangle_{\mathcal{H}} = \sum_{\pm} \sum_{s=1,2} \int_{\mathbb{R}^3} \frac{d^3 k}{\omega_k} (\pm \omega_k - z)^{-1} g_{\pm}(k, s, z) \quad (4.15)$$

Due to the fact that $g_{\pm}$ is analytic in z and continuous in k

Fubini's theorem can be used on equation (4.15) which gives

$$\begin{aligned} & \lim_{\epsilon \searrow 0} \frac{1}{2\pi i} \int_{\alpha}^{\beta} d\mu \langle f, [R(\mu+i\epsilon) - R(\mu-i\epsilon)] f \rangle_{\mathcal{H}} \\ &= \sum_{\pm} \sum_{s=1,2} \lim_{\epsilon \searrow 0} \int_{\mathbb{R}^3} \frac{d^3 k}{\omega_k} \frac{1}{2\pi i} \int_{\alpha}^{\beta} d\mu h_{\pm}(k, s, \mu, \epsilon) \end{aligned} \quad (4.16)$$

where

$$\begin{aligned} h_{\pm}(k, s, \mu, \epsilon) &= (\pm\omega_k - (\mu+i\epsilon))^{-1} g_{\pm}(k, s, \mu+i\epsilon) \\ &\quad - (\pm\omega_k - (\mu-i\epsilon))^{-1} g_{\pm}(k, s, \mu-i\epsilon) \end{aligned} \quad (4.17)$$

Since $g_{\pm}$ is analytic in z Taylor's formula can be used and together

with Lebesgue's dominated convergence theorem equation (4.16)

becomes

$$\begin{aligned} & \lim_{\epsilon \searrow 0} \frac{1}{2\pi i} \int_{\alpha}^{\beta} d\mu \langle f, [R(\mu+i\epsilon) - R(\mu-i\epsilon)] f \rangle_{\mathcal{H}} \\ &= \sum_{\pm} \sum_{s=1,2} \int_{\mathbb{R}^3} \frac{d^3 k}{\omega_k} : \pi^{-1} \lim_{\epsilon \searrow 0} \int_{\alpha}^{\beta} d\mu \Psi_{\pm}(k, s, \mu, \epsilon) \frac{\epsilon}{(\mu \mp \omega_k)^2 + \epsilon^2} \end{aligned} \quad (4.18)$$

where

$$\begin{aligned}
 \Psi_{\pm}(k,s,\mu,\epsilon) = & g_{\pm}(k,s,\mu) - g_{\pm}'(k,s,\mu)(\mu \mp \omega_k) \\
 & - \epsilon(1/2[\operatorname{Re}(g_{\pm}''(\xi_{\pm})) + \operatorname{Im}(g_{\pm}''(\eta_{\pm}))](\pm \omega_k - (\mu - i\epsilon)) \\
 & - 1/2[\operatorname{Re}(g_{\pm}''(\xi_{\pm}')) + \operatorname{Im}(g_{\pm}''(\eta_{\pm}'))](\pm \omega_k - (\mu - i\epsilon)))
 \end{aligned} \tag{4.19}$$

here $g_{\pm}'$ denote the partial derivative of $g_{\pm}$, $g_{\pm}''$ the second partial derivative of $g_{\pm}$ in the z -variable and $0 \leq \xi_{\pm}, \xi_{\pm}', \eta_{\pm}, \eta_{\pm}' \leq \epsilon$.

Finally since $f_{\epsilon}(y) = \pi^{-1}\epsilon(y^2 + \epsilon^2)^{-1}$ is a delta family (as $\epsilon \searrow 0$) equation (4.18) gives

$$\begin{aligned}
 \lim_{\epsilon \searrow 0} \frac{1}{2\pi i} \int_{\alpha}^{\beta} d\mu \langle f, [R(\mu + i\epsilon) - R(\mu - i\epsilon)] f \rangle_{\mathcal{H}} \\
 = \sum_{\pm} \sum_{s=1,2} \int_{\alpha < \pm \omega_k < \beta} \frac{d^3 k}{\omega_k} \Psi_{\pm}(k,s,\pm \omega_k, 0)
 \end{aligned} \tag{4.20}$$

i.e.

$$\lim_{\epsilon \searrow 0} \frac{1}{2\pi i} \int_{\alpha}^{\beta} d\bar{\mu} \langle f, [R(\bar{\mu} + i\epsilon) - R(\bar{\mu} - i\epsilon)] f \rangle_{\mathcal{H}}$$

$$= \left\{ \begin{array}{ll} \sum_{s=1,2} \int_{\alpha < \omega_k < \beta} \frac{d^3 k}{\omega_k} \quad | \hat{f}_+(k,s) |^2, & m < \alpha < \beta \\ \sum_{s=1,2} \int_{-\beta < \omega_k < -\alpha} \frac{d^3 k}{\omega_k} \quad | \hat{f}_-(k,s) |^2, & \alpha < \beta < -m \\ \sum_{s=1} \int_{\omega_k < \beta} \frac{d^3 k}{\omega_k} \quad | \hat{f}_+(k,s) |^2 \\ + \sum_{s=1,2} \int_{\omega_k < -\alpha} \frac{d^3 k}{\omega_k} \quad | \hat{f}_-(k,s) |^2, & \alpha < -m, \beta > m \\ 0 & , \text{ otherwise} \end{array} \right. \quad (4.21)$$

duing to the fact that $\Psi_{\pm}(k,s,\pm\omega_k,0) = g_{\pm}(k,s,\pm\omega_k) = | \hat{f}_{\pm}(k,s) |^2$.
From equation (4.1) and (4.17) it then follows that

$$\langle f, [E(\beta) - E(\alpha)]f \rangle_{\mathcal{H}}$$

$$= \begin{cases} \sum_{s=1,2} \int_{\alpha < \omega_k < \beta} \frac{d^3k}{\omega_k} |\hat{f}_+(k,s)|^2, & m < \alpha < \beta \\ \sum_{s=1,2} \int_{-\beta < \omega_k < -\alpha} \frac{d^3k}{\omega_k} |\hat{f}_-(k,s)|^2, & \alpha < \beta < -m \\ \sum_{s=1,2} \int_{\omega_k < \beta} \frac{d^3k}{\omega_k} |\hat{f}_+(k,s)|^2 \\ + \sum_{s=1,2} \int_{\omega_k < -\alpha} \frac{d^3k}{\omega_k} |\hat{f}_-(k,s)|^2, & \alpha < -m, \beta > m \\ 0 & , \text{otherwise} \end{cases} \quad (4.22)$$

Letting $\alpha \rightarrow -\infty$ and $\beta \rightarrow \infty$ formular (4.23) gives

$$\|f\|_{\mathcal{H}}^2 = \sum_{\pm} \sum_{s=1,2} \int_{\mathbb{R}^3} \frac{d^3k}{\omega_k} |\hat{f}_{\pm}(k,s)|^2 \quad (4.24)$$

for $f \in C_0^\infty(\mathbb{R}^3, \mathbb{C}^4)$. Now the map defined in definition 4.3 can be extended by continuity to an isometry.

Definition 4.4. Let $\widehat{\mathcal{H}} = \widehat{\mathcal{H}}_+ \oplus \widehat{\mathcal{H}}_-$, $\widehat{\mathcal{H}}_\pm = L^2(\mathfrak{M}_\pm, \mathbb{C}^2)$ where $\mathfrak{M}_\pm$ are the mass-hyperboloids and the measure is $\frac{d^3k}{\omega_k}$. Define

$$F : \mathcal{H} \rightarrow \widehat{\mathcal{H}} \text{ by } Ff = (\widehat{f}_+, \widehat{f}_-).$$

It follows from formula (2.7) that the map $F : \mathcal{H} \rightarrow \widehat{\mathcal{H}}$ in fact is onto, i.e. F is unitary.

Theorem 4.5. The unitary map $F : \mathcal{H} \rightarrow \widehat{\mathcal{H}}$, called the "light-cone Fourier transform", diagonalizes H , i.e.

$$FHf = (\omega_k \widehat{f}_+ , -\omega_k \widehat{f}_-) \quad (4.25)$$

for $f \in \mathcal{D}(H)$.

Proof. Let $f \in C_0^\infty(\mathbb{R}^3, \mathbb{C}^4)$ then

$$\begin{aligned} \widehat{Hf}_\pm(k,s) &= \langle u_\pm(\cdot, k, s), Hf(\cdot) \rangle_{\mathcal{H}} \\ &= \langle Du_\pm(\cdot, k, s), f(\cdot) \rangle_{L^2} \\ &= \pm \omega_k \langle u(\cdot, k, s), Qf(\cdot) \rangle_{L^2} \\ &= \pm \omega_k \widehat{f}_\pm(k, s) \end{aligned} \quad (4.26)$$

where $Du_\pm$ is understood in weak sense. Since H is closed and F is continuous, formula (4.25) holds for all $f \in \mathcal{D}(H)$.

The light-cone Fourier transform defines a kind of "duality" between the light-cone and the mass-hyperboloids.

Recall that formula (2.3) gives rise to a unitary map from $\hat{\mathcal{H}}$ onto $L^2(\mathbb{R}^3, \mathbb{C}^4)$ of $t = \text{constant}$ data, i.e. there is a one-to-one correspondance between weak solutions $u(t, x)$ of the Dirac equation (2.1) with $u(t, \cdot) \in L^2(\mathbb{R}^3, \mathbb{C}^4)$ and the solution $e^{-iHt}f$, $f \in \mathcal{H}$, of equation (3.13).

5. Conclusion.

We have shown that the characteristic Cauchy problem for the Dirac equation

$$(i\gamma^0 \frac{\partial}{\partial t} + i\gamma \cdot \nabla - m\mathbf{1})u(t,x) = 0$$

$$u(-|x|, x) = f(x) \quad , \quad x \in \mathbb{R}^3 \quad (5.1)$$

has a unique weak solution with $u(t, \cdot) \in L^2(\mathbb{R}^3, \mathbb{C}^4)$ for all $f \in \mathcal{H}$ and that all "L²-solutions" are obtained in this way.

The Dirac equation was written as an evolution equation for light-cone data

$$i \frac{du}{dt} = Hu \quad (5.2)$$

and the spectral theory for H was developed including a "light-cone Fourier transform" which diagonalized H .

I want to acknowledge Department of mathematics, University of Copenhagen, Denmark where this letter is done and Department of mathematics, University of California at Santa Barbara where the final form has taken place. Especialy I want to thank Lars-Erik Lundberg Department of mathematics, University of Copenhagen, Denmark for suggesting the problem and for giving me lots of advice.

References.

- [1] Lundberg, L.-E. : The Klein-Gordon Equation with
Light-Cone Data. Comm. Math. Phys. 62 , 107 (1978).
- [2] Riesz, m.: L'intégrale de Riemann-Lionville et le problem de
Cauchy. Acta Math. 81, 1 (1949).
- [3] Strichartz, R. S.: Harmonic analysis on hyperboloids. J. Funct.
Anal. 12, 341 (1973).
- [4] Hörmander, L.: Linear partial differential operators, p.151.
Berlin, Göttingen, Heidelberg, New York: Springer 1964.

- 1/78 "TANKER OM EN PRAKSIS" - et matematikprojekt.
Projektrapport af: Anne Jensen, Lena Lindenskov, Marianne Kesselhahn og Nicolai Lomholt.
Vejleder: Anders Madsen
- 2/78 "OPTIMERING" - Menneskets forøgede beherskelsermuligheder af natur og samfund.
Projektrapport af: Tom J. Andersen, Tommy R. Andersen, Gert Krenøe og Peter H. Lassen
Vejleder: Bernhelm Boss.
- 3/78 "OPCAVESAMLING", breddekursus i fysik.
Af: Lasse Rasmussen, Aage Bonde Kræmmer og Jens Højgaard Jensen.
- 4/78 "TRE ESSAYS" - om matematikundervisning, matematiklæreruddannelsen og videnskabsrindalismen.
Af: Mogens Niss
Nr. 4 er p.t. udgået.
- 5/78 "BIBLIOGRAFISK VEJLEDNING til studiet af DEN MODERNE FYSIKS HISTORIE".
Af: Helge Kragh.
Nr. 5 er p.t. udgået.
- 6/78 "NOGLE ARTIKLER OG DEBATINDLÆG OM - læreruddannelse og undervisning i fysik, og - de naturvidenskabelige fags situation efter studenteroprøret".
Af: Karin Beyer, Jens Højgaard Jensen og Bent C. Jørgensen.
- 7/78 "MATEMATIKKENS FORHOLD TIL SAMFUNDSØKONOMIEN".
Af: B.V. Gnedenko.
Nr. 7 er udgået.
- 8/78 "DYNAMIK OG DIAGRAMMER". Introduktion til energy-bond-graph formalismen.
Af: Peder Voetmann Christiansen.
- 9/78 "OM PRAKSIS' INDFLYDELSE PÅ MATEMATIKKENS UDVIKLING". - Motiver til Kepler's: "Nova Stereometria Doliorum Vinariorum".
Projektrapport af: Lasse Rasmussen.
Vejleder: Anders Madsen.
-
- 10/79 "TERMODYNAMIK I GYMNASIET".
Projektrapport af: Jan Christensen og Jeanne Mortensen.
Vejledere: Karin Beyer og Peder Voetmann Christiansen.
- 11/79 "STATISTISKE MATERIALER".
Af: Jørgen Larsen.
- 12/79 "LINEÆRE DIFFERENTIALLIGNINGER OG DIFFERENTIALLIGNINGSSYSTEMER".
Af: Mogens Brun Heefelt.
Nr. 12 er udgået.
- 13/79 "CAVENDISH'S FORSØG I GYMNASIET".
Projektrapport af: Gert Kreinøe.
Vejleder: Albert Chr. Paulsen.
- 14/79 "BOOKS ABOUT MATHEMATICS: History, Philosophy, Education, Models, System Theory, and Works of".
Af: Else Høyrup.
Nr. 14 er p.t. udgået.
- 15/79 "STRUKTUREL STABILITET OG KATASTROFER i systemer i og udenfor termodynamisk ligevægt".
Specialeopgave af: Leif S. Striegler.
Vejleder: Peder Voetmann Kristiansen.
- 16/79 "STATISTIK I KRÆFTFORSKNINGEN".
Projektrapport af: Michael Olsen og Jørn Jensen.
Vejleder: Jørgen Larsen.
- 17/79 "AT SPØRGE OG AT SVARE i fysikundervisningen".
Af: Albert Christian Paulsen.
- 18/79 "MATHEMATICS AND THE REAL WORLD", Proceedings af an International Workshop, Roskilde University Centre, Denmark, 1978.
Preprint.
Af: Bernhelm Booss og Mogens Niss (eds.)
- 19/79 "GEOMETRI, SKOLE OG VIRKELIGHED".
Projektrapport af: Tom J. Andersen, Tommy R. Andersen og Per H.H. Larsen.
Vejleder: Mogens Niss.
- 20/79 "STATISTISKE MODELLER TIL BESTEMMELSE AF SIKRE DOSER FOR CARCINOGENE STOFFER".
Projektrapport af: Michael Olsen og Jørn Jensen.
Vejleder: Jørgen Larsen
- 21/79 "KONTROL I GYMNASIET-FORMÅL OG KONSEKVENSER".
Projektrapport af: Crilles Bacher, Per S. Jensen, Preben Jensen og Torben Nysteen.
- 22/79 "SEMIOTIK OG SYSTEMEGENSKABER (1)".
1-port lineært response og støj i fysikken.
Af: Peder Voetmann Christiansen.
- 23/79 "ON THE HISTORY OF EARLY WAVE MECHANICS - with special emphasis on the role of reality".
Af: Helge Kragh.
-
- 24/80 "MATEMATIKOPFATTELSE HOS 2. G'ERE".
a+b 1. En analyse. 2. Interviewmateriale.
Projektrapport af: Jan Christensen og Knud Lindhardt Rasmussen.
Vejleder: Mogens Niss.
- 25/80 "EKSAMENSOPGAVER", Dybdemodul/fysik 1974-79.
- 26/80 "OM MATEMATISKE MODELLER".
En projektrapport og to artikler.
Af: Jens Højgaard Jensen m.fl.
- 27/80 "METHODODOLOGY AND PHILOSOPHY OF SCIENCE IN PAUL DIRAC'S PHYSICS".
Af: Helge Kragh.
- 28/80 "DIELEKTRISK RELAXATION - et forslag til en ny model bygget på væskernes viscoelastiske egenskaber".
Projektrapport af: Gert Kreinøe.
Vejleder: Niels Boye Olsen.
- 29/80 "ODIN - undervisningsmateriale til et kursus i differentialligningsmodeller".
Projektrapport af: Tommy R. Andersen, Per H.H. Larsen og Peter H. Lassen.
Vejleder: Mogens Brun Heefelt.
- 30/80 "FUSIONSENERGIEN - - - ATOMSAMFUNDETS ENDESTATION".
Af: Oluf Danielsen.
Nr. 30 er udgået.
- 31/80 "VIDENSKABSTEORETISKE PROBLEMER VED UNDERVISNINGSSYSTEMER BASERET PÅ MÆNDELÆRE".
Projektrapport af: Troels Lange og Jørgen Karrebæk.
Vejleder: Stig Andur Pedersen.
Nr. 31 er p.t. udgået.
- 32/80 "POLYMERE STOFFERS VISCOELASTISKE EGENSKABER - BELYST VED HJÆLP AF MEKANISKE IMPEDANSMÅLINGER - GER MOSSBAUEREFFEKTMALINGER".
Projektrapport af: Crilles Bacher og Preben Jensen.
Vejledere: Niels Boye Olsen og Peder Voetmann Christiansen.
- 33/80 "KONSTITUERING AF FAG INDEN FOR TEKNISK - NATURVIDENSKABELIGE UDDANNELSER. I-II".
Af: Arne Jakobsen.
- 34/80 "ENVIRONMENTAL IMPACT OF WIND ENERGY UTILIZATION".
ENERGY SERIES NO. I.
Af: Bent Sørensen
Nr. 34 er udgået.

- 35/80 "HISTORISKE STUDIER I DEN NYERE ATOMFYSIKS UDVIKLING".
Af: Helge Kragh.
- 36/80 "HVAD ER MENINGEN MED MATEMATIKUNDERVISNINGEN?".
Fire artikler.
Af: Mogens Niss.
- 37/80 "RENEWABLE ENERGY AND ENERGY STORAGE".
ENERGY SERIES NO. 2.
Af: Bent Sørensen.
-
- 38/81 "TIL EN HISTORIETEORI OM NATURERKENDELSE, TEKNOLOGI OG SAMFUND".
Projektrapport af: Erik Gade, Hans Hedal, Henrik Lau og Finn Physant.
Vejledere: Stig Andur Pedersen, Helge Kragh og Ib Thiersen.
Nr. 38 er p.t. udgået.
- 39/81 "TIL KRITIKKEN AF VEKSTØKONOMIEN".
Af: Jens Højgaard Jensen.
- 40/81 "TELEKOMMUNIKATION I DANMARK - oplæg til en teknologivurdering".
Projektrapport af: Arne Jørgensen, Bruno Petersen og Jan Vedde.
Vejleder: Per Nørgaard.
- 41/81 "PLANNING AND POLICY CONSIDERATIONS RELATED TO THE INTRODUCTION OF RENEWABLE ENERGY SOURCES INTO ENERGY SUPPLY SYSTEMS".
ENERGY SERIES NO. 3.
Af: Bent Sørensen.
- 42/81 "VIDENSKAB TEORI SAMFUND - En introduktion til materialistiske videnskabsopfattelser".
Af: Helge Kragh og Stig Andur Pedersen.
- 43/81 1. "COMPARATIVE RISK ASSESSMENT OF TOTAL ENERGY SYSTEMS".
2. "ADVANTAGES AND DISADVANTAGES OF DECENTRALIZATION".
ENERGY SERIES NO. 4.
Af: Bent Sørensen.
- 44/81 "HISTORISKE UNDERSØGELSER AF DE EKSPERIMENTELLE FORUDSÆTNINGER FOR RUTHERFORDS ATOMMODEL".
Projektrapport af: Niels Thor Nielsen.
Vejleder: Bent C. Jørgensen.
-
- 45/82 Er aldrig udkommet.
- 46/82 "EKSEMPLARISK UNDERVISNING OG FYSISK ERKENDELSE-
1+11 ILLUSTRERET VED TO EKSEMPLER".
Projektrapport af: Torben O. Olsen, Lasse Rasmussen og Niels Dreyer Sørensen.
Vejleder: Bent C. Jørgensen.
- 47/82 "BARSEBÄCK OG DET VÆRST OFFICIELT-TÆNKELIGE UHELD".
ENERGY SERIES NO. 5.
Af: Bent Sørensen.
- 48/82 "EN UNDERSØGELSE AF MATEMATIKUNDERVISNINGEN PÅ ADGANGSKURSUS TIL KØBENHAVNS TEKNIKUM".
Projektrapport af: Lis Eilertzen, Jørgen Karrebæk, Troels Lange, Preben Nørregaard, Lissi Pedersen, Laust Rishøj, Lill Røn og Isac Showiki.
Vejledere: Mogens Niss.
- 49/82 "ANALYSE AF MULTISPEKTRALE SATELLITBILLEDER".
Projektrapport af: Preben Nørregaard.
Vejledere: Jørgen Larsen og Rasmus Ole Rasmussen.
- 50/82 "HERSLEV - MULIGHEDER FOR VEDVARENDE ENERGI I EN LANDSBY".
ENERGY SERIES NO. 6.
Rapport af: Bent Christensen, Bent Hove Jensen, Dennis B. Møller, Bjarne Laursen, Bjarne Lillethorup og Jacob Mørch Pedersen.
Vejleder: Bent Sørensen.
- 51/82 "HVAD KAN DER GÅR'S FOR AT AFHJÆLPE PICERS BLOKERING OVERFOR MATEMATIK?".
Projektrapport af: Lis Eilertzen, Lissi Pedersen, Lill Røn og Susanne Stender.
- 52/82 "DESUSPENSION OF SPLITTING ELLIPTIC SYMBOLS".
Af: Bernhelm Booss og Krzysztof Wojciechowski.
- 53/82 "THE CONSTITUTION OF SUBJECTS IN ENGINEERING EDUCATION".
Af: Arne Jacobsen og Stig Andur Pedersen.
- 54/82 "FUTURES RESEARCH" - A Philosophical Analysis of Its Subject-Matter and Methods.
Af: Stig Andur Pedersen og Johannes Witt-Hansen.
- 55/82 "MATEMATISKE MODELLER" - Litteratur på Roskilde Universitetsbibliotek.
En biografi.
Af: Else Højrup.

Vedr. tekst nr. 55/82 se også tekst nr. 62/83.
- 56/82 "EN - TO - MANGE" -
En undersøgelse af matematisk økologi.
Projektrapport af: Troels Lange.
Vejleder: Anders Madsen.
-
- 57/83 "ASPECT EKSPERIMENTET"-
Skjulte variable i kvantemekanikken?
Projektrapport af: Tom Juul Andersen.
Vejleder: Peder Voetmann Christiansen.
Nr. 57 er udgået.
- 58/83 "MATEMATISKE VANDRINGER" - Modelbetragtninger over spredning af dyr mellem småbiotoper i agerlandet.
Projektrapport af: Per Hammershøj Jensen og Lene Vagn Rasmussen.
Vejleder: Jørgen Larsen.
- 59/83 "THE METHODOLOGY OF ENERGY PLANNING".
ENERGY SERIES NO. 7.
Af: Bent Sørensen.
- 60/83 "MATEMATISK MODEKSPERTISE"- et eksempel.
Projektrapport af: Erik O. Gade, Jørgen Karrebæk og Preben Nørregaard.
Vejleder: Anders Madsen.
- 61/83 "FYSIKS IDEOLOGISKE FUNKTION, SOM ET EKSEMPEL PÅ EN NATURVIDENSKAB - HISTORISK SET".
Projektrapport af: Annette Post Nielsen.
Vejledere: Jens Højrup, Jens Højgaard Jensen og Jørgen Vogelius.
- 62/83 "MATEMATISKE MODELLER" - Litteratur på Roskilde Universitetsbibliotek.
En biografi 2. rev. udgave.
Af: Else Højrup.
- 63/83 "CREATING ENERGY FUTURES: A SHORT GUIDE TO ENERGY PLANNING".
ENERGY SERIES No. 8.
Af: David Crossley og Bent Sørensen.
- 64/83 "VON MATEMATIK UND KRIEG".
Af: Bernhelm Booss og Jens Højrup.
- 65/83 "ANVENDT MATEMATIK - TEORI ELLER PRAKSIS".
Projektrapport af: Per Hedegård Andersen, Kirsten Habekost, Carsten Holst-Jensen, Annelise von Moos, Else Marie Pedersen og Erling Møller Pedersen.
Vejledere: Bernhelm Booss og Klaus Grünbaum.
- 66/83 "MATEMATISKE MODELLER FOR PERIODISK SELEKTION I ESCHERICHIA COLI".
Projektrapport af: Hanne Lisbet Andersen, Ole Richard Jensen og Klavs Frisdaal.
Vejledere: Jørgen Larsen og Anders Hede Madsen.
- 67/83 "ELEPSOIDE METODEN - EN NY METODE TIL LINEÆR PROGRAMMERING?".
Projektrapport af: Lone Billmann og Lars Boye.
Vejleder: Mogens Brun Heefelt.
- 68/83 "STOKASTISKE MODELLER I POPULATIONSGENETIK" - til kritikken af teoriladede modeller.
Projektrapport af: Lise Odgård Gade, Susanne Hansen, Michael Hviid og Frank Mølgård Olsen.
Vejleder: Jørgen Larsen.

- 69/83 "ELEVFORUDSÆTNINGER I FYSIK"
- en test i l.g med kommentarer.
Af: Albert C. Paulsen.
- 70/83 "INDLÆRINGS - OG FORMIDLINGSPROBLEMER I MATEMATIK PÅ VOKSENUNDERVISNINGSNIVEAU".
Projektrapport af: Hanne Lisbet Andersen, Torben J. Andreasen, Svend Åge Houmann, Helle Glerup Jensen, Keld Fl. Nielsen, Lene Vagn Rasmussen.
Vejleder: Klaus Grünbaum og Anders Hede Madsen.
- 71/83 "PIGER OG FYSIK".
- et problem og en udfordring for skolen?
Af: Karin Beyer, Sussanne Blegaa, Birthe Olsen, Jette Reich og Mette Vedelsby.
- 72/83 "VERDEN IFØLGE PEIRCE" - to metafysiske essays, om og af C.S Peirce.
Af: Peder Voetmann Christiansen.
- 73/83 "EN ENERGIANALYSE AF LANDBRUG"
- økologisk contra traditionelt.
ENERGY SERIES NO. 9
Specialeopgave i fysik af: Bent Høve Jensen.
Vejleder: Bent Sørensen.
-
- 74/84 "MINIATURISERING AF MIKROELEKTRONIK" - om videnskabeliggjort teknologi og nytten af at lære fysik.
Projektrapport af: Bodil Harder og Linda Szkotak Jensen.
Vejledere: Jens Højgaard Jensen og Bent C. Jørgensen.
- 75/84 "MATEMATIKUNDERVISNINGEN I FREMTIDENS GYMNASIUM"
- Case: Lineær programmering.
Projektrapport af: Morten Blomhøj, Klavs Frisdahl og Frank Mølgaard Olsen.
Vejledere: Mogens Brun Heefelt og Jens Bjørneboe.
- 76/84 "KERNEKRAFT I DANMARK?" - Et høringssvar indkaldt af miljøministeriet, med kritik af miljøstyrelsens rapporter af 15. marts 1984.
ENERGY SERIES No. 10
Af: Niels Boye Olsen og Bent Sørensen.
- 77/84 "POLITISKE INDEKS - FUP ELLER FAKTA?"
Opinionsundersøgelser belyst ved statistiske modeller.
Projektrapport af: Svend Åge Houmann, Keld Nielsen og Susanne Stender.
Vejledere: Jørgen Larsen og Jens Bjørneboe.
- 78/84 "JÆVNSTRØMSLEDNINGSEVNE OG GITTERSTRUKTUR I AMORFT GERMANIUM".
Specialrapport af: Hans Hedal, Frank C. Ludvigsen og Finn C. Physant.
Vejleder: Niels Boye Olsen.
- 79/84 "MATEMATIK OG ALMENDANNELSE".
Projektrapport af: Henrik Ooster, Mikael Wennerberg Johansen, Povl Kattler, Birgitte Lydholm og Morten Overgaard Nielsen.
Vejleder: Bernhelm Booss.
- 80/84 "KURSUSMATERIALE TIL MATEMATIK B".
Af: Mogens Brun Heefelt.
- 81/84 "FREKVENSAFHÆNGIG LEDNINGSEVNE I AMORFT GERMANIUM".
Specialrapport af: Jørgen Wind Petersen og Jan Christensen.
Vejleder: Niels Boye Olsen.
- 82/84 "MATEMATIK - OG FYSIKUNDERVISNINGEN I DET AUTOMATISEREDE SAMFUND".
Rapport fra et seminar afholdt i Hvidovre 25-27 april 1983.
Red.: Jens Højgaard Jensen, Bent C. Jørgensen og Mogens Niss.
- 83/84 "ON THE QUANTIFICATION OF SECURITY":
PEACE RESEARCH SERIES NO. 1
Af: Bent Sørensen
nr. 83 er p.t. udgået
- 84/84 "NOGLE ARTIKLER OM MATEMATIK, FYSIK OG ALMENDANNELSE".
Af: Jens Højgaard Jensen, Mogens Niss m. fl.
- 85/84 "CENTRIFUGALREGULATORER OG MATEMATIK".
Specialrapport af: Per Hedegård Andersen, Carsten Holst-Jensen, Else Marie Pedersen og Erling Møller Pedersen.
Vejleder: Stig Andur Pedersen.
- 86/84 "SECURITY IMPLICATIONS OF ALTERNATIVE DEFENSE OPTIONS FOR WESTERN EUROPE".
PEACE RESEARCH SERIES NO. 2
Af: Bent Sørensen.
- 87/84 "A SIMPLE MODEL OF AC HOPPING CONDUCTIVITY IN DISORDERED SOLIDS".
Af: Jeppe C. Dyre.
- 88/84 "RISE, FALL AND RESURRECTION OF INFINITESIMALS".
Af: Detlef Laugwitz.
- 89/84 "FJERNVARMEOPTIMERING".
Af: Bjarne Lillethorup og Jacob Mørch Pedersen.
- 90/84 "ENERGI I L.G - EN TEORI FOR TILRETTELÆGGELSE".
Af: Albert Chr. Paulsen.
-
- 91/85 "KVANTETEORI FOR GYMNASIET".
1. Lærervejledning
Projektrapport af: Biger Lundgren, Henning Sten Hansen og John Johansson.
Vejleder: Torsten Meyer.
- 92/85 "KVANTETEORI FOR GYMNASIET".
2. Materiale
Projektrapport af: Biger Lundgren, Henning Sten Hansen og John Johansson.
Vejleder: Torsten Meyer.
- 93/85 "THE SEMIOTICS OF QUANTUM - NON - LOCALITY".
Af: Peder Voetmann Christiansen.
- 94/85 "TREENIGHEDEN BOURBAKI - generalen, matematikeren og ånden".
Projektrapport af: Morten Blomhøj, Klavs Frisdahl og Frank M. Olsen.
Vejleder: Mogens Niss.
- 95/85 "AN ALTERNATIV DEFENSE PLAN FOR WESTERN EUROPE".
PEACE RESEARCH SERIES NO. 3
Af: Bent Sørensen
- 96/85 "ASPEKTER VED KRAFTVARMEFORSYNING".
Af: Bjarne Lillethorup.
Vejleder: Bent Sørensen.
- 97/85 "ON THE PHYSICS OF A.C. HOPPING CONDUCTIVITY".
Af: Jeppe C. Dyre.
- 98/85 "VALGMULIGHEDER I INFORMATIONSSALDEREN".
Af: Bent Sørensen.
- 99/85 "Der er langt fra Q til R".
Projektrapport af: Niels Jørgensen og Mikael Klintorp.
Vejleder: Stig Andur Pedersen.
- 100/85 "TALSYSTEMETS OPBYGNING".
Af: Mogens Niss.
- 101/85 "EXTENDED MOMENTUM THEORY FOR WINDMILLS IN PERTURBATIVE FORM".
Af: Ganesh Sengupta.
- 102/85 OPSTILLING OG ANALYSE AF MATEMATISKE MODELLER, BELYST VED MODELLER OVER KØRS FODEROPTAGELSE OG - OMSÆTNING".
Projektrapport af: Lis Eilertzen, Kirsten Habekost, Lill Røn og Susanne Stender.
Vejleder: Klaus Grünbaum.

- 103/85 "ØDSLE KOLDKRIGERE OG VIDENSKABENS LYSE IDEER".
Projektrapport af: Niels Ole Dam og Kurt Jensen.
Vejleder: Bent Sørensen.
- 104/85 "ANALOGREGNEMASKINEN OG LORENZLIGNINGER".
Af: Jens Jæger.
- 105/85 "THE FREQUENCY DEPENDENCE OF THE SPECIFIC HEAT OF THE GLASS TRANSITION".
Af: Tage Christensen.

"A SIMPLE MODEL OF AC HOPPING CONDUCTIVITY".
Af: Jeppe C. Dyre.
Contributions to the Third International Conference on the Structure of Non - Crystalline Materials held in Grenoble July 1985.
- 106/85 "QUANTUM THEORY OF EXTENDED PARTICLES".
Af: Bent Sørensen.
- 107/85 "EN MYG GØR INGEN EPIDEMI".
- flodblindhed som eksempel på matematisk modellering af et epidemiologisk problem.
Projektrapport af: Per Hedegård Andersen, Lars Boye, Carsten Holst Jensen, Else Marie Pedersen og Erling Møller Pedersen.
Vejleder: Jesper Larsen.
- 108/85 "APPLICATIONS AND MODELLING IN THE MATHEMATICS CURRICULUM" - state and trends -
Af: Mogens Niss.
- 109/85 "COX I STUDIETIDEN" - Cox's regressionsmodel anvendt på studenteroplysninger fra RUC.
Projektrapport af: Mikael Wennerberg Johansen, Poul Katler og Torben J. Andreassen.
Vejleder: Jørgen Larsen.
- 110/85 "PLANNING FOR SECURITY".
Af: Bent Sørensen.
- 111/85 "JORDEN RUNDT PÅ FLADE KORT".
Projektrapport af: Birgit Andresen, Beatriz Quinones og Jimmy Staal.
Vejleder: Mogens Niss.
- 112/85 "VIDENSKABELIGGØRELSE AF DANSK TEKNOLOGISK INNOVATION FRA 1950 - BELYST VED EKSEMPLER".
Projektrapport af: Erik Odgaard Gade, Hans Hedal, Frank C. Ludvigsen, Annette Post Nielsen og Finn Physant.
Vejleder: Claus Bryld og Bent C. Jørgensen.
- 113/85 "DESUSPENSION OF SPLITTING ELLIPTIC SYMBOLS II".
Af: Bernhelm Booss og Krzysztof Wojciechowski.
- 114/85 "ANVENDELSE AF GRAFISKE METODER TIL ANALYSE AF KONTINGENSTABELLER".
Projektrapport af: Lone Billmann, Ole R. Jensen og Arne-Lise von Moos.
Vejleder: Jørgen Larsen.
- 115/85 "MATEMATIKKENS UDVIKLING OP TIL RENESSANCEN".
Af: Mogens Niss.
- 116/85 "A PHENOMENOLOGICAL MODEL FOR THE MEYER-NELDEL RULE".
Af: Jeppe C. Dyre.
- 117/85 "KRAFT & FJERNVARMEOPTIMERING".
Af: Jacob Mørch Pedersen.
Vejleder: Bent Sørensen.
- 118/85 "TILFÆLDIGHEDEN OG NØDVENDIGHEDEN IFØLGE PEIRCE OG FYSIKKEN".
Af: Peder Voetmann Christiansen.
- 119/86 "DET ER GANSKE VIST - - EUKLIDS FEMTE POSTULAT KUNNE NOK SKABE RØRE I ANDEDAMMEN".
Af: Iben Maj Christiansen
Vejleder: Mogens Niss.
- 120/86 "ET ANTAL STATISTISKE STANDARDMODELLER".
Af: Jørgen Larsen.
- 121/86 "SIMULATION I KONTINUERT TID".
Af: Peder Voetmann Christiansen.
- 122/86 "ON THE MECHANISM OF GLASS IONIC CONDUCTIVITY".
Af: Jeppe C. Dyre.
- 123/86 "GYMNASIEFYSIKKEN OG DEN STORE VERDEN".
Fysiklærerforeningen, IMFUFA, RUC.
- 124/86 "OPGAVESAMLING I MATEMATIK".
Samtlige opgaver stillet i tiden 1974-jan. 1986.
- 125/86 "UVBY, 8 - systemet - en effektiv fotometrisk spektral-klassifikation af B-, A- og F-stjerner".
Projektrapport af: Birger Lundgren.
- 126/86 "OM UDVIKLINGEN AF DEN SPECIELLE RELATIVITETSTEORI".
Projektrapport af: Lise Odgaard & Linda Szkotak Jensen
Vejledere: Karin Beyer & Stig Andur Pedersen.
- 127/86 "GALOIS' BIDRAG TIL UDVIKLINGEN AF DEN ABSTRAKTE ALGEBRA".
Projektrapport af: Pernille Sand, Heine Larsen & Lars Frandsen.
Vejleder: Mogens Niss.
- 128/86 "SMÅKRYB" - om ikke-standard analyse.
Projektrapport af: Niels Jørgensen & Mikael Klintorp.
Vejleder: Jeppe Dyre.
- 129/86 "PHYSICS IN SOCIETY"
Lecture Notes 1983 (1986)
Af: Bent Sørensen
- 130/86 "Studies in Wind Power"
Af: Bent Sørensen
- 131/86 "FYSIK OG SAMFUND" - Et integreret fysik/historie-projekt om naturanskuelsens historiske udvikling og dens samfundsmæssige betingethed.
Projektrapport af: Jakob Heckscher, Søren Brønd, Andy Wierød.
Vejledere: Jens Høyrup, Jørgen Vogelius, Jens Højgaard Jensen.
- 132/86 "FYSIK OG DANNEELSE"
Projektrapport af: Søren Brønd, Andy Wierød.
Vejledere: Karin Beyer, Jørgen Vogelius.
- 133/86 "CHERNOBYL ACCIDENT: ASSESSING THE DATA. ENERGY SERIES NO. 15."
Af: Bent Sørensen.
-
- 134/87 "THE D.C. AND THE A.C. ELECTRICAL TRANSPORT IN AsSeTe SYSTEM"
Authors: M.B.El-Den, N.B.Olsen, Ib Høst Pedersen, Petr Visčor
- 135/87 "INTUITIONISTISK MATEMATIKS METODER OG ERKENDELSESTEORETISKE FORUDSÆTNINGER"
MATEMATIKSPECIALE: Claus Larsen
Vejledere: Anton Jensen og Stig Andur Pedersen
- 136/87 "Mystisk og naturlig filosofi: En skitse af kristendommens første og andet møde med græsk filosofi"
Projektrapport af Frank Colding Ludvigsen
Vejledere: Historie: Ib Thiersen
Fysik: Jens Højgaard Jensen
- 137/87 "HOPMODELLER FOR ELEKTRISK LEDNING I UORDNEDE FASTE STOFFER" - Resume af licentiatafhandling
Af: Jeppe Dyre
Vejledere: Niels Boye Olsen og Peder Voetmann Christiansen.

- 138/87 "JOSEPHSON EFFECT AND CIRCLE MAP."
Paper presented at The International Workshop on Teaching Nonlinear Phenomena at Universities and Schools, "Chaos in Education". Balaton, Hungary, 26 April-2 May 1987.
By: Peder Voetmann Christiansen
- 139/87 "Machbarkeit nichtbeherrschbarer Technik durch Fortschritte in der Erkennbarkeit der Natur"
Af: Bernhelm Booss-Bavnbek
Martin Bohle-Carbonell
- 140/87 "ON THE TOPOLOGY OF SPACES OF HOLOMORPHIC MAPS"
By: Jens Gravesen
- 141/87 "RADIOMETERS UDVIKLING AF BLODGASAPPARATUR - ET TEKNOLOGIHISTORISK PROJEKT"
Projektrapport af Finn C. Physant
Vejleder: Ib Thiersen
- 142/87 "The Calderón Projektor for Operators With Splitting Elliptic Symbols"
by: Bernhelm Booss-Bavnbek og
Krzysztof P. Wojciechowski
- 143/87 "Kursusmateriale til Matematik på NAT-BAS"
af: Mogens Brun Heefelt
- 144/87 "Context and Non-Locality - A Peircan Approach
Paper presented at the Symposium on the Foundations of Modern Physics The Copenhagen Interpretation 60 Years after the Como Lecture. Joensuu, Finland, 6 - 8 august 1987.
By: Peder Voetmann Christiansen
- 145/87 "AIMS AND SCOPE OF APPLICATIONS AND MODELLING IN MATHEMATICS CURRICULA"
Manuscript of a plenary lecture delivered at ICMTA 3, Kassel, FRG 8.-11.9.1987
By: Mogens Niss
- 146/87 "BESTEMMELSE AF BULKRESISTIVITETEN I SILICIUM"
- en ny frekvensbaseret målemetode.
Fysikspeciale af Jan Vedde
Vejledere: Niels Boye Olsen & Petr Višćor
- 147/87 "Rapport om BIS på NAT-BAS"
redigeret af: Mogens Brun Heefelt
- 148/87 "Naturvidenskabsundervisning med Samfundsperspektiv"
af: Peter Colding-Jørgensen DLH
Albert Chr. Paulsen
- 149/87 "In-Situ Measurements of the density of amorphous germanium prepared in ultra high vacuum"
by: Petr Višćor
- 150/87 "Structure and the Existence of the first sharp diffraction peak in amorphous germanium prepared in UHV and measured in-situ"
by: Petr Višćor
- 151/87 "DYNAMISK PROGRAMMERING"
Matematikprojekt af:
Birgit Andresen, Keld Nielsen og Jimmy Staal
Vejleder: Mogens Niss
- 152/87 "PSEUDO-DIFFERENTIAL PROJECTIONS AND THE TOPOLOGY OF CERTAIN SPACES OF ELLIPTIC BOUNDARY VALUE PROBLEMS"
by: Bernhelm Booss-Bavnbek
Krzysztof P. Wojciechowski
- 153/88 "HALVLEDERTEKNOLOGIENS UDVIKLING MELLEM MILITÆRE OG CIVILE KRÆFTER"
Et eksempel på humanistisk teknologihistorie
Historiespeciale
Af: Hans Heddal
Vejleder: Ib Thiersen
- 154/88 "MASTER EQUATION APPROACH TO VISCOUS LIQUIDS AND THE GLASS TRANSITION"
By: Jeppe Dyre
- 155/88 "A NOTE ON THE ACTION OF THE POISSON SOLUTION OPERATOR TO THE DIRICHLET PROBLEM FOR A FORMALLY SELFADJOINT DIFFERENTIAL OPERATOR"
by: Michael Pedersen
- 156/88 "THE RANDOM FREE ENERGY BARRIER MODEL FOR AC CONDUCTION IN DISORDERED SOLIDS"
by: Jeppe C. Dyre
- 157/88 "STABILIZATION OF PARTIAL DIFFERENTIAL EQUATIONS BY FINITE DIMENSIONAL BOUNDARY FEEDBACK CONTROL: A pseudo-differential approach."
by: Michael Pedersen
- 158/88 "UNIFIED FORMALISM FOR EXCESS CURRENT NOISE IN RANDOM WALK MODELS"
by: Jeppe Dyre
- 159/88 "STUDIES IN SOLAR ENERGY"
by: Bent Sørensen
- 160/88 "LOOP GROUPS AND INSTANTONS IN DIMENSION TWO"
by: Jens Gravesen
- 161/88 "PSEUDO-DIFFERENTIAL PERTURBATIONS AND STABILIZATION OF DISTRIBUTED PARAMETER SYSTEMS: Dirichlet feedback control problems"
by: Michael Pedersen
- 162/88 "PIGER & FYSIK - OG MEGET MERE"
Af: Karin Beyer, Sussanne Blegaa, Birthe Olsen, Jette Reich, Mette Vedelsby
- 163/88 "EN MATEMATISK MODEL TIL BESTEMMELSE AF PERMEABILITETEN FOR BLOD-NETHINDE-BARRIEREN"
Af: Finn Langberg, Michael Jarden, Lars Frellesen
Vejleder: Jesper Larsen
- 164/88 "Vurdering af matematisk teknologi
Technology Assessment
Technikfolgenabschätzung"
Af: Bernhelm Booss-Bavnbek, Glen Pate med
Martin Bohle-Carbonell og Jens Højgaard Jensen
- 165/88 "COMPLEX STRUCTURES IN THE NASH-MOSER CATEGORY"
by: Jens Gravesen

166/88 "Grundbegreber i Sandsynligheds-
regningen"

Af: Jørgen Larsen

167a/88 "BASISSTATISTIK 1. Diskrete modeller"

Af: Jørgen Larsen

167b/88 "BASISSTATISTIK 2. Kontinuerte
modeller"

Af: Jørgen Larsen

168/88 "OVERFLADEN AF PLANETEN MARS"

Laboratorie-simulering og MARS-analoger
undersøgt ved Mössbauerspektroskopi.

Fysikspeciale af:

Birger Lundgren

Vejleder: Jens Martin Knudsen
Fys.Lab./HCØ

169/88 "CHARLES S. PEIRCE: MURSTEN OG MØRTEL
TIL EN METAFYSIK."

Fem artikler fra tidsskriftet "The Monist"
1891-93.

Introduktion og oversættelse:

Peder Voetmann Christéansen

170/88 "OPGAVESAMLING I MATEMATIK"

Samtlige opgaver stillet i tiden
1974 - juni 1988