

The calderón projektor for operators with splitting elliptic symbols

Booss-Bavnbek, Bernhelm; Wojciechowski, Krzysztof P.

Publication date:
1987

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Booss-Bavnbek, B., & Wojciechowski, K. P. (1987). *The calderón projektor for operators with splitting elliptic symbols*. Roskilde Universitet.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

TEKST NR 142

1987

Bernhelm Booß-Bavnbek
Krzysztof P. Wojciechowski

The Calderón Projector
for Operators With
Splitting Elliptic Symbols

TEKSTER fra

IMFUFA

ROSKILDE UNIVERSITETSCENTER

INSTITUT FOR STUDIET AF MATEMATIK OG FYSIK SAMT DERES
FUNKTIONER I UNDERVISNING, FORSKNING OG ANVENDELSER

IMFUFA, Roskilde Universitetscenter. Postboks 260, DK-4000 Roskilde
Bernhelm Boøß-Bavnbeek. Krzysztof P. Wojciechowski
**THE CALDERÓN PROJECTOR FOR OPERATORS WITH SPLITTING
ELLIPTIC SYMBOL**

IMFUFAtekst nr. 142/1987. 16 sider. ISSN 0106-6242

Summary. We shall investigate a class of "splitting" elliptic non-classical pseudo-differential operators of symbol class $S^{1,0}$, i.e. elliptic operators which take the form $A = G_A(y, t)(\partial_t + B_t)$ near the boundary Y of a smooth oriented Riemannian compact manifold X , where t denotes the normal coordinate, B_t is an elliptic (classical) pseudo-differential operator of order 1 over Y , and G_A is a bundle isomorphism over Y . We shall show the existence of a precise pseudo-differential projection, the Calderón projector, onto the space of Cauchy data of the kernel of A over Y .

THE CALDERÓN PROJECTOR FOR OPERATORS WITH SPLITTING ELLIPTIC SYMBOLS

BERNHLM BOOSS-BAVNBEK, KRZYSZTOF P. WOJCIECHOWSKI

Summary.

We shall investigate a class of "splitting" elliptic non-classical pseudo-differential operators of symbol class $S^{1,0}$, i.e. elliptic operators which take the form $A = G_A(y, t)(\partial_t + B_t)$ near the boundary Y of a smooth oriented Riemannian compact manifold X , where t denotes the normal coordinate, B_t is an elliptic (classical) pseudo-differential operator of order 1 over Y , and G_A is a bundle isomorphism over Y . We shall show the existence of a precise pseudo-differential projection, the Calderón projector, onto the space of Cauchy data of the kernel of A over Y .

0 Introduction.

More than 20 years ago A.P. Calderón observed that certain subspaces of the kernel of an elliptic differential operator A over a compact manifold X with boundary Y admit a pseudo-differential projection (see the short announcement of Calderón [5] and his beautiful lectures [6]). The construction of the Calderón projector was worked out in detail in Seeley [14] (for the case of an invertible elliptic pseudo-differential operator which is equal to a differential operator near Y), and in Hörmander [9]; and it was later extended in Grubb [7] to differential systems elliptic in the sense of Douglis and Nirenberg. In Hörmander [11] the construction of Calderón was generalized to a class of pseudo-differential operators with what we call splitting elliptic symbols (using a symbol class from Grubb [8]). These works, however, approach the Calderón projector only on the symbol level, i.e. the construction leads only to an approximate projection (modulo smoothing operators).

In this note we shall show that one obtains a true pseudo-differential Calderón projector for a certain class of elliptic operators of first order with "splitting symbol" near the boundary Y . There are different methods of proving this fact. We can follow the argumentation of the Calderón Lectures; this was done earlier by the authors in [4] in connection with a systematic reexamination of elliptic boundary value problems using the machinery of "elliptic towers" taken from Birman-Solomyak [1].

The present construction uses some ideas of Palais [12] and the "doubling construction" (continuation of an elliptic symbol splitting near the boundary to the double) of [18] and [4] and gives a more direct approach to the Calderón projector.

To some extent, the ease of this construction and the vigour of the Calderón projector in treating elliptic boundary value problems allows us to dispense with the elaborate machinery of the Boutet de Monvel calculus (see [15], [2]). Moreover it turns out that our way of looking at the Calderón projector and at elliptic boundary value problems extends to elliptic problems on manifolds with singularities. This will be explained separately.

The plan of this paper is as follows: In Section 1 we introduce the class $\text{Ell}^1(X, Y)$ of operators with splitting elliptic symbols over a compact smooth oriented manifold X with smooth boundary Y . Section 2 contains two versions of our Continuation Lemma. The second leads immediately to the construction of the Calderón projector in Section 3. Section 4 presents some concluding remarks and a non-trivial example of an elliptic differential equation of second order which leads naturally to a (non-classical) pseudo-differential operator of first order with splitting elliptic symbol.

1 Operators With Splitting Elliptic Symbol.

Let X be a smooth compact oriented manifold with boundary Y . Let E and F be smooth complex vector bundles over X . We fix a Riemannian structure on X and a Hermitian structure on E and F . We use the Riemannian structure to construct a collar neighbourhood $N = Y \times I$ of the boundary Y , and we denote the (inward) normal coordinate by $t \in I = [0, 1]$. We identify Y with $Y \times \{0\}$ and $E|_N$ and $F|_N$ with the liftings of $E|_Y$ and $F|_Y$ to N . We denote the strip $Y \times [0, t]$ by X_t for $t \in I$.

1.1 DEFINITION. (a) A linear map $A : C^\infty(X; E) \rightarrow C^\infty(X; F)$ belongs to the class $\text{Pdiff}^1(X, Y)$ of (non-classical) pseudo-differential operators with *splitting symbol*, if it can be written in the form $A = A^b + A^i$ where A^i is a pseudo-differential operator of order 1 over X with distributional kernel $K^i(x, x')$ such that

$$\text{supp } K^i \subset (X \setminus X_{1/2}) \times (X \setminus X_{1/2}).$$

Moreover we shall assume that the operator A^b can be written in the form

$$A^b = \phi(t)G_A(y, t)(\partial_t + B_t),$$

where $B_t : C^\infty(Y; E|_Y) \rightarrow C^\infty(Y; F|_Y)$ is a pseudo-differential operator of first order and $G_A(\cdot, t) : E|_Y \rightarrow F|_Y$ is a bundle isomorphism for any $t \in I$, and where $\phi(t)$ is a smooth function of t vanishing for $t > 2/3$. If necessary, we can reparametrize the collar. Without loss of generality we may assume that $\phi(t) = 1$ for small t .

(b) In the following we assume that the families $\{G_A(\cdot, t)\}_{t \in I}$ and $\{B_t\}_{t \in I}$ depend smoothly on t . This assumption can be weakened somewhat, see Polking [13], but we will not go into the details in this paper.

Moreover, we will assume that the families extend to smooth families $\{B_t\}_{t \in [-\epsilon, 1]}$ and $\{G_A(\cdot, t)\}_{t \in [-\epsilon, 1]}$ for some $\epsilon > 0$. Extending $\{G_A(\cdot, t)\}$ is trivial. As shown in Seeley [15], a family $\{B_t\}_{t \in I}$ of differential operators can always be extended smoothly. However, the continuation of a family of pseudo-differential operators requires a more refined analysis which our assumption helps us avoid.

(c) The *principal symbol* σ_A of A is (by definition) the sum of the principal symbols of A^i and A^b . Close to Y it has the form

$$\sigma(x, \xi) = \phi(t)G_A(y, t)(i\tau + b(y, \eta)).$$

where $x = (y, t) \in X_{1/2}$, $y \in Y$, $0 \leq 1/2$, $\xi = (\eta, \tau) \in T^*(X)_x$, and $b(y, \eta)$ is the principal symbol of the operator B . Hence the *total symbol* $p(x, \xi)$ (which is the principal symbol + lower order terms) will locally satisfy estimates of the form

$$|D_\xi^\alpha D_x^\beta p(x, \xi)| \leq C_{\alpha\beta} (1 + |\xi|)^{m-\alpha''} (1 + |\eta|)^{m'-|\alpha'|}$$

with $m = 1$ and $m' = 0$, where α, β are multiindices, $\alpha = (\alpha_1, \dots, \alpha_n)$, $\alpha' = (\alpha_1, \dots, \alpha_{n-1})$, $|\alpha'| = \alpha_1 + \dots + \alpha_{n-1}$, and $\alpha'' = \alpha_n$. So our operator A has a symbol which belongs to the class $S^{m, m'}$, for $m = 1$ and $m' = 0$, which is defined in Grubb [8] and further investigated in Hörmander [11, Section 20.1, pp 242 ff].

(d) We say that A is *elliptic*, $A \in \text{Ell}^1(X, Y)$, if the principal symbol σ_A is an isomorphism.

(e) In the same way we define the operator classes $\text{Pdiff}^1(M, Y)$ and $\text{Ell}^1(M, Y)$ for any closed manifold M which contains Y as a submanifold of codimension 1 such that Y has a bicollar neighbourhood in M .

(f) Finally we introduce the classes $\text{Pdiff}^{-1}(X, Y)$ and $\text{Ell}^{-1}(X, Y)$ by assuming that the total symbol $p(x, \xi)$ belongs to the splitting symbol class $S^{-1, 0}$, i.e. $p(x, \xi)$ satisfies the inequality

$$|D_\xi^\alpha D_x^\beta p(x, \xi)| \leq C_{\alpha\beta} (1 + |\xi|)^{m-\alpha''} (1 + |\eta|)^{m'-|\alpha'|}$$

for $m = -1$ and $m' = 0$, and the principal symbol σ_A is invertible.

Note that all differential operators of first order belong to our class Pdiff^1 and that an operator A of class Pdiff^1 is a pseudo-differential operator of first order in the classical sense, i.e. it fulfills the symbol estimates

$$|D_\xi^\alpha D_x^\beta p(x, \xi)| \leq C_{\alpha\beta} (1 + |\xi|)^{1-|\alpha|}$$

if and only if A^b is a differential operator for a suitable reparametrization of the collar of Y .

We recall the following technical result without proof; it is a special case of Grubb [8, Théorème 2.1] and Hörmander [11, Proposition 20.1.11]:

12 THE CALCULUS OF SPLITTING ELLIPTIC SYMBOLS. Let M be a closed manifold with submanifold Y of codimension 1, let E and F be vector bundles over M , and let $A : C^\infty(M; E) \rightarrow C^\infty(M; F)$ belong to the class $\text{Ell}^1(M, Y)$. Then there exists a parametrix T of A which belongs to the class $\text{Ell}^{-1}(M, Y)$, i.e. an operator $T : C^\infty(M; F) \rightarrow C^\infty(M; E)$ with splitting elliptic symbol of order -1 such that $TA = Id - K_1$ and $AT = Id - K_2$ where $K_1, K_2 \in \text{Pdiff}^{-1}(M, Y)$.

Following the fundamental tenets of numerical analysis we build our analysis upon Green's Formula and the concept of the spaces of Cauchy data.

13 THEOREM (GREEN'S FORMULA). For any $A \in \text{Pdiff}^1(X, Y)$ with $A = G_A(y, t)(\partial t + B)$ near Y and for any $u \in C^\infty(X; E)$ and $v \in C^\infty(X; F)$ we have

$$\langle Au, v \rangle_F - \langle u, A^*v \rangle_E = - \int_Y (G_A(y, 0)u(y), v(y))_{F_y}$$

where A^* denotes the formal-adjoint of A and $\langle \dots \rangle_E$ and $\langle \dots \rangle_F$ denote the inner products in the section spaces defined by the Hermitian structure of the bundles and the volume form on X given by the Riemannian metric.

PROOF: We can just follow the arguments given in Palais [12, Theorem 17.1], Seeley [14, pp. 793 ff], or Hörmander [17, p. 183]. $\square$

1.4 DEFINITION. For $A \in \text{Ell}^1(X, Y)$ we define

(a) the space of *interior solutions* of the equation $Au = 0$ by

$$\ker_X A := \{u \in C^1(X; E) \mid Au = 0 \text{ and } u|_Y = 0\};$$

(b) the space of *Cauchy data* (along Y) by

$$H(A) := \{u|_Y \mid u \in C^\infty(X; E) \text{ and } Au = 0\}.$$

1.51 REMARK. "In general", namely when A satisfies the unique continuation property, the space of interior solutions consists only of the zero section, cf. Taylor [16, Chapter XIV].

1.52 REMARK. From Green's Formula it follows immediately that

$$H(A) \perp (G_A)^{-1}(H(A^*)).$$

Actually, the spaces are orthogonal complementary subspaces in $C^\infty(Y; E|_Y)$ as shown in by the authors in [4, Proposition 3.2] and - differently - in [19, Theorem 1].

2 "Symmetrization" and Continuation of Operators.

Operators with splitting elliptic symbols can be "symmetrized" near the boundary in the sense of the following Lemma. This was noted already in our [4, Lemma 3.1] from a topological perspective. Now we give an explicit deformation on the operator level:

2.1 LEMMA. Let X, Y, E , and F be as in Section 1 and let $A : C^\infty(X; E) \rightarrow C^\infty(X; F)$ belong to the class $\text{Ell}^1(X, Y)$ such that A takes the form $G_A(y, t)(\partial_t + B_t)$ on the collar $N = Y \times I$ where t denotes the (inward) normal coordinate, $B = \{B_t\}_{t \in I}$ is a family of elliptic pseudo-differential operators of first order $B_t : C^\infty(Y; E|_Y) \rightarrow C^\infty(Y; F|_Y)$, and $G_A(\cdot, t) : E|_Y \rightarrow F|_Y$ is a bundle isomorphism for any $t \in I$. As in 1.1(b) we assume that $\{B_t\}$ and $\{G_A(\cdot, t)\}$ depend smoothly on t and extend to small negative t . Then any such A extends to an $A' \in \text{Ell}^1(X', Y')$ such that A' takes the form $G_A(y, 0)(\partial_t + B')$ on the new collar $Y \times [-1, -1 + \epsilon]$ for some $\epsilon > 0$ where $B' : C^\infty(Y; E|_Y) \rightarrow C^\infty(Y; F|_Y)$ is an elliptic self-adjoint operator (which does not depend on t) and $X' = (X \setminus N) \cup (Y \times [-1, 1])$ denotes a larger manifold (diffeomorphic to X and with boundary $Y' = Y \times \{-1\}$; moreover we assume that E and F are defined on the whole of X').

PROOF: Since A is elliptic, the principal symbols $b_t(y, \eta)$ of B_t , $t \in I$, have no purely imaginary eigenvalues. Moreover, because of 1.1(b) we can extend this family

to a family $\{B_t\}_{t \in [-\epsilon, 1]}$ for some small $\epsilon > 0$ such that the principal symbols of B_t for negative t have no purely imaginary eigenvalues either. Now we further deform this family on the interval $[-\epsilon, -\epsilon/2]$ such that $B_t = B_{-\epsilon}$ in a neighbourhood of $-\epsilon$. We denote the principal symbol of $B_{-\epsilon}$ by b .

Now, let $y \in Y$, $(y, \eta) \in (T^*Y)_y$, $\eta \neq 0$, and let $p_+(y, \eta) : E_y \rightarrow E_y$ denote the projection onto the direct sum of the eigenspaces of the linear transformation $b(y, \eta) : E_y \rightarrow E_y$ corresponding to the eigenvalues with a positive real part. We may assume that p is a family of orthogonal projections. This can be obtained by a suitable deformation of the Hermitian structure of $E|_{Y \times [-\epsilon, -\epsilon/2]}$.

Then we have

$$b = (2p_+ - Id)^2 b = (2p_+ - Id) q_0$$

where $(2p_+ - Id)$ is an elliptic symbol of order 0 and $q_0 := (2p_+ - Id)b$ has the following property:

There exists a constant $C > 0$ such that

$$\Re(\langle q_0(y, \eta)v, v \rangle_{E_y}) \geq C\|v\|^2$$

for any $y \in Y$, $(y, \eta) \in S_y Y$ and $v \in E_y$, where $\Re$ denotes the real part.

Now let f be a smooth non-negative function on $\mathbb{R}$ equal to 0 in a neighbourhood of 0 and $f(t) = 1$ for $t > 1 - \delta$ with $\delta > 0$. Then the family of symbols

$$q_t(y, \eta) = f(t)f(|\eta|)|\eta| + (1 - f(t))q_0(y, \eta)$$

gives a smooth homotopy of elliptic symbols between $q_0 = (2p_+ - Id)b$ and $f(|\eta|)|\eta|$. Therefore we obtain a homotopy $r_t := (2p_+ - Id)q_t$ between b and the self-adjoint symbol $(2p_+ - Id)f(|\eta|)|\eta|$. We lift $\{r_t\}$ to a smooth homotopy $\{R_t\}$ of pseudo-differential operators (see e.g. Palais [12, Chapter XVI]) with $\sigma_L(R_t) = r_t$.

Clearly we may assume that $R_t = R_i$ in some neighbourhoods of $i \in \{0, 1\}$. We also may assume that $R_1^* = R_1$. In fact: We can replace $\{R_t\}$ near 1 (where it is the constant family equal to R_1) by a new family $\tilde{R}_t := (1 - f(t))R_1 + f(t)\frac{1}{2}(R_1 + R_1^*)$.

Now we make the deformation on the other end where $R_t = R_0$. Since $r_0 = b$, the operator $R_0 - B_{-\epsilon}$ is of order 0. So, we may connect R_0 and $B_{-\epsilon}$ by a family $\tilde{R}_t := (1 - f(t))B_{-\epsilon} + f(t)R_0$.

As a result of all these deformations we get (after reparametrization) a family of operators $\{\tilde{R}_t\}$ with principal symbols without purely imaginary eigenvalues which connects $B_{-\epsilon}$ with a self-adjoint operator $B' := R_1$. Moreover, this family is constant near $i = 0$, hence glueing is possible. So, after further reparametrization, we finally get a family which joins $B = \{B_t\}$ with the self-adjoint B' . This ends the proof. $\square$

The following Proposition provides the main tool for our approach to the Calderón projector.

22 PROPOSITION. *Let X, Y, N, E , and F be as in Section 1. Then any $A \in \text{Ell}^1(X, Y)$ extends to an $\tilde{A} \in \text{Ell}^1(\tilde{X}, Y)$ such that*

$$\tilde{A}|_{X \cup (Y \times [-1/4, 0])} = A|_{X \cup (Y \times [-1/4, 0])}.$$

Here $\tilde{X}$ denotes the closed double of X . Moreover, we assume that $X = (X \setminus N) \cup (Y \times [0, 1])$, that $X' = (X \setminus N) \cup (Y \times [-1, 1])$ is a larger manifold (diffeomorphic to X and with boundary $Y' = Y \times \{-1\}$), and that E, F and A are defined on the whole of X' .

PROOF: First, we apply Lemma 21 and deform A into an $A' \in \text{Ell}^1(X', Y')$ with

$$A'|_{X \cup (Y \times [-1/4, 0])} = A|_{X \cup (Y \times [-1/4, 0])}$$

and with a self-adjoint split summand B' near Y' . We denote the principal symbol of B' by b' .

Then we apply the glueing construction as it is described by the authors in [18] and [4]: Over the double $\tilde{X}' = (X')_+ \cup (X')_-$ with $(X')_+ = (X')_- = X'$ we have a well defined operator

$$\tilde{A} = A' \cup (A')^* := \begin{cases} A' & \text{on } (X')_+ \\ (A')^* & \text{on } (X')_- \end{cases}$$

acting from C^∞ sections of the bundle $E \cup_G F$ to C^∞ sections of the bundle $F \cup_{G^{-1}} E$ (with $G := (G_A)|_Y$). Note that $(A')^* = (-\partial_t + B')(G_A)^{-1}$ in a collar neighbourhood of $Y \times \{-1\}$, hence the operators A' and $(A')^*$ and the symbols

$$\sigma_{\tilde{A}}(-1, y; \tau, \eta) = \begin{cases} \sigma_{A'}(-1, y_+; \tau, \eta) = G_A(i\tau + b'(y_+, \eta)) \\ \sigma_{(A')^*}(-1, y_-; \tau, \eta) = (-i\tau + b'(y_-, \eta))(G_A)^{-1} \end{cases}$$

fit together under the identification $(-1, y_+; \tau, \eta) \sim (-1, y_-; -\tau, \eta)$ of the cotangent sphere bundles near $Y' = Y \times \{-1\}$ and yield an elliptic symbol $\sigma_{\tilde{A}} = \sigma_{A'} \cup \sigma_{(A')^*}$.

The operator $\tilde{A}$ thus becomes the operator we are looking for, if we identify $\tilde{X}'$, the double of X' , with $\tilde{X}$, the double of X , in such a way that $(X')_+$ goes over in X' and $(X')_- \setminus Y'$ goes over in $\tilde{X} \setminus X'$. $\square$

23 COROLLARY. Let A be an operator of class $\text{Ell}^1(X, Y)$ where X is a smooth oriented compact manifold with boundary Y . Then the space $\ker_X A$ is finite-dimensional and consists of smooth sections.

PROOF: By the preceding Proposition we can assume that A is the restriction of an elliptic operator $\tilde{A}$ on some closed manifold M containing X . We embed $\ker_X A$ in $\ker \tilde{A}$ by extending any interior solution over the whole of M by 0. Since the operators A and $\tilde{A}$ are of first order, we obtain a true solution of the homogeneous equation over M by the continuation (see Palais [12, Lemma 17.A]). The space $\ker \tilde{A}$ is finite-dimensional and consists of smooth sections. $\square$

241 REMARK. Note that

$$\begin{aligned} \ker(A' \cup A'^*) &= \{(u, v) \mid u \in C^\infty(X'; E), \quad v \in C^\infty(X'; F), \quad G_A(u|_{Y \times \{-1\}}) = \\ &= v|_{Y \times \{-1\}}, \text{ and } A'u = 0, \quad A'^*v = 0\} \end{aligned}$$

consists of the direct sum $\ker_{X'} A' \oplus \ker_{X'} A'^*$ of the spaces of interior solutions. Because of the mutual orthogonality of the spaces $H(A')$ and $(G_A)^{-1}(H(A'^*))$, see Remark 152, there exists no solution of the equation $(A' \cup A'^*)w = 0$ with $w|_Y$ not vanishing identically. Now it is also easy to determine the range $\text{ran}(A' \cup A'^*)$ and the index $\text{index}(A' \cup A'^*)$. Since $A' \cup A'^*$ has an elliptic symbol, the orthogonal complement of $\text{ran}(A' \cup A'^*)$ in $L^2(\tilde{X}'; F \cup_{G^{-1}} E)$ is identical with $\ker(A' \cup A'^*)^* \subset C^\infty(\tilde{X}'; F \cup_{G^{-1}} E)$. We have $(A' \cup A'^*)^* = A'^* \cup A'$, so

$$\text{coker}(A' \cup A'^*) \cong \ker_{X'} A'^* \oplus \ker_{X'} A' \stackrel{j}{\cong} \ker_{X'} A' \oplus \ker_{X'} A'^* \cong \ker(A' \cup A'^*),$$

hence $\text{index}(A' \cup A'^*) = 0$. Here j denotes the involution of the double of X' which interchanges the two factors $(X')_+$ and $(X')_-$.

242 REMARK. Our construction differs from the usual doubling construction (see e.g. Hörmander [11, Section 20.3, pp. 257 f]) which simply reflects the bundles and the operator along the boundary Y in the same way as the manifold. This results in a singularity along Y . The "supersymmetry" idea of looking at A and A^* simultaneously was suggested by A. P. Calderón and exploited already by Seeley in [14, Section 8] with his corrections given in [15, Appendix]. He considers the continuation of the matrix

$$\begin{pmatrix} 0 & A^* \\ A & 0 \end{pmatrix}$$

over the double through the identity (on the symbol level). Our construction is topologically different from Seeley's: (1) The operator $A \cup A^*$ can be defined on the operator level as a true continuation of A , and (2) it is not self-adjoint in general. (Although more explicit and more subtle than Seeley's construction, it does not provide additional information about the index because its index also vanishes.)

243 REMARK. If A satisfies the unique continuation property, then there exists no non-trivial interior solution and the continuation $A' \cup A'^*$ of A over the double is invertible because of Remark 241. In general, we get an invertible "approximative" continuation of A as a second Corollary to Proposition 22:

25 COROLLARY. Let X, Y, E, F be as above. For any $A \in \text{Ell}^1(X, Y)$ there exists a bijective operator $A'' \in \text{Ell}^1(\tilde{X}, Y)$ over the double $\tilde{X}$ with the following properties:

- (i) $A''|_X - A|_X$ is an operator of finite rank.
- (ii) The Cauchy data space $H_+(A'')$ of A'' along Y is equal to the Cauchy data space $H(A)$ of A along Y .
- (iii) The inverse of A'' belongs to the class $\text{Ell}^{-1}(\tilde{X}, Y)$.

PROOF: To (i). From Remark 241 we recall that $A' \cup A'^*$ is a Fredholm operator with $\text{index}(A' \cup A'^*) = 0$ and that the reflection j defines an isomorphism

$$\ker(A' \cup A'^*) \stackrel{j}{\cong} \ker_{X'} A' \oplus \ker_{X'} A'^* \cong \ker_{X'} A'^* \oplus \ker_{X'} A' \cong (\text{ran } A' \cup A'^*)^\perp.$$




Fig. 1. The decomposition of $\tilde{X}$

The standard method of getting an invertible operator from a Fredholm operator with vanishing index consists in identifying kernel and cokernel and then adding the orthogonal projection onto the kernel to the operator. This is very easy in our situation. We define $A'' := A' \cup A'^* + jk$, where $k : C^\infty(\tilde{X}; E \cup_G F) \rightarrow \ker_{X'} A'$ is the orthogonal projection. More explicitly, we write

$$A''w = A''(u, v) := (A'u + jk_2v, A'^*v + jk_1u),$$

where $w = (u, v)$, $u \in C^\infty(X'; E)$, $v \in C^\infty(X'; F)$ with $G_A(u|_{Y \times \{-1\}}) = v|_{Y \times \{-1\}}$, and k_1 and k_2 are the orthogonal projections of $C^\infty(X'; E)$ and $C^\infty(X'; F)$ onto the finite-dimensional subspaces $\ker_{X'} A'$ and $\ker_{X'} (A')^*$ of interior solutions. The difference jk between the operators A'' and $A' \cup A'^*$ is of finite rank and hence a smoothing operator. Therefore A'' belongs to $\text{Ell}^1(\tilde{X}, Y)$, too.

To (ii). Next we compare the Cauchy data spaces

$$H_+(A'') = \{w|_Y \mid w \in C^\infty(\tilde{X}; E \cup_G F) \text{ and } (A''w)|_X = 0\}$$

and

$$H(A) = \{u|_Y \mid u \in C^\infty(X; E) \text{ and } Au = 0\}.$$

Let us denote the operator $A' \cup A'^*$ by $\tilde{A}$. We split $C^\infty(\tilde{X}; E \cup_G F)$ into its two components $\ker \tilde{A}$ and $(\ker \tilde{A})^\perp$. Then we have

$$A''w = \tilde{A}w + jkw = \begin{cases} \tilde{A}w & \text{for } w \in (\ker \tilde{A})^\perp \\ jw & \text{for } w \in \ker \tilde{A}. \end{cases}$$

Hence the contributions to the Cauchy data spaces $H_+(A'')$ and $H(A) = H_+(\tilde{A})$ coming from sections in $(\ker \tilde{A})^\perp$ are identical, namely equal to

$$\{w|_Y \mid w \in (\ker \tilde{A})^\perp \text{ and } (\tilde{A}w)|_X = 0\}.$$

The contributions coming from $\ker \tilde{A}$ are also identical: The contribution to $H(A)$ consists of

$$\{w|_Y \mid w \in \ker \tilde{A}\} = \{w|_Y \mid w \in \ker_{(X')_+} \tilde{A}\}.$$

The contribution to $H_+(A'')$ consists of

$$\{w|_Y \mid w \in \ker \tilde{A} \text{ and } (jw)|_X = 0\} = \{w|_Y \mid w \in \ker_{\tilde{X} \setminus X_-} \tilde{A}\}.$$

Since $\ker \tilde{A}$ decomposes into the sum

$$\ker \tilde{A} = \ker_{(X')_+} \tilde{A} + \ker_{(X')_-} \tilde{A},$$

only elements in $\ker_{(X')_+} \tilde{A}$ contribute to $(\ker_{\tilde{X} \setminus X_-} \tilde{A})|_Y$. See Figures 1 and 2.




Fig. 2. The decomposition of $\ker_{\tilde{X} \setminus X_-} \tilde{A}$

This proves the equality.

To (iii). Let us denote the inverse of A'' by S . Now, exactly in the same way as in Palais [12, Lemma 17.B] we prove that $S \in \text{Pdiff}^{-1}(\tilde{X}, Y)$. First we apply Proposition 12 and choose a parametrix $T \in \text{Ell}^{-1}(\tilde{X}, Y)$ of A'' , i.e.

$$TA'' = Id - K, \text{ where } K \in \text{Pdiff}^{-1}(\tilde{X}, Y).$$

Now we have the equality

$$(Id + K + K^2 + \dots + K^m)TA'' = Id - K^{m+1}, m \geq 1,$$

and we put the operator S on both sides (from the right). We get

$$(Id + K + \dots + K^m)T = S - K^{m+1}S.$$

Thus we have $S = T + KT + \dots + K^mT + K^{m+1}S$. The operators K^jT , $j = 0, \dots, m$, belong to the class $\text{Pdiff}^{-j-1}(\tilde{X}, Y) \subset \text{Pdiff}^{-1}(\tilde{X}, Y)$, and $K^{m+1}S$ is the composition of an operator of order -1 with an operator of order $-m-1$, hence it is an operator of order $-m-2$. So we are able to approximate a pseudolocal operator $S = (A'')^{-1}$ by operators belonging to the class $\text{Pdiff}^{-1}(\tilde{X}, Y)$ up to any order. This proves that S belongs to $\text{Pdiff}^{-1}(\tilde{X}, Y)$ by a similar argument as in Hörmander [10, Theorem 5.26]. The ellipticity of the symbol follows as usual from the Fredholm properties. $\square$

26 REMARK. Note that the construction of $\tilde{A} = A' \cup A''$ does not necessarily take us out of the class of differential operators. For example $\tilde{A}$ is a well defined differential operator, if B is differential and self-adjoint. The same is not true for A'' . In fact, if $A'' \neq \tilde{A}$, i.e. if there exist non-trivial interior solutions, then the operator A'' is not local and thus definitely not a differential operator - even though A and B may be differential. Then, Corollary 25 tells us, A'' still is pseudolocal and belongs to our class Pdiff^1 .

3 The Calderón Projector - Revisited.

By the preceding Continuation Lemma 25 it is now very easy to carry out a slight modification of the constructions given in Palais [12], Seeley [14], and Hörmander [11]. The purpose of our modification is two-fold: (1) we get a *true* Calderón projector - not just an approximative one as in Hörmander [11]. And (2) our construction applies to *arbitrary* operators with splitting elliptic symbols. Thus the construction of the Calderón projector is no longer restricted to invertible operators differential over the whole manifold as in Palais [12] or to invertible operators differential near the boundary as in Seeley [14].

31 THEOREM. Let X be a compact smooth oriented Riemannian manifold with smooth boundary Y and let E and F be smooth Hermitian vector bundles over X . Let $A : C^\infty(X; E) \rightarrow C^\infty(X; F)$ be an operator of class $\text{Ell}^1(X, Y)$ with $A = G_A(y, t)(\partial_t + B_t)$ near Y where $y \in Y, t$ the (inward) normal coordinate, B_t an elliptic pseudo-differential operator of first order over Y , and $G_A(., t)$ a bundle isomorphism from $E|_Y$ to $F|_Y$. Then there exists

- (i) a pseudo-differential operator of 0-th order $P(A) : C^\infty(Y; E|_Y) \rightarrow C^\infty(Y; E|_Y)$ such that
- (ii) $P(A) = P(A)^* = P(A)^2$ and
- (iii) $\text{ran } P(A) = H(A)$, where $H(A)$ is the Cauchy data space of A over Y .
- (iv) At each point (y, η) of the covariant sphere bundle SY the principal symbol of $P(A)$ is the projection $p(y, \eta) : E_y \rightarrow E_y$ onto the direct sum of the eigenspaces of the linear transformation $b(y, \eta) : E_y \rightarrow E_y$ corresponding to the eigenvalues with positive real part. Here b denotes the principal symbol of the operator B_0 .

PROOF: We apply Continuation Lemma 25. This yields an invertible operator $A'' \in \text{Ell}^1(\tilde{X}, Y)$. We repeat the arguments of Hörmander [11, Section 20.1, pp 234-237 and pp 242-249] with the following construction: We define $P(A'') := \gamma S A''^c$, where $A''^c g := G_A(., 0)g \otimes \delta_Y$ for $g \in C^\infty(Y; E|_Y)$, δ_Y is the Dirac measure in the parameter t , $S = (A'')^{-1}$, and $\gamma : C^\infty(\tilde{X}; E \cup_G F) \rightarrow C^\infty(Y; E|_Y)$ is the restriction. Note the simplification in the form of A''^c which is due to our restriction to operators of order 1. A second simplification is due to our choice of S where we do not take a parametrix, but, instead, take the inverse of A'' , which exists and also has splitting elliptic symbol as shown in 25. Thus we obtain a pseudo-differential operator

$$P(A'') : C^\infty(Y; E|_Y) \rightarrow C^\infty(Y; E|_Y)$$

of 0-th order fulfilling (ii) and (iv) and with $\text{ran } P(A'') = H_+(A'')$. Since $H_+(A'') = H(A)$ we can define $P(A) := P(A'')$. $\square$

32 REMARK. Since our proof closely follows the arguments of Hörmander [11] one may ask whether one can deduce our result - the existence of a pseudo-differential projector $P(A)$ onto the space of Cauchy data - just from Hörmander's result, i.e. from the existence of an idempotent pseudo-differential symbol $q(y, \eta)$ with its leading part equal to $p(y, \eta)$, the principal symbol of $P(A)$. In fact, any appropriate pseudo-differential projection Q with closed range in $H(A)$ and with $\sigma_L(Q) = p$ can be deformed into the Calderón projector by a series of modifications with suitable operators of finite rank. However, this obviously is not true if the closedness of the range of Q is not assured.

33 REMARK. If $A \in \text{Ell}^1(X, Y)$ takes the form $A = G_A(\partial_t + B)$ near Y with B self-adjoint, then the spectral projection $sP(B)$ is a well-defined pseudo-differential operator of order zero. It maps $C^\infty(Y; E|_Y)$ onto the span $\{e_j\}_{j>0}$ where the $\{e_j\}$ are an orthonormal system of the eigenfunctions of B with positive eigenvalues, and the principal symbols of the spectral projection $sP(B)$ and of the Calderón projector $P(A)$ coincide, see Taylor [16, Chapter V] and our [3, Section 2]. Note, that the Calderón projector $P(A)$ depends solely on the choice of a normal vector field on X near Y which provides the splitting of A . In contrast, the spectral projection $sP(B)$ is additionally dependent on the Riemannian structure of Y and on the Hermitian structure of E . In fact, the spectral projections may belong to different connected components (in the L^2 -operator topology) of the space of pseudo-differential projections with the given principal symbol. This will be explained in a subsequent paper.

4 Examples and Concluding Remarks.

The main tool for the analysis of elliptic boundary value problems for operators of the class $\text{Ell}^1(X, Y)$ is provided by Theorem 31 (and Corollary 25). With this tool we can prove the same regularity theorems and the finiteness of the index as was done for differential operators in the beautiful lectures Seeley [15, Chapter 6]. The results of [15] concerning the realizations of the form A_R with (local or global) elliptic boundary condition R also remain valid for $A \in \text{Ell}^1(X, Y)$.

It is not difficult to extend the results of Polking [13] so that they become valid for any manifold with boundary; and not just for the cylinder $M \times I$ where M is a closed manifold.

These are (up to now) the main applications of our results. We would like to end with two easy examples which show that the class $\text{Ell}^1(X, Y)$ is really a useful class of operators. Operators of class $\text{Ell}^1(X, Y)$ namely appear naturally as the result of a reduction of operators of higher order:

41 PROPOSITION. Let A_i , $i = 1, 2$, be differential operators of order i (scalar or acting on the sections of a vector bundle) over a closed smooth oriented manifold M . Then the second order differential operator $A = \partial_t^2 + A_1 \partial_t + A_2$ over the cylinder $X = M \times I$ with boundary $Y = M \times \{0, 1\}$ can be reduced to a (non-classical) pseudo-differential operator A' with splitting symbol such that $\ker A \cong \ker A'$. Here ∂_t denotes the differentiation with respect to t , i.e. the I -direction. Moreover, A is elliptic if and only if A' belongs to the class $\text{Ell}^1(X, Y)$.

PROOF: We choose a positive self-adjoint first order elliptic operator Δ over M with principal symbol $\sigma_\Delta(y, \eta) = |\eta|$. For example we may take $\Delta = Q^*Q + 1$ with suitable Q with $\sigma_Q(y, \eta) = |\eta|^{1/2}$. Here y denotes the coordinate in M and $\eta \in T^*M_y$. Then we obtain the following reduction scheme (cf. Taylor [16, Chapter IV and Chapter V]):

Near $M \times \{0\}$ we have

$$Au = 0 \iff A'w = 0,$$

where

$$A' = \begin{pmatrix} Id & 0 \\ 0 & Id \end{pmatrix} \partial_t + \begin{pmatrix} 0 & -\Delta \\ A_2 \Delta^{-1} & A_1 \end{pmatrix}$$

and

$$w = \begin{pmatrix} \Delta u \\ \partial_t u \end{pmatrix}.$$

In fact

$$A' w = \begin{pmatrix} \partial_t \Delta u - \Delta \partial_t u \\ Au \end{pmatrix} = \begin{pmatrix} 0 \\ Au \end{pmatrix}.$$

The total symbol $a(x, \xi)$ of A' has the following form:

$$a(x, \xi) = \begin{pmatrix} i\tau & -|\eta| \\ a_2(y, \eta) |\eta|^{-1} & i\tau + a_1(y, \eta) \end{pmatrix} \\ + \text{lower order purely pseudo-differential terms,}$$

where $x = (y, t)$, $\xi = (\eta, \tau) \in (T^*X)_x$, and a_i denotes the principal symbol of A_i . Thus we obtain

$$|D_\xi^\alpha D_x^\beta a(x, \xi)| \leq C_{\alpha\beta} (1 + |\xi|)^{1-\alpha''} (1 + |\eta|)^{-|\alpha'|}$$

for all multiindices $\alpha = (\alpha_1, \dots, \alpha_{n-1}, \alpha'')$, $\alpha' = (\alpha_1, \dots, \alpha_{n-1})$ and β , so $A' \in \text{Pdiff}^1(X, Y)$. Moreover, for the principal symbols we have the relation

$$\sigma_A(x, \xi) = -\tau^2 + i\tau a_1(y, \eta) + a_2(y, \eta) = \det \sigma_{A'}(x, \xi),$$

hence A' belongs to the class $\text{Ell}^1(X, Y)$ if and only if A is elliptic. $\square$

42 REMARK. For elliptic A (and A') we find

$$H(A') = \{(\Delta u|_{\partial X}, (\partial_t u)|_{\partial X}) \mid u \in \ker A\}.$$

So, the spaces of Cauchy data for the first order operator A' and for the second order operator A are isomorphic under the transformation $(r_1, r_2) \mapsto (\Delta r_1, r_2)$, and we get for the Calderón projector

$$P(A') : L^2(M \times \{0, 1\}; E) \oplus L^2(M \times \{0, 1\}; E) \longrightarrow H(A')$$

the condition

$$P(\mathcal{A}')(r_1, r_2) = (r_1, r_2) \iff u|_{M \times \{i\}} = \Lambda^{-1} r_{1i} \text{ and } (\partial_t u)|_{M \times \{i\}} = r_{2i},$$

for all $u \in C^\infty(M \times I; E)$ and $i \in \{0, 1\}$ where $r_1 = (r_{10}, r_{11})$ and $r_2 = (r_{20}, r_{21})$.

43 REMARK. $\mathcal{A}$ is a differential operator. If additionally it is elliptic and scalar it then fulfills the unique continuation property. Hence the equivalent operator $\mathcal{A}'$ has no non-trivial interior solutions.

44 EXAMPLE. Let M be a closed smooth oriented Riemannian manifold. Let E be a Hermitian vector bundle over M , and let $B : C^\infty(M; E) \rightarrow C^\infty(M; E)$ be a self-adjoint elliptic differential operator of first order. Over the cylinder $X = M \times I$ we consider the operator

$$\mathcal{A} = \partial_t^2 - B^2 + \epsilon B \partial_t$$

where ϵ is a complex parameter. We want to make the following observations:

(i) As noticed above, we obtain

$$\sigma_{\mathcal{A}}(x, \xi) = -\tau^2 + i\tau\epsilon b(y, \eta) - b^2(y, \eta)$$

for $x = (y, t) \in X$ and $\xi = (\eta, \tau) \in (T^*X)_x$ where b denotes the principal symbol of B . Then a necessary and sufficient condition for the ellipticity of $\mathcal{A}$ is that the equation

$$-\tau^2 + i\tau\epsilon\lambda - \lambda^2 = 0$$

has no real solution τ for any eigenvalue λ of the endomorphism $b(y, \eta) : E_y \rightarrow E_y$. So, we have ellipticity for example for all real ϵ and for purely imaginary ϵ with $|\epsilon| \leq 2$. In the last case $\mathcal{A}$ becomes self-adjoint.

(ii) Now assume that $\mathcal{A}$ is elliptic. We apply Proposition 41 and obtain an equivalent first order operator with splitting symbol

$$\mathcal{A}' = \begin{pmatrix} Id & 0 \\ 0 & Id \end{pmatrix} \partial_t + \begin{pmatrix} 0 & -\Lambda \\ -B^2 \Lambda^{-1} & \epsilon B \end{pmatrix}$$

where we choose $\Lambda := \sqrt{1 + B^2}$. Then we have

$$\mathcal{A}' = \begin{pmatrix} Id & 0 \\ 0 & Id \end{pmatrix} \partial_t + B$$

with

$$B = \begin{pmatrix} 0 & -\sqrt{1 + B^2} \\ \frac{B^2}{\sqrt{1 + B^2}} & \epsilon B \end{pmatrix} : C^\infty(E \oplus E) \rightarrow C^\infty(E \oplus E).$$

(iii) Recall that elliptic self-adjoint operators of positive order have a discrete spectrum of finite multiplicity $\{\lambda_k\}_{k \in \mathbb{Z}}$. Moreover, there exists no essential spectrum, and the eigenvectors $\{\phi_k\}$ span the whole $L^2 E$. Let $\{\lambda_k, \phi_k\}_{k \in \mathbb{Z}}$ denote such a spectral decomposition of $L^2 E$ related to our self-adjoint operator B . Then

$\{\sqrt{1+\lambda_k^2}, \phi_k\}_{k \in \mathbb{Z}}$ is a spectral decomposition related to the operator $\sqrt{1+B^2}$.

We want to construct a set of solutions of $Au = 0$ from that information.

Let E_k denote the one-dimensional space spanned by ϕ_k . Then B maps $E_k \oplus E_j$ into $E_k \oplus E_j$ for all $k, j \in \mathbb{Z}$. We define $B_{kj} := B|_{E_k \oplus E_j}$ and obtain $B = \sum_{k,j} B_{kj}$.

For any $c, c' \in \mathbb{C}$ we have

$$B(c\phi_k, c'\phi_k) = (\gamma\phi_k, \gamma'\phi_k)$$

with

$$\begin{pmatrix} \gamma \\ \gamma' \end{pmatrix} = \begin{pmatrix} 0 & -\sqrt{1+\lambda_k^2} \\ \frac{-\lambda_k^2}{\sqrt{1+\lambda_k^2}} & \epsilon\lambda_k \end{pmatrix} \begin{pmatrix} c \\ c' \end{pmatrix}.$$

So the operator B_{kk} takes the form

$$B_{kk} = \lambda_k \begin{pmatrix} 0 & \left(\frac{-\lambda_k}{\sqrt{1+\lambda_k^2}}\right)^{-1} \\ \frac{-\lambda_k^2}{\sqrt{1+\lambda_k^2}} & \epsilon \end{pmatrix}.$$

We determine the eigenvalues and eigenfunctions of B_{kk} . For $a \in \mathbb{R}$ we have

$$\begin{pmatrix} 0 & a^{-1} \\ a & \epsilon \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} a^{-1}v \\ au + \epsilon v \end{pmatrix},$$

hence

$$\mu u = a^{-1}v \quad \text{and} \quad \mu v = au + \epsilon v$$

if and only if $\mu^2 - \epsilon\mu - 1 = 0$, i.e.

$$\mu_{\pm} = \frac{\epsilon}{2} \pm \sqrt{1 + \frac{\epsilon^2}{4}}.$$

The corresponding eigenvectors are

$$\begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 1 \\ a \left(\frac{\epsilon}{2} \pm \sqrt{1 + \frac{\epsilon^2}{4}} \right) \end{pmatrix}.$$

Therefore, the eigenvalues of B_{kk} are $\mu_{kk} = \lambda_k \mu_{\pm}$ and the eigenfunctions are

$$\psi_{kk} = \left(\phi_k, -\frac{\lambda_k \mu_{\pm}}{\sqrt{1+\lambda_k^2}} \phi_k \right).$$

With similar - but more laborious - work we determine the eigenvalues and eigenfunctions of the partial operators B_{kj} for $j \neq k$.

(iv) As explained by the authors in [4, Example 3.5(a)], any solution of

$$\mathcal{A}'w = \left(\begin{pmatrix} Id & 0 \\ 0 & Id \end{pmatrix} \partial_t + \mathcal{B} \right) w = 0$$

has the form

$$w = \sum_{\alpha_{kj} \in \mathbb{Z}} \exp(-t\mu_{kj\pm}) \psi_{kj\pm},$$

and the space of Cauchy data $H(\mathcal{A}')$ is the subspace of $L^2(M \times \{0\}; E \oplus E) \oplus L^2(M \times \{1\}; E \oplus E)$ with the orthonormal base

$$\frac{1}{\sqrt{1 + e^{2\mu_{kj\pm}}}} \begin{pmatrix} \psi_{kj\pm} & e^{\mu_{kj\pm}} \psi_{kj\pm} \end{pmatrix} \begin{matrix} \text{over } M \times \{0\} \\ \text{over } M \times \{1\} \end{matrix}.$$

(v) We finally obtain the solutions of $\mathcal{A}u = 0$ by back transformation following the rules given in the proof of Proposition 41. So, any $w \in \ker \mathcal{A}'$ provides a $u \in \ker \mathcal{A}$. In particular we obtain for any $k \in \mathbb{Z}$ a solution

$$u_k = \exp \left(-t\lambda_k \left(\epsilon/2 \pm \sqrt{1 + \epsilon^2/4} \right) \right) (1 + \lambda_k^2)^{-\frac{1}{2}} \phi_k.$$

ACKNOWLEDGEMENT. The authors wish to thank Gerd Grubb for her very helpful comments and suggestions.

REFERENCES

- (1) Birman, M. and Solomyak, A., On subspaces which admit pseudodifferential projections. *Vestnik Leningrad University* 82 no. 1 (1982), 18-25 (Russian).
- (2) Boöf-Baynbek, B., *Elliptic Boundary Value Problems - Revisited*. Lecture Notes, Roskilde, 1987 (preprint).
- (3) Boöf, B. and Wojciechowski, K., Desuspension of splitting elliptic symbols. I. *Ann. Glob. Analysis and Geometry* 3 (1985), 337-383.
- (4) -, Desuspension of splitting elliptic symbols. II. *Ann. Glob. Analysis and Geometry* 4 (1986), forthcoming.
- (5) Calderón, A. P., Boundary value problems for elliptic equations. In: *Outlines of the Joint Soviet-American Symposium on Partial Differential Equations*, Novosibirsk, 1963, 303-304.
- (6) -, *Lecture Notes on Pseudo-Differential Operators and Elliptic Boundary Value Problems. I*. Consejo Nacional de Investigaciones Científicas y Técnicas, Inst. Argentino de Matematica, Buenos Aires, 1976.
- (7) Grubb, G., Boundary problems for systems of partial differential operators of mixed order. *J. Functional Analysis* 26 (1977), 131-165.
- (8) -, Problèmes aux limites pseudo-différentiels dépendant d'un paramètre. *C.R. Acad. Sci. Paris* 292 (1981), 581-583.

- (9) Hörmander, L., Pseudo-differential operators and non-elliptic boundary problems. *Ann. of Math.* 83 (1966), 129-209.
- (10) -, *The Analysis of Linear Partial Differential Operators. I.* Springer-Verlag, Berlin (West), 1983.
- (11) -, *The Analysis of Linear Partial Differential Operators. III.* Springer-Verlag, Berlin (West), 1985.
- (12) Palais, R. S. (ed.), *Seminar on the Atiyah-Singer Index Theorem.* Ann. of Math. Studies 57. Princeton University Press, Princeton, 1965.
- (13) Polking, J. C., Boundary value problems for parabolic systems of partial differential equations. *Amer. Math. Soc. Proc. Symp. Pure Math.* 10 (1967), 243-274.
- (14) Seeley, R. T., Singular integrals and boundary value problems. *Amer. J. Math.* 88 (1966), 781-809.
- (15) -, Topics in pseudo-differential operators. In: *CIME Conference on Pseudo-Differential Operators (Stresa 1968)*. Ed. Cremonese, Rome, 1969, 167-305.
- (16) Taylor, M. E., *Pseudodifferential Operators.* Princeton University Press, Princeton, 1981.
- (17) Treves, F., *Pseudodifferential and Fourier Integral Operators. I.* Plenum Press, New York, 1980.
- (18) Wojciechowski, K. P., Elliptic operators and relative K-homology groups on manifolds with boundary. *C.R. Math. Rep. Acad. Sci. Canada* 7 (1985), 149-154.
- (19) -, *Cauchy data spaces and boundary map in relative K-homology for manifolds with boundary.* Notes, Warsaw, 1986 (preprint).

BBB, IMFUFA, Roskilde Universitetscenter, DK-4000 Roskilde, Denmark

KPW, Instytut Matematyki, Uniwersytet Warszawski, PL-00-901 PKiN Warszawa, Poland

- 1/78 "TANKER OM EN PRAKSIS" - et matematikprojekt.
Projektrapport af: Anne Jensen, Lena Lindenskov, Marianne Kesselhahn og Nicolai Lomholt.
Vejleder: Anders Madsen
- 2/78 "OPTIMERING" - Menneskets forøgede beherskelsermuligheder af natur og samfund.
Projektrapport af: Tom J. Andersen, Tommy R. Andersen, Gert Krenø og Peter H. Lassen
Vejleder: Bernhelm Boss.
- 3/78 "OPCAVESAMLING", breddekursus i fysik.
Af: Lasse Rasmussen, Aage Bonde Kræmmer og Jens Højgaard Jensen.
- 4/78 "TRE ESSAYS" - om matematikundervisning, matematiklæreruddannelsen og videnskabsrindalismen.
Af: Mogens Niss
Nr. 4 er p.t. udgået.
- 5/78 "BIBLIOGRAFISK VEJLEDNING til studiet af DEN MODERNE FYSIKS HISTORIE".
Af: Helge Kragh.
Nr. 5 er p.t. udgået.
- 6/78 "NOGLE ARTIKLER OG DEBATINDLÆG OM - læreruddannelse og undervisning i fysik, og - de naturvidenskabelige fags situation efter studenteroprøret".
Af: Karin Beyer, Jens Højgaard Jensen og Bent C. Jørgensen.
- 7/78 "MATEMATIKKENS FORHOLD TIL SAMFUNDSØKONOMIEN".
Af: B.V. Gnedenko.
Nr. 7 er udgået.
- 8/78 "DYNAMIK OG DIAGRAMMER". Introduktion til energy-bond-graph formalismen.
Af: Peder Voetmann Christiansen.
- 9/78 "OM PRAKSIS' INDFLYDELSE PÅ MATEMATIKKENS UDVIKLING". - Motiver til Kepler's: "Nova Stereometria Doliorum Vinariorum".
Projektrapport af: Lasse Rasmussen.
Vejleder: Anders Madsen.
-
- 10/79 "TERMODYNAMIK I GYMNASIET".
Projektrapport af: Jan Christensen og Jeanne Mortensen,
Vejledere: Karin Beyer og Peder Voetmann Christiansen.
- 11/79 "STATISTISKE MATERIALER".
Af: Jørgen Larsen.
- 12/79 "LINEÆRE DIFFERENTIALLIGNINGER OG DIFFERENTIALLIGNINGSSYSTEMER".
Af: Mogens Brun Heefelt.
Nr. 12 er udgået.
- 13/79 "CAVENDISH'S FORSØG I GYMNASIET".
Projektrapport af: Gert Kreinø.
Vejleder: Albert Chr. Paulsen.
- 14/79 "BOOKS ABOUT MATHEMATICS: History, Philosophy, Education, Models, System Theory, and Works of".
Af: Else Høyrup.
Nr. 14 er p.t. udgået.
- 15/79 "STRUKTUREL STABILITET OG KATASTROFER i systemer i og udenfor termodynamisk ligevægt".
Specialeopgave af: Leif S. Striegler.
Vejleder: Peder Voetmann Christiansen.
- 16/79 "STATISTIK I KRÆFTFORSKNINGEN".
Projektrapport af: Michael Olsen og Jørn Jensen.
Vejleder: Jørgen Larsen.
- 17/79 "AT SPØRGE OG AT SVARE i fysikundervisningen".
Af: Albert Christian Paulsen.
- 18/79 "MATHEMATICS AND THE REAL WORLD", Proceedings af an International Workshop, Roskilde University Centre, Denmark, 1978.
Preprint.
Af: Bernhelm Booss og Mogens Niss (eds.)
- 19/79 "GEOMETRI, SKOLE OG VIRKELIGHED".
Projektrapport af: Tom J. Andersen, Tommy R. Andersen og Per H.H. Larsen.
Vejleder: Mogens Niss.
- 20/79 "STATISTISKE MODELLER TIL BESTEMMELSE AF SIKRE DOSER FOR CARCINOGENE STOFFER".
Projektrapport af: Michael Olsen og Jørn Jensen.
Vejleder: Jørgen Larsen
- 21/79 "KONTROL I GYMNASIET-FORMÅL OG KONSEKVENSER".
Projektrapport af: Crilles Bacher, Per S.Jensen, Preben Jensen og Torben Nysteen.
- 22/79 "SEMIOTIK OG SYSTEMEGENSKABER (1)".
1-port lineært response og støj i fysikken.
Af: Peder Voetmann Christiansen.
- 23/79 "ON THE HISTORY OF EARLY WAVE MECHANICS - with special emphasis on the role of reality".
Af: Helge Kragh.
-
- 24/80 "MATEMATIKOPFATTELSE HOS 2.G'ERE".
a+b 1. En analyse. 2. Interviewmateriale.
Projektrapport af: Jan Christensen og Knud Lindhardt Rasmussen.
Vejleder: Mogens Niss.
- 25/80 "EKSAMENSOPGAVER", Dybdemodulet/fysik 1974-79.
- 26/80 "OM MATEMATISKE MODELLER".
En projektrapport og to artikler.
Af: Jens Højgaard Jensen m.fl.
- 27/80 "METHODOLOGY AND PHILOSOPHY OF SCIENCE IN PAUL DIRAC'S PHYSICS".
Af: Helge Kragh.
- 28/80 "DIELEKTRISK RELAXATION - et forslag til en ny model bygget på væskernes viscoelastiske egenskaber".
Projektrapport af: Gert Kreinø.
Vejleder: Niels Boye Olsen.
- 29/80 "ODIN - undervisningsmateriale til et kursus i differentiaalligningsmodeller".
Projektrapport af: Tommy R. Andersen, Per H.H. Larsen og Peter H. Lassen.
Vejleder: Mogens Brun Heefelt.
- 30/80 "FUSIONSENERGIEN - - - ATOMSAMFUNDETS ENDESTATION".
Af: Oluf Danielsen.
Nr. 30 er udgået.
- 31/80 "VIDENSKABSTEORETISKE PROBLEMER VED UNDERVISNINGSSYSTEMER BASERET PÅ MÆNGDELÆRE".
Projektrapport af: Troels Lange og Jørgen Karrebæk.
Vejleder: Stig Andur Pedersen.
Nr. 31 er p.t. udgået.
- 32/80 "POLYMERE STOFFERS VISCOELASTISKE EGENSKABER - BELYST VED HJÆLP AF MEKANISKE IMPEDANSMÅLINGER MØSSBAUEREFFEKTIVITETER".
Projektrapport af: Crilles Bacher og Preben Jensen.
Vejledere: Niels Boye Olsen og Peder Voetmann Christiansen.
- 33/80 "KONSTITUERING AF FAG INDEN FOR TEKNISK - NATURVIDENSKABELIGE UDDANNELSER. I-II".
Af: Arne Jakobsen.
- 34/80 "ENVIRONMENTAL IMPACT OF WIND ENERGY UTILIZATION".
ENERGY SERIES NO. I.
Af: Bent Sørensen
Nr. 34 er udgået.

- 35/80 "HISTORISKE STUDIER I DEN NYERE ATOMFYSIKS UDVIKLING".
Af: Helge Kragh.
- 36/80 "HVAD ER MENINGEN MED MATEMATIKUNDERVISNINGEN?".
Fire artikler.
Af: Mogens Niss.
- 37/80 "RENEWABLE ENERGY AND ENERGY STORAGE".
ENERGY SERIES NO. 2.
Af: Bent Sørensen.
-
- 38/81 "TIL EN HISTORIETEORI OM NATURERKENDELSE, TEKNOLOGI OG SAMFUND".
Projektrapport af: Erik Gade, Hans Hedal, Henrik Lau og Finn Physant.
Vejledere: Stig Andur Pedersen, Helge Kragh og Ib Thiersen.
Nr. 38 er p.t. udgået.
- 39/81 "TIL KRITIKKEN AF VÆKSTØKONOMIEN".
Af: Jens Højgaard Jensen.
- 40/81 "TELEKOMMUNIKATION I DANMARK - oplæg til en teknologivurdering".
Projektrapport af: Arne Jørgensen, Bruno Petersen og Jan Vedde.
Vejleder: Per Nørregaard.
- 41/81 "PLANNING AND POLICY CONSIDERATIONS RELATED TO THE INTRODUCTION OF RENEWABLE ENERGY SOURCES INTO ENERGY SUPPLY SYSTEMS".
ENERGY SERIES NO. 3.
Af: Bent Sørensen.
- 42/81 "VIDENSKAB TEORI SAMFUND - En introduktion til materialistiske videnskabsopfattelser".
Af: Helge Kragh og Stig Andur Pedersen.
- 43/81 1. "COMPARATIVE RISK ASSESSMENT OF TOTAL ENERGY SYSTEMS".
2. "ADVANTAGES AND DISADVANTAGES OF DECENTRALIZATION".
ENERGY SERIES NO. 4.
Af: Bent Sørensen.
- 44/81 "HISTORISKE UNDERSØGELSER AF DE EKSPERIMENTELLE FORUDSÆTNINGER FOR RUTHERFORDS ATOMMODEL".
Projektrapport af: Niels Thor Nielsen.
Vejleder: Bent C. Jørgensen.
-
- 45/82 Er aldrig udkommet.
- 46/82 "EKSEMPLARISK UNDERVISNING OG FYSISK ERKENDELSE-1+1 ILLUSTRERET VED TO EKSEMPLER".
Projektrapport af: Torben O. Olsen, Lasse Rasmussen og Niels Dreyer Sørensen.
Vejleder: Bent C. Jørgensen.
- 47/82 "BARBERÄCK OG DET VÆRST OFFICIELT-TÆNKELIGE UHELD".
ENERGY SERIES NO. 5.
Af: Bent Sørensen.
- 48/82 "EN UNDERSØGELSE AF MATEMATIKUNDERVISNINGEN PÅ ADGANGSKURSUS TIL KØBENHAVNS TEKNISKUM".
Projektrapport af: Lis Ellertzen, Jørgen Karrebæk, Troels Lange, Preben Nørregaard, Lissi Pedersen, Laust Rishøj, Lill Røn og Isac Showiki.
Vejleder: Mogens Niss.
- 49/82 "ANALYSE AF MULTISPEKTRALE SATELLITBILLEDER".
Projektrapport af: Preben Nørregaard.
Vejledere: Jørgen Larsen og Rasmus Ole Rasmussen.
- 50/82 "HERSLEV - MULIGHEDER FOR VEDVARENDE ENERGI I EN LANDSBY".
ENERGY SERIES NO. 6.
Rapport af: Bent Christensen, Bent Hove Jensen, Dennis B. Møller, Bjarne Laursen, Bjarne Lillethorup og Jacob Mørch Pedersen.
Vejleder: Bent Sørensen.
- 51/82 "HVAD KAN DER GØRES FOR AT AFHJÆLPE PIGERS BLOKERING OVERFOR MATEMATIK?".
Projektrapport af: Lis Ellertzen, Lissi Pedersen, Lill Røn og Susanne Stender.
- 52/82 "DESUSPENSION OF SPLITTING ELLIPTIC SYMBOLS".
Af: Bernhelm Booss og Krzysztof Wojciechowski.
- 53/82 "THE CONSTITUTION OF SUBJECTS IN ENGINEERING EDUCATION".
Af: Arne Jacobsen og Stig Andur Pedersen.
- 54/82 "FUTURES RESEARCH" - A Philosophical Analysis of Its Subject-Matter and Methods.
Af: Stig Andur Pedersen og Johannes Witt-Hansen.
- 55/82 "MATEMATISKE MODELLER" - Litteratur på Roskilde Universitetsbibliotek.
En biografi.
Af: Else Højrup.

Vedr. tekst nr. 55/82 se også tekst nr. 62/83.
- 56/82 "EN - TO - MANGE" -
En undersøgelse af matematisk økologi.
Projektrapport af: Troels Lange.
Vejleder: Anders Madsen.
-
- 57/83 "ASPECT EKSPERIMENTET"-
Skjulte variable i kvantemekanikken?
Projektrapport af: Tom Juul Andersen.
Vejleder: Peder Voetmann Christiansen.
Nr. 57 er udgået.
- 58/83 "MATEMATISKE VANDRINGER" - Modelbetragtninger over spredning af dyr mellem småbiotoper i agerlandet.
Projektrapport af: Per Hammershøj Jensen og Lene Vagn Rasmussen.
Vejleder: Jørgen Larsen.
- 59/83 "THE METHODOLOGY OF ENERGY PLANNING".
ENERGY SERIES NO. 7.
Af: Bent Sørensen.
- 60/83 "MATEMATISK MODEKSPERTISE"- et eksempel.
Projektrapport af: Erik O. Gade, Jørgen Karrebæk og Preben Nørregaard.
Vejleder: Anders Madsen.
- 61/83 "FYSIKS IDEOLOGISKE FUNKTION, SOM ET EKSEMPEL PÅ EN NATURVIDENSKAB - HISTORISK SET".
Projektrapport af: Annette Post Nielsen.
Vejledere: Jens Højrup, Jens Højgaard Jensen og Jørgen Vogelius.
- 62/83 "MATEMATISKE MODELLER" - Litteratur på Roskilde Universitetsbibliotek.
En biografi 2. rev. udgave.
Af: Else Højrup.
- 63/83 "CREATING ENERGY FUTURES: A SHORT GUIDE TO ENERGY PLANNING".
ENERGY SERIES NO. 8.
Af: David Crossley og Bent Sørensen.
- 64/83 "VON MATEMATIK UND KRIEG".
Af: Bernhelm Booss og Jens Højrup.
- 65/83 "ANVENDT MATEMATIK - TEORI ELLER PRAKSIS".
Projektrapport af: Per Hedegård Andersen, Kirsten Habekost, Carsten Holst-Jensen, Annelise von Moos, Else Marie Pedersen og Erling Møller Pedersen.
Vejledere: Bernhelm Booss og Klaus Grünbaum.
- 66/83 "MATEMATISKE MODELLER FOR PERIODISK SELEKTION I ESCHERICHIA COLI".
Projektrapport af: Hanne Lisbet Andersen, Ole Richard Jensen og Klavs Frisdahl.
Vejledere: Jørgen Larsen og Anders Hede Madsen.
- 67/83 "ELEPSOIDE METODEN - EN NY METODE TIL LINEÆR PROGRAMMERING?".
Projektrapport af: Lone Billmann og Lars Boye.
Vejleder: Mogens Brun Heefelt.
- 68/83 "STOKASTISKE MODELLER I POPULATIONSGENETIK" - til kritikken af teoriladede modeller.
Projektrapport af: Lise Odgård Gade, Susanne Hansen, Michael Hviid og Frank Mølgård Olsen.
Vejleder: Jørgen Larsen.

- 69/83 "ELEVFORUDSÆTNINGER I FYSIK"
- en test i l.g med kommentarer.
Af: Albert C. Paulsen.
- 70/83 "INDLÆRINGS - OG FORMIDLINGSPROBLEMER I MATEMATIK PÅ VOKSENUNDERVISNINGSNIVEAU".
Projektrapport af: Hanne Lisbet Andersen, Torben J. Andreasen, Svend Åge Houmann, Helle Glorup Jensen, Keld Fl. Nielsen, Lene Vagn Rasmussen.
Vejleder: Klaus Grünbaum og Anders Hede Madsen.
- 71/83 "PIGER OG FYSIK"
- et problem og en udfordring for skolen?
Af: Karin Beyer, Sussanne Blegaa, Birthe Olsen, Jette Reich og Mette Vedelsby.
- 72/83 "VERDEN IFØLGE PEIRCE" - to metafysiske essays, om og af C.S Peirce.
Af: Peder Voetmann Christiansen.
- 73/83 "'EN ENERGIANALYSE AF LANDERUG"
- økologisk contra traditionelt.
ENERGY SERIES NO. 9
Specialeopgave i fysik af: Bent Hove Jensen.
Vejleder: Bent Sørensen.
-
- 74/84 "MINIATURISERING AF MIKROELEKTRONIK" - om videnskabeliggjort teknologi og nytten af at lære fysik.
Projektrapport af: Bodil Harder og Linda Szkotak Jensen.
Vejledere: Jens Højgaard Jensen og Bent C. Jørgensen.
- 75/84 "MATEMATIKUNDERVISNINGEN I FREMTIDENS GYMNASIUM"
- Case: Lineær programmering.
Projektrapport af: Morten Blomhøj, Klavs Frisdahl og Frank Mølgaard Olsen.
Vejledere: Mogens Brun Heefelt og Jens Bjørneboe.
- 76/84 "KERNEKRAFT I DANMARK?" - Et høringssvar indkaldt af miljøministeriet, med kritik af miljøstyrelsens rapporter af 15. marts 1984.
ENERGY SERIES No. 10
Af: Niels Boye Olsen og Bent Sørensen.
- 77/84 "POLITISKE INDEKS - FUP ELLER FAKTA?"
Opinionsundersøgelser belyst ved statistiske modeller.
Projektrapport af: Svend Åge Houmann, Keld Nielsen og Susanne Stender.
Vejledere: Jørgen Larsen og Jens Bjørneboe.
- 78/84 "JÆVNSTRØMSLEDNINGSEVNE OG GITTERSTRUKTUR I AMORFT GERMANIUM".
Specialrapport af: Hans Heddal, Frank C. Ludvigsen og Finn C. Physant.
Vejleder: Niels Boye Olsen.
- 79/84 "MATEMATIK OG ALMENDANNELSE".
Projektrapport af: Henrik Coster, Mikael Wennerberg Johansen, Povl Kattler, Birgitte Lydholm og Morten Overgaard Nielsen.
Vejleder: Bernhelm Booss.
- 80/84 "KURSUSMATERIALE TIL MATEMATIK B".
Af: Mogens Brun Heefelt.
- 81/84 "FREKVENSAFHÆNGIG LEDNINGSEVNE I AMORFT GERMANIUM".
Specialrapport af: Jørgen Wind Petersen og Jan Christensen.
Vejleder: Niels Boye Olsen.
- 82/84 "MATEMATIK - OG FYSIKUNDERVISNINGEN I DET AUTOMATISEREDE SAMFUND".
Rapport fra et seminar afholdt i Hvidovre 25-27 april 1983.
Red.: Jens Højgaard Jensen, Bent C. Jørgensen og Mogens Niss.
- 83/84 "ON THE QUANTIFICATION OF SECURITY":
PEACE RESEARCH SERIES NO. 1.
Af: Bent Sørensen
nr. 83 er p.t. udgået
- 84/84 "NOGLE ARTIKLER OM MATEMATIK, FYSIK OG ALMENDANNELSE".
Af: Jens Højgaard Jensen, Mogens Niss m. fl.
- 85/84 "CENTRIFUGALREGULATORER OG MATEMATIK".
Specialrapport af: Per Hedegård Andersen, Carsten Holst-Jensen, Else Marie Pedersen og Erling Møller Pedersen.
Vejleder: Stig Andur Pedersen.
- 86/84 "SECURITY IMPLICATIONS OF ALTERNATIVE DEFENSE OPTIONS FOR WESTERN EUROPE".
PEACE RESEARCH SERIES NO. 2
Af: Bent Sørensen.
- 87/84 "A SIMPLE MODEL OF AC HOPPING CONDUCTIVITY IN DISORDERED SOLIDS".
Af: Jeppe C. Dyre.
- 88/84 "RISE, FALL AND RESURRECTION OF INFINITESIMALS".
Af: Detlef Laugwitz.
- 89/84 "FJERNVARMEOPTIMERING".
Af: Bjarne Lillethorup og Jacob Mørch Pedersen.
- 90/84 "ENERGI I L.G - EN TEORI FOR TILRETTELÆGGELSE".
Af: Albert Chr. Paulsen.
-
- 91/85 "KVANTETEORI FOR GYMNASIET".
1. Lærervejledning
Projektrapport af: Biger Lundgren, Henning Sten Hansen og John Johansson.
Vejleder: Torsten Meyer.
- 92/85 "KVANTETEORI FOR GYMNASIET".
2. Materiale
Projektrapport af: Biger Lundgren, Henning Sten Hansen og John Johansson.
Vejleder: Torsten Meyer.
- 93/85 "THE SEMIOTICS OF QUANTUM - NON - LOCALITY".
Af: Peder Voetmann Christiansen.
- 94/85 "TREENIGHEDEN BOURBAKI - generalen, matematikeren og ånden".
Projektrapport af: Morten Blomhøj, Klavs Frisdahl og Frank M. Olsen.
Vejleder: Mogens Niss.
- 95/85 "AN ALTERNATIVE DEFENSE PLAN FOR WESTERN EUROPE".
PEACE RESEARCH SERIES NO. 3
Af: Bent Sørensen
- 96/85 "ASPEKTER VED KRAFTVARMEFORSYNING".
Af: Bjarne Lillethorup.
Vejleder: Bent Sørensen.
- 97/85 "ON THE PHYSICS OF A.C. HOPPING CONDUCTIVITY".
Af: Jeppe C. Dyre.
- 98/85 "VALGMULIGHEDER I INFORMATIONSAALDEREN".
Af: Bent Sørensen.
- 99/85 "Der er langt fra Q til R".
Projektrapport af: Niels Jørgensen og Mikael Klintorp.
Vejleder: Stig Andur Pedersen.
- 100/85 "TALSYSTEMETS OPBYGNING".
Af: Mogens Niss.
- 101/85 "EXTENDED MOMENTUM THEORY FOR WINDMILLS IN PERTURBATIVE FORM".
Af: Ganesh Sengupta.
- 102/85 OPSTILLING OG ANALYSE AF MATEMATISKE MODELLER, BELYST VED MODELLER OVER KØRS FODEROPTAGELSE OG - OMSÆTNING".
Projektrapport af: Lis Eilertzen, Kirsten Habekost, Lilli Røn og Susanne Stender.
Vejleder: Klaus Grünbaum.

- 103/85 "ØDSLE KOLDKRIGERE OG VIDENSKABENS LYSE IDEER".
Projekttraktat af: Niels Ole Dam og Kurt Jensen.
Vejleder: Bent Sørensen.
- 104/85 "ANALOGREGNEMASKINEN OG LORENZLIGNINGER".
Af: Jens Jøger.
- 105/85 "THE FREQUENCY DEPENDENCE OF THE SPECIFIC HEAT OF THE GLASS TRANSITION".
Af: Tage Christensen.
- "A SIMPLE MODEL OF AC HOPPING CONDUCTIVITY".
Af: Jeppe C. Dyre.
Contributions to the Third International Conference on the Structure of Non - Crystalline Materials held in Grenoble July 1985.
- 106/85 "QUANTUM THEORY OF EXTENDED PARTICLES".
Af: Bent Sørensen.
- 107/85 "EN MYG GØR INGEN EPIDEMI".
- flodblindhed som eksempel på matematisk modellering af et epidemiologisk problem.
Projekttraktat af: Per Hedegård Andersen, Lars Boye, Carsten Holst Jensen, Else Marie Pedersen og Erling Møller Pedersen.
Vejleder: Jesper Larsen.
- 108/85 "APPLICATIONS AND MODELLING IN THE MATHEMATICS CURRICULUM" - state and trends -
Af: Mogens Niss.
- 109/85 "COX I STUDIETIDEN" - Cox's regressionsmodel anvendt på studenteroplysninger fra RUC.
Projekttraktat af: Mikael Wennerberg Johansen, Poul Katler og Torben J. Andreassen.
Vejleder: Jørgen Larsen.
- 110/85 "PLANNING FOR SECURITY".
Af: Bent Sørensen
- 111/85 "JORDEN RUNDT PÅ FLADE KORT".
Projekttraktat af: Birgit Andresen, Beatriz Quinones og Jimmy Staal.
Vejleder: Mogens Niss.
- 112/85 "VIDENSKABELIGGØRELSE AF DANSK TEKNOLOGISK INNOVATION FRA 1950 - BELYST VED EKSEMPLER".
Projekttraktat af: Erik Odgaard Gade, Hans Hødal, Frank C. Ludvigsen, Annette Post Nielsen og Finn Physant.
Vejleder: Claus Bryld og Bent C. Jørgensen.
- 113/85 "DESUSPENSION OF SPLITTING ELLIPTIC SYMBOLS 11".
Af: Bernhelm Booss og Krzysztof Wojciechowski.
- 114/85 "ANVENDELSE AF GRAFISKE METODER TIL ANALYSE AF KONFIGURATIONER".
Projekttraktat af: Lone Billmann, Ole R. Jensen og Arne-Lise von Moos.
Vejleder: Jørgen Larsen.
- 115/85 "MATEMATIKKENS UDVIKLING OP TIL RENESSANCEN".
Af: Mogens Niss.
- 116/85 "A PHENOMENOLOGICAL MODEL FOR THE MEYER-NELDEL RULE".
Af: Jeppe C. Dyre.
- 117/85 "KRAFT & FJERNVARMEOPTIMERING".
Af: Jacob Mørch Pedersen.
Vejleder: Bent Sørensen
- 118/85 "TILFÆLDIGHEDEN OG NØDVENDIGHEDEN IFØLGE PEIRCE OG FYSIKKEN".
Af: Peder Voetmann Christiansen
- 120/86 "ET ANTAL STATISTISKE STANDARDMODELLER".
Af: Jørgen Larsen
- 121/86 "SIMULATION I KONTINUERT TID".
Af: Peder Voetmann Christiansen.
- 122/86 "ON THE MECHANISM OF GLASS IONIC CONDUCTIVITY".
Af: Jeppe C. Dyre.
- 123/86 "GYMNASIEFYSIKKEN OG DEN STORE VERDEN".
Fysiklærerforeningen, IMFUA, RUC.
- 124/86 "OPGAVESAMLING I MATEMATIK".
Samtlige opgaver stillet i tiden 1974-jan. 1986.
- 125/86 "UVBY, 8 - systemet - en effektiv fotometrisk spektral-klassifikation af B-, A- og F-stjerner".
Projekttraktat af: Birger Lundgren.
- 126/86 "OM UDVIKLINGEN AF DEN SPECIELLE RELATIVITETSTEORI".
Projekttraktat af: Lise Odgaard & Linda Szkotak Jensen
Vejledere: Karin Beyer & Stig Andur Pedersen.
- 127/86 "GALOIS' BIDRAG TIL UDVIKLINGEN AF DEN ABSTRAKTE ALGEBRA".
Projekttraktat af: Pernille Sand, Heine Larsen & Lars Frandsen.
Vejleder: Mogens Niss.
- 128/86 "SMÅKRYB" - om ikke-standard analyse.
Projekttraktat af: Niels Jørgensen & Mikael Klinton.
Vejleder: Jeppe Dyre.
- 129/86 "PHYSICS IN SOCIETY"
Lecture Notes 1983 (1986)
Af: Bent Sørensen
- 130/86 "Studies in Wind Power"
Af: Bent Sørensen
- 131/86 "FYSIK OG SAMFUND" - Et integreret fysik/historie-projekt om naturanskuelsens historiske udvikling og dens samfundsmæssige betingethed.
Projekttraktat af: Jakob Heckscher, Søren Brønd, Andy Wierød.
Vejledere: Jens Høyrup, Jørgen Vogelius, Jens Højgaard Jensen.
- 132/86 "FYSIK OG DANDELSE"
Projekttraktat af: Søren Brønd, Andy Wierød.
Vejledere: Karin Beyer, Jørgen Vogelius.
- 133/86 "CHERNOBYL ACCIDENT: ASSESSING THE DATA. ENERGY SERIES NO. 15."
Af: Bent Sørensen.
-
- 134/87 "THE D.C. AND THE A.C. ELECTRICAL TRANSPORT IN AsSeTe SYSTEM"
Authors: M.B.El-Den, N.B.Olsen, Ib Høst Pedersen, Petr Visčor
- 135/87 "INTUITIONISTISK MATEMATIKS METODER OG ERKENDELSESTEORETISKE FORUDSÆTNINGER"
MATEMATIKSPECIALE: Claus Larsen
Vejledere: Anton Jensen og Stig Andur Pedersen
- 136/87 "Mystisk og naturlig filosofi: En skitse af kristendommens første og andet møde med græsk filosofi"
Projekttraktat af Frank Colding Ludvigsen
Vejledere: Historie: Ib Thiersen
Fysik: Jens Højgaard Jensen
- 137/87 "HOPMODELLER FOR ELEKTRISK LEDNING I UORDNEDE FASTE STOFFER" - Resume af licentiatafhandling
Af: Jeppe Dyre
Vejledere: Niels Boye Olsen og Peder Voetmann Christiansen.
- 119/86 "DET ER GANSKE VIST - - EUKLIDS FEMTE POSTULAT KUNNE NOK SKABE RØRE I ANDEDAMMEN".
Af: Ib M. Christiansen
Vejleder: Mogens Niss.

138/87 "JOSEPHSON EFFECT AND CIRCLE MAP."

Paper presented at The International
Workshop on Teaching Nonlinear Phenomena
at Universities and Schools, "Chaos in
Education". Balaton, Hungary; 26 April-2 May 1987.

By: Peder Voetmann Christiansen

139/87 "Machbarkeit nichtbeherrschbarer Technik
durch Fortschritte in der Erkennbarkeit
der Natur"

Af: Bernhelm Booss-Bavnbek
Martin Bohle-Carbonell

140/87 "ON THE TOPOLOGY OF SPACES OF HOLOMORPHIC MAPS"

By: Jens Gravesen

141/87 "RADIOMETERS UDVIKLING AF BLODGASAPPARATUR -
ET TEKNOLOGIHISTORISK PROJEKT"

Projektrapport af Finn C. Physant
Vejleder: Ib Thiersen

142/87 "The Calderón Projektor for Operators With
Splitting Elliptic Symbols"

by: Bernhelm Booss-Bavnbek og
Krzysztof P. Wojciechowski