

Measuring and modelling dry deposition of ammonia to deciduous forest using high temporal and spatial resolution techniques

Hansen, Kristina; Sørensen, Lise Lotte; Hertel, Ole; Andersen, Helle Vibeke; Geels, Camilla; Jensen, Bjarne; Hildan, Kaj Morten; Søren, Lund; Boegh, Eva

Publication date:
2011

Document Version
Early version, also known as pre-print

Citation for published version (APA):
Hansen, K., Sørensen, L. L., Hertel, O., Andersen, H. V., Geels, C., Jensen, B., Hildan, K. M., Søren, L., & Boegh, E. (2011). *Measuring and modelling dry deposition of ammonia to deciduous forest using high temporal and spatial resolution techniques*. Poster session presented at Nitrogen and Global Change, Edinburgh, United Kingdom. <http://www.nitrogen2011.org/>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

Measuring and modelling dry deposition of ammonia to deciduous forest using high temporal and spatial resolution techniques

Kristina Hansen^{1,2}, Lise L. Sørensen², Ole Hertel^{1,2}, Helle V. Andersen², Camilla Geels², Bjarne Jensen², Morten K. Hildan², Søren W. Lund³, and Eva Boegh¹

1. Department of Environmental, Social and Spatial Change, Roskilde University, P.O. Box 260, Universitetsvej 1, DK-4000 Roskilde. 2. National Environmental Research Institute, Aarhus University, P.O. Box 358, Frederiksborgvej 399, DK-4000 Roskilde. 3. Risø National Laboratory for Sustainable Energy, Technical University of Denmark, P.O. Box 49, Frederiksborgvej 399, DK-4000 Roskilde
krih@dmu.dk

Aim

- To compare vertical NH₃ concentrations and fluxes measured above deciduous forest using Conditional Time Average Gradient (COTAG) and Relaxed Eddy Accumulation (REA).
- To examine the performance of the Danish applied local-scale deposition model OML-DEP on calculating dry deposition of NH₃ to deciduous forest, by comparing calculations with new flux measurements.

Conclusion

- Measurements and model calculations are in good agreement on estimating atmospheric NH₃ concentration.
- REA indicate a clearly bi-directional NH₃ flux while fluxes determined using COTAG are non significant and associated with high uncertainty.
- There is a further need for improving the description of dry deposition in local-scale models whereby the NH₃ vegetative emission and its contribution to the atmospheric NH₃ flux is considered.


Introduction

Calculations of ammonia (NH₃) deposition to Danish nature resorts indicate exceedance of critical loads for N deposition particular to forests (ref.1). A continuous high N load in terrestrial ecosystems can cause critical effects to the biodiversity (ref. 3).


Modeling dry deposition processes of NH₃ above forest is challenging due to the complex forest-atmosphere interactions along with a lacking knowledge of vertical NH₃ fluxes between vegetative surfaces and the atmosphere. Therefore further measurements of fluxes are still needed to improve the current knowledge of NH₃ fluxes and to verify the local-scale deposition models.

Method

Vertical fluxes of NH₃ was measured above a beech forest (*Fagus sylvatica*) in Denmark from 10 August to 11 November 2010 using the long-term COTAG technique (29.8 m and 34 m) and the high time resolution technique REA (33 m). Calculations of concentration and dry deposition are performed using the local-scale deposition model (OML-DEP) applied in DAMOS (ref. 2). The DAMOS calculations are based on state-of-the-art emission inventories with hourly time resolution and a spatial resolution down to single farm level (ref. 4).




Concentrations




NH₃ concentration above Lille Bøgeskov measured using COTAG Denuder (29.8 m) in four sampling periods and REA WEDD (33 m) only in the fourth sampling period and calculated by OML-DEP through the full measuring period.

OML-DEP calculations indicate an overall decreasing trend in the measuring period. These results are in good agreement with the concentration measured using COTAG that indicates mean concentrations decreasing from 1.1 to 0.7 µg NH₃-N m⁻³.

REA WEDD measurements are only performed in the fourth sampling period and indicate an increasing concentration in the end of the measuring period, but the mean concentrations assessed by the three different techniques are all in good agreement through the fourth sampling period and deviate only by 0.56 to 0.68 µg NH₃-N m⁻³ from each other.



Fluxes





NH₃ fluxes above Lille Bøgeskov for 10 August – 11 November 2010 along with leaf area index (LAI) and CO₂ fluxes.

High temporal resolution measurements of NH₃ flux using REA indicate a bidirectional flux that could be related to LAI. After defoliation the NH₃ flux begins changing direction from deposition to emission equally to the pattern of CO₂ fluxes.

OML-DEP doesn't take natural contributions as the NH₃ vegetative emission into account in calculating NH₃ deposition fluxes. Furthermore, the calculated flux seems underestimated due to an underestimation of the turbulent transport to the canopy.

The conditional COTAG fluxes indicate non significant deposition fluxes and the comparable conditional averaged REA fluxes indicate emission fluxes. Though the uncertainty on the estimated COTAG fluxes indicate that the flux could be oppositely directed.



The measurement site (Lille Bøgeskov, Sorø) in Denmark in January 2011.

References

1. Frohn et al. (2008), Nitrogen load of nature areas in Eastern Jutland (In Danish: Kvalstofbelastning af naturområder i Østjylland), National Environmental Research Institute, University of Aarhus, Roskilde, Denmark.
2. Olesen (2005), International J. Environment and Pollution, 5, 412-417.
3. Stevens et al. (2004), Science, 303, 1876-1879.
4. Skjøth et al. (2011), ACPD, 11, 1-37.

Acknowledgement

The NitroEurope IP project (0174841-2) has supported this project financial.