

Dynamics of dissolved organic matter in a shallow Danish estuary

Nielsen, Søren Laurentius; Pedersen, Troels Møller; Markager, Stiig; Sand-Jensen, Kaj

Publication date:
2013

Document Version
Peer reviewed version

Citation for published version (APA):
Nielsen, S. L., Pedersen, T. M., Markager, S., & Sand-Jensen, K. (2013). *Dynamics of dissolved organic matter in a shallow Danish estuary*.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

DYNAMICS OF DISSOLVED ORGANIC MATTER IN A SHALLOW DANISH ESTUARY

Søren Laurentius Nielsen, ENSPAC, Roskilde University, Denmark
Troels Møller Pedersen, ENSPAC, Roskilde University, Denmark
Stiig Markager, Aarhus University, Denmark
Kaj Sand-Jensen, University of Copenhagen, Denmark




This study presents data on dissolved organic matter, collected through an annual cycle in a shallow Danish estuary, Roskilde Fjord.


The different DOM fractions initially appeared to behave conservatively when evaluated against salinity. However, isotopic mixing models based on stable isotope ratios of carbon and nitrogen reveal that conservative mixing models based on DOM concentration and salinity mask estuarine processes altering the DOM pool.

Our data suggest an accumulation of DON especially in the inner parts of the estuary potentially as release from both sediment and phytoplankton.


Roskilde Fjord, Denmark, is microtidal, and salinity varies from about 7 psu in the innermost part to some 23 psu at the mouth, opening to Kattegat. Roskilde Fjord receives freshwater input from several smallish streams along the whole estuary




All DOM fractions – DOC, DON and CDOM (expressed as a_{375}) would initially appear to be conservatively mixed – i.e. no production or decomposition in Roskilde Fjord. Note initial disappearance of 20 – 30 % of DOC and CDOM




The $\delta^{13}\text{C}$ -DOM mixing curve (A and B) follows a non-conservative isotope mixing both in autumn and in spring. The $\delta^{15}\text{N}$ -DOM mixing curve shows ^{15}N depletion in autumn at the inner stations while the two outer stations follow the expected conservative distribution (C). In spring, DON concentrations at all stations are depleted compared to the expected conservative distribution.



Average estuarine $\delta^{15}\text{N}$ -DOM and $\delta^{13}\text{C}$ -DOM values from autumn show that DOM at the inner station (E1) has the highest terrestrial signal. DOM at St. 2, on the other hand, is depleted in ^{13}C which could be explained by autochthonous DOC production. St. 3 and 4 had similar $\delta^{13}\text{C}$ -DOM values but the DOM at St. 4 was more depleted of ^{15}N . During spring the pattern changed and the isotope composition of DOM at St. 1 and St. 2 (E1 and E2) was enriched in $\delta^{13}\text{C}$ and depleted of $\delta^{15}\text{N}$ compared with autumn averages.



Schematic relationship between the ratio of DIN:DOC in loadings and the autochthonous to allochthonous DOC ratio (auDOC:alDOC) illustrating the potential effect of residence time and production rate. The line represents the ratio assuming that autochthonous DOC is produced from DIN at the Redfield ratio (C:N=6.2). The annual average of DIN:DOC loadings for Horsens Fjord and Roskilde Fjord are 0.45 and 0.19 (mol basis), respectively. This is equal to a auDOC:alDOC ratio at Redfield (*) of 2.68 for Horsens Fjord and 1.22 for Roskilde Fjord. Estimated auDOC:alDOC are approximately 4 in both Roskilde Fjord and Horsens Fjord placing them above the Redfield line (□)



The mechanisms controlling concentrations of DON, DOC and CDOM seem to be dilution most of the year (conservative mixing). However, isotopic mixing models reveal that conservative mixing models based on DOM concentration and salinity mask estuarine processes altering the DOM pool. Isotope analysis of DOM reveals a high terrestrial/refractory signal in November-December whereas biochemical processes alter the DOM pool in spring.