

HOXB4 Gene Expression Is Regulated by CDX2 in Intestinal Epithelial Cells

Jørgensen, Steffen; Coshun, Mehmet; Mikkelsen Homburg, Keld; Pedersen, Ole B.V.; Troelsen, Jesper

Published in:
P L o S One

DOI:
[10.1371/journal.pone.0164555](https://doi.org/10.1371/journal.pone.0164555)

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Jørgensen, S., Coshun, M., Mikkelsen Homburg, K., Pedersen, O. B. V., & Troelsen, J. (2016). HOXB4 Gene Expression Is Regulated by CDX2 in Intestinal Epithelial Cells. *P L o S One*, 11(10), Article 0164555. <https://doi.org/10.1371/journal.pone.0164555>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

RESEARCH ARTICLE

HOXB4 Gene Expression Is Regulated by CDX2 in Intestinal Epithelial Cells

Steffen Jørgensen^{1,2}, Mehmet Coskun³, Keld Mikkelsen Homburg², Ole B. V. Pedersen², Jesper T. Troelsen^{1*}

1 Department of Science and Environment, Roskilde University, Roskilde, Denmark, **2** Department of Clinical Immunology, Naestved Hospital, Naestved, Region Zealand, Denmark, **3** Department of Gastroenterology, Medical Section, Herlev Hospital, University of Copenhagen, Copenhagen, Denmark

* troelsen@ruc.dk



OPEN ACCESS

Citation: Jørgensen S, Coskun M, Homburg KM, Pedersen OBV, Troelsen JT (2016) *HOXB4* Gene Expression Is Regulated by CDX2 in Intestinal Epithelial Cells. PLoS ONE 11(10): e0164555. doi:10.1371/journal.pone.0164555

Editor: Shrikant Anant, University of Kansas School of Medicine, UNITED STATES

Received: June 23, 2016

Accepted: September 6, 2016

Published: October 18, 2016

Copyright: © 2016 Jørgensen et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: This work was funded by the Local Research Foundation, Hospital South, Naestved, Region Zealand; a Regional Zealand Research Grant; Fonden til Lægevidenskabens Fremme, and by the Danish Research Council (FSS).

Competing Interests: The authors declare no competing interests.

Abstract

The mammalian *Caudal*-related homeobox transcription factor 2 (CDX2) plays a key role in the homeobox regulatory network and is essential in regulating the expression of several homeobox (*HOX*) genes during embryonic development, particularly in the gut. Genome-wide CDX2 chromatin immunoprecipitation analysis and expression data from Caco2 cells also suggests a role for CDX2 in the regulation of *HOXB4* gene expression in the intestinal epithelium. Thus, the aim of this study was to investigate whether *HOXB4* gene expression is regulated by CDX2 in the intestinal epithelium. We demonstrated binding of CDX2 to four different CDX2 binding sites in an enhancer region located upstream of the *HOXB4* transcription start site. Mutations in the CDX2 binding sites reduced *HOXB4* gene activity, and knock down of endogenous CDX2 expression by shRNA reduced *HOXB4* gene expression. This is the first report demonstrating the CDX2 regulation of *HOXB4* gene expression in the developed intestinal epithelium, indicating a possible role for *HOXB4* in intestinal homeostasis.

Introduction

Homeobox (*HOX*) proteins are important regulators during development and the formation of the anterior and posterior body plane in vertebrates. The 39 different *HOX* genes in humans can be divided into four different clusters, *HOXA*, *HOXB*, *HOXC* and *HOXD*, which are located on different chromosomes[1]. Each cluster contains 9–11 *HOX* genes. *HOX* genes are transcription factors belonging to the *HOX* protein family that share a DNA-binding domain, termed the *HOX* domain. The *HOX* domain is comprised of 60 amino acids that bind DNA in a sequence-dependent manner[2]. In adults, *HOX* gene expression is organ-specific, and the deregulation of specific *HOX* genes has been associated with cancer development[3]; these genes include *HOXA9*, which has been implicated in the development of acute myeloid leukemia[4], *HOXD3*, which increases invasiveness and metastasis in lung cancer[5], and both *HOXA4* and *HOXD10*, which are overexpressed in colorectal cancer[6].

Another member of the *HOX* family is *HOXB4*, which is expressed in different tissues such as colon and liver[7] and is a possible marker in cervical cancer[8]. In hematopoietic stem cells,

overexpression of HOXB4 increases proliferation, and expression is regulated by the transcription factor GATA-2[9]. However, despite the central role of HOXB4 in these processes, the regulation of HOXB4 is still poorly understood.

The *Caudal*-related homeobox 2 (CDX2) gene is specifically expressed early after fertilization in the trophectoderm and is important for the correct formation of the blastocyst and subsequent implantation in the uterus[10]. During later development, the CDX2 gene is specifically expressed posteriorly in the tailbud. The gene encoding the transcription factor CDX2 is also a member of the paralogous gene family. DNA interactions mediated by the homeodomain of CDX2 can activate gene expression in a manner similar to the HOX proteins. CDX2 also plays an important role during development and establishment of the gastro-intestinal tract, and loss of CDX2 during development is fatal[11]. Later in life, CDX2 expression is limited to the intestinal epithelium and plays an important role in maintaining the crypt/villus axis and cellular differentiation along the axis[12]. Loss of Cdx2 expression in the mouse intestinal epithelium inhibits the development of enterocytes and is fatal, underscoring the importance of CDX2 in intestinal homeostasis[13]. Moreover, CDX2 acts as a tumor suppressor, and inhibition of CDX2 expression increases the invasiveness of colon cancer[14]. We conducted the study presented here to determine whether CDX2 is involved in the gene expression of HOXB4.

Materials and Methods

Cell lines

SW480 (ATCC: CCL-228), Caco2 (ATCC: HTB-37) and HEK293 (ATCC: CRL-1573) cells were cultured as monolayers in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% heat inactivated fetal calf serum (FCS), 100 µg/ml streptomycin and 100 U/ml penicillin. All cell lines were cultured in a humidified atmosphere under 5% CO₂.

Viral transduction and CDX2 knockdown

An SW480 cell line with stable CDX2 knock down was created using the lentiviral pLKO.1 puro shRNA expression vector. The shRNA expression vector (pLKO.1 puro) was a kind gift from Dr. Bob Weinberg (Plasmid #8453, Addgene, Cambridge, MA, USA)[15]. pLKO.1 puro was digested with *AgeI* and *EcoRI*, and the DNA oligos CDX2 shRNA 735 Forward and CDX2 shRNA 735 Reverse were hybridized and inserted into pLKO.1. A lentiviral control vector containing scrambled non-target shRNA (pLKO.1 scramble) was used as a negative control and was a kind gift from Dr. David Sabatini (Plasmid #1864, Addgene)[16]. Next, 6x10⁶ HEK293 cells were plated in 6-well plates one day prior to transfection. HEK293 were co-transfected with either scrambled shRNA or CDX2 shRNA expressing lentivirus, psPAX, which was a gift from Malin Parmar (Plasmid #35002, Addgene), and pMD2.G, which was a gift from Didier Trono (Plasmid #12259, Addgene). The transfected HEK293 cells were maintained for 48 hours in DMEM supplemented with heat inactivated 10% FCS. The supernatant was then sterile filtered using a 0.45-µm filter and added to 5x10⁵ SW480 cells, followed by transduction for 24 hours. After 24 hours, the medium was replaced with DMEM supplemented with 10% heat inactivated FCS and 10 µg/ml puromycin (Life Technologies, Carlsbad, CA, USA), and the cells were reseeded twice per week for 3 weeks. Thereafter, cells were reseeded twice per week and cultured in DMEM supplemented with 10% heat inactivated FCS. CDX2 knock down was verified by RT-qPCR and western blotting.

HOXB4 promoter/enhancer cloning

A potential CDX2 binding region in the *HOXB4* enhancer was identified using previously published chromatin immunoprecipitation-sequencing (ChIP-seq) data [17], which was uploaded to the UCSC Genome Browser (version hg18).

The BAC clone RP11-111C6, which covers chromosome 17, positions -48583316 to -48578320 (GENBANK ID: NC_000017.11), was obtained from the BACPAC Resource Centre (Oakland, CA, USA) and used as a template to amplify the *HOXB4* promoter and enhancer region. The *HOXB4* promoter and enhancer region was cloned using an In-Fusion™ cloning kit (Clontech, Saint-Germain-en-Laye, France). A 4997-bp fragment was amplified using Phusion HOTSTART II HD polymerase (Thermo Fisher Scientific, Waltham, MA, USA), the Infusion HOXB4 Enhancer/promoter forward primer and the Infusion HOXB4 reverse primer (Table 1). The pGL4.10 luciferase reporter plasmid (Promega, Nacika, Sweden) was digested with *HindIII* according to the manufacturer's protocol (Thermo Fisher Scientific), and the promoter and enhancer were inserted using the In-Fusion™ HD cloning kit (Clontech) according to the manufacturer's protocol (Clontech). A 2850-bp fragment excluding the up-stream CDX2 binding sites was amplified using Phusion HOTSTART II HD polymerase (Thermo

Table 1. DNA oligo sequences used in this study.

Name	Sequence
CDX2 shRNA 735 Forward	5' - CGGCAATATCGAGTGGTGTACACCTCGAGGTGTACACCACTCGATATTTGTTTTCG - 3'
CDX2 shRNA 735 Reverse	5' - AATTCAAAACAAATATCGAGTGGTGTACACCTCGAGGTGTACACCACTCGATATTTG - 3'
Infusion HOXB4 Promoter Reverse Primer	5' - CCGGATTGCCAAGCTGGTGGTGTAAATAAAGTCCTTTTCG - 3'
Infusion HOXB4 Enhancer/Promoter Forward Primer	5' - CTCGGCGGCCAAGCTATTGATGAGAACGATTCTTCGG - 3'
Infusion HOXB4 Enhancer/Promoter Reverse Primer	5' - CCGGATTGCCAAGCTTAATTTCTGGGAATTGCCACAAA - 3'.
Infusion HOXB4 Move Forward	5' - CTAGGCGAGTTCATA TATGTGTTTAGAAGGAAAGGAC - 3'
Infusion HOXB4 Move Reverse	5' - GTCTGCCTGACTGCCTGTGGTCAGATCGATTGGC - 3'
EMSA-CDX2-Site-1 Forward	5' - TCAGGCCTTTCTGGGTGCCT - 3'
EMSA-CDX2-Site-1 Reverse	5' - AAGGCACCCAGAAAGGCCTG - 3'
EMSA-CDX2-Site-2 Forward	5' - GGTGGTGTAAATAAAGTCCT - 3'
EMSA-CDX2-Site-2 Reverse	5' - AAGGACTTTTATTACACCAC - 3'
EMSA-CDX2-Site-3 Forward	5' - GGATATTTAATGACGGGCAA - 3'
EMSA-CDX2-Site-3 Reverse	5' - ATTGCCCGTCATTAAATATC - 3'
EMSA-CDX2-Site-4 Forward	5' - TGAGGAACCATAAAAGATGC - 3'
EMSA-CDX2-Site-4 Reverse	5' - TGCATCTTTTATGGTTCCTC - 3'
EMSA Sucrase-isomaltase probe Forward	5' - ACTGACTTTATTAACCTTTGTGACC - 3'
EMSA Sucrase-isomaltase probe Reverse	5' - AGGTCACAAAGTTAATAAAGTCAG - 3'
EMSA unspecific probe Forward	5' - AACGTAGCTGATCGAATCGGTTAC - 3'
EMSA unspecific probe Reverse	5' - AGTAACCGATTTCGATCAGCTACGT - 3'
HOXB4 ChIP PCR Forward	5' - GCCAGATGATATAGGGCCCC - 3'
HOXB4 ChIP PCR Reverse	5' - AAGGTGGGAAATGGGGTTC - 3'
HOXB4 qPCR Forward	5' - CTGGATGCGCAAAGTTCAC
HOXB4 qPCR Reverse	5' - AGCGGTGTAGTGAAATTCCTT - 3'
CDX2 qPCR Forward	5' - ACTACAGTCGCTACATCACCA - 3'
CDX2 qPCR Reverse	5' - GAAGACACCGACTCAAGGG - 3'
RPLP0 qPCR Forward	5' - GCAATGTTGCCAGTGTCTG - 3'
RPLP0 qPCR Reverse	5' - GCCTTGACCTTTTCAGCAA - 3'
HOXB4 CDX2 ChIP PCR Forward	5' - GCCAGATGATATAGGGCCCC - 3'
HOXB4 CDX2 ChIP PCR Reverse	5' - AAGGTGGGAAATGGGGTTC - 3'

doi:10.1371/journal.pone.0164555.t001

Fisher Scientific), the Infusion HOXB4 promoter forward primer and the Infusion HOXB4 promoter reverse primer (Table 1). The pGL4.10 luciferase reporter plasmid (Promega) was digested with *HindIII* according to the manufacturer's protocol (Fisher Scientific), and the HOXB4 promoter was inserted using the In-Fusion™ HD cloning kit (Clontech) according to the manufacturer's protocol (Clontech).

CDX2 binding site mutations

Four different constructs containing the HOXB4 promoter/enhancer region (4997-bp fragment) and in which each of the CDX2 binding sites, sequence a/cATAAAa/t[18], were replaced with the *EcoRI* cleavage sequence, GAATTCG, were synthesized by Eurofins (Eurofins MWG Operon, Ebersberg, Germany). The HOXB4 pGL4.10 reporter plasmid was digested with *NdeI* (Thermo Fisher Scientific) and *PstI* (Thermo Scientific) according to the manufacturer's protocol. Each of the four CDX2 mutated constructs was used as template, and a 1724-bp fragment containing the mutated CDX2 binding site was amplified using Phusion HotStart II polymerase (Thermo Fisher Scientific), following the manufacturer's protocol, and the Infusion HOXB4 Move forward and reverse primers (Table 1). The fragments were inserted in the *NdeI*- and *PstI*-digested HOXB4 pGL4.10 reporter plasmid using the In-Fusion™ HD cloning kit (Clontech) according to the manufacturer's instructions, resulting in four different plasmids with each of the CDX2 binding sites mutated.

Reporter gene assays

One day prior to transfection, SW480 cells were seeded in 24-well plates at a density of 5×10^4 cells/well and transiently transfected the following day. The cells were transfected using polyethylenimine at a final concentration of 2 μ M and 0.3 μ g DNA/well. Twenty-five nanograms of luciferase reporter gene plasmid and 50 ng of β -galactosidase expression plasmid were used per well, and the DNA concentration was adjusted to 0.3 μ g with pBluescript SK+ (Stratagene, La Jolla, CA, USA). Forty-eight hours post-transfection, cells were harvested and lysed; luciferase and β -galactosidase activities were determined using the Dual Light™ luminescent reporter gene assay (Applied Biosystems, Naerum, Denmark), and luminescence was determined using a GloMax® 96 Microplate Luminometer (Promega) according to the manufacturer's instructions.

Protein extraction and immunoblotting

Total proteins were extracted from the lentiviral-modified shRNA-expressing SW480 cells (scrambled shRNA or CDX2 shRNA) using Trizol (Life Technologies) as described by the manufacturer. Ten micrograms of total protein were analyzed by SDS-PAGE using a NuPAGE 4–12% Bis-Tris protein gel (Life Technologies) followed by transfer to a PVDF membrane via electroblotting using the iBlot Western blotting system (Life Technologies). The membrane was incubated with primary antibodies against human CDX2 (1:300, mouse monoclonal, Bio-Genex Laboratories, Fremont, CA) or glyceraldehyde 3-phosphate dehydrogenase (GAPDH, 1:300, mouse monoclonal, Millipore, Darmstadt, Germany).

Electrophoretic mobility shift assay

The amplified region of the HOXB4 promoter and enhancer was analyzed using the JASPAR database with a relative profile score threshold setting of 80% [19]. Four different probe sets were designed to cover CDX2 binding sites in the HOXB4 enhancer at position -2662 to -2672 (site-4) on the minus strand, position -2737 to -2747 (site-3) on the plus strand, position -2836 to -2846 (site-2) on the minus strand, and position -3030 to -3040 (site-1) on the plus

strand. The electrophoretic mobility shift assay (EMSA) was performed as previously described [20]. Briefly, nuclear extracts from confluent Caco2 cells were incubated with p32-labeled probes for site-1, -2, -3, or -4. A competition assay was performed using either an unlabeled sucrose isomaltase probe containing a known CDX2 binding site (positive control) or unlabeled HOXB4 probes corresponding to each of the four identified potential CDX2 binding sites (site-1, -2, -3-, or -4). A supershift assay was performed using a human CDX2 antibody (BioGenex) or influenza hemagglutinin (HA) as a negative control (described below).

RNA extraction and RT-qPCR

Total RNA was extracted from SW480 cells using an E.Z.N.A. Total RNA KIT I (OMEGA BIOTEK, Norcross, GA). RNA extraction was performed according to the manufacturer's instructions. cDNA was synthesized according to the manufacturer's instructions, using qSCRIPT cDNA Supermix (Quanta Biosciences, Gaithersburg, MD). Relative quantification was calculated using the $\Delta\Delta C_T$ method, and ribosomal protein large P0 (RPLP0) was used as a reference gene[21].

Chromatin immunoprecipitation

Confluent Caco2 cells were cross-linked, sonicated, and immunoprecipitated with an antibody specific for either human CDX2 (BioGenex) or an antibody specific for the influenza hemagglutinin (HA) epitope (rabbit polyclonal α -HA; Santa Cruz Biotechnology Inc., Heidelberg, Germany) as a negative control, as previously described[22]. Purified immunoprecipitated DNA was compared to input DNA, which corresponded to non-immunoprecipitated sheared cross-linked chromatin, via qPCR analysis. The primers used to amplify the human genomic sequences of *HOXB4* are listed in Table 1, and quantification of the chromatin immunoprecipitation (ChIP)-DNA was performed using qPCR in triplicate with a Stratagene MX3002 (Stratagene) and a QuantiTect SYBR Green PCR Kit (Qiagen, Duesseldorf, Germany). Enrichment was calculated as described elsewhere[23].

Statistical analysis

Statistical analysis was performed using unpaired Student's *t*-tests, and values are shown as the mean and standard deviation (SD). P-values of less than 0.05 were considered significant.

Results

CDX2 binds to the *HOXB4* enhancer

Genome-wide CDX2 ChIP-seq data from a published report[17] on the intestinal cell line Caco2 revealed a potential CDX2 binding region in the *HOXB4* enhancer, located from 2640 to 3050 bp upstream of the *HOXB4* transcription start site (Fig 1A). Thus, to verify whether CDX2 binds to this region, we performed qPCR on chromatin obtained from Caco2 cells immunoprecipitated with an anti-CDX2 or an anti-HA (negative control) antibody. In ChIP assays, chromatin obtained from anti-CDX2 antibody immunoprecipitates showed significant enrichment of the *HOXB4* enhancer (49-fold, $P < 0.05$) compared to the negative control (Fig 1B).

HOXB4 expression is modulated by CDX2

To determine whether endogenous *HOXB4* expression is regulated by CDX2, we performed stable shRNA-mediated knock down of endogenous *CDX2* in SW480 cells. This cell line was chosen because we were unable to generate a Caco2 cell line with stable *CDX2* knock down, a limitation that has also been reported by other groups[18]. In SW480 cells, endogenous CDX2




Fig 1. Illustration of the CDX2 binding region in the *HOXB4* enhancer region and analysis of CDX2 interactions with the *HOXB4* enhancer region. Fig 1A) Genome browser CDX2 ChIP-seq peaks from a dataset generated by Boyd et al. [17]. Fig 1B) ChIP-PCR with primers covering the CDX2 binding region within the *HOXB4* enhancer. Equal amounts of input DNA were used for CDX2 ChIP-PCR. Hemagglutinin (HA) was used as a negative control. The mean ($n = 4$) and SD are shown. * $P < 0.05$.

doi:10.1371/journal.pone.0164555.g001

expression was effectively knocked down (Fig 2A). Moreover, both *CDX2* and *HOXB4* mRNA levels were decreased in SW480^{CDX2} compared to SW480^{NEG} cells (data not shown), indicating that CDX2 is involved in maintaining the endogenous expression of *HOXB4* in SW480 cells.

To verify the possible role of CDX2 in the regulation of *HOXB4* expression, a luciferase reporter construct containing the human *HOXB4* enhancer and promoter as well as another construct containing only the *HOXB4* promoter were created. The activity of the reporter constructs containing the *HOXB4* promoter or *HOXB4* promoter and enhancer were measured both in SW480^{CDX2} and SW480^{NEG} cells. The *HOXB4* promoter exhibited activity on its own when comparing the expression levels in SW480^{CDX2} and SW480^{NEG} cells (Fig 2B). The activity of the luciferase construct containing both the *HOXB4* enhancer and promoter was significantly reduced ($P < 0.01$) when examining the expression levels in SW480^{CDX2} compared to SW480^{NEG} cells (Fig 2B). This result indicates that CDX2 is involved in the transcriptional regulation of *HOXB4*.

To further investigate the role of CDX2 in *HOXB4* expression, we performed a co-transfection experiment in SW480 cells. The *HOXB4* enhancer and promoter reporter construct was




Fig 2. The effects of *CDX2* knock down and overexpression on *HOXB4* expression in the SW480 cell line and functional analysis of 4 different *CDX2* binding sites in the *HOXB4* enhancer region. Fig 2A) Western blot analysis of *CDX2* expression levels in SW480 cells treated with scrambled shRNA and cells in which *CDX2* was knocked down. GAPDH was used as a control to analyze equal amounts of total cellular protein. Fig 2B) The effects of *CDX2* knock down on the promoter activities of reporter constructs containing the *HOXB4* promoter alone or the *HOXB4* promoter including the enhancer region. Fig 2C) Transient transfection of the *HOXB4* promoter and enhancer reporter construct in SW480 cells with and without co-transfection of a *CDX2* expression plasmid. The mean (n = 4) and SD are shown. * $P < 0.05$ and ** $P < 0.01$

doi:10.1371/journal.pone.0164555.g002

transiently transfected into SW480 cells with or without the *CDX2* expression plasmid. When overexpressing *CDX2*, luciferase activity driven by the *HOXB4* enhancer and promoter construct was significantly higher (2.4-fold, $P < 0.01$) relative to cells without co-transfection of the *CDX2* expression vector (Fig 2C).

CDX2 interacts with the *HOXB4* enhancer region

We used computational DNA sequence analysis to identify potential *CDX2* binding sites in the *HOXB4* enhancer region. This analysis revealed four predicted and potential *CDX2* binding sites within the *HOXB4* enhancer region (Fig 3A). To further analyze the function of each of the four *CDX2* binding sites, mutations were introduced into each *CDX2* binding site (*HOXB4* site-1, -2, -3, and -4). The enhancer activity of each construct was measured using transient transfection in SW480 cells, and mutation of all four sites individually significantly reduced *HOXB4* enhancer and promoter luciferase activity ($P < 0.01$ for all) compared to the wild type reporter construct (Fig 3B). When comparing the reduction in enhancer activity between the four *CDX2* binding sites, mutation of *HOXB4* site 2 reduced the activity to a lesser extent compared to *HOXB4* site-1, -3 and -4. ($P < 0.05$)

In addition to the functional analysis of the *HOXB4* enhancer/promoter constructs containing mutations within the *CDX2* binding sites (*HOXB4* site-1, -2, -3 or -4), we performed EMSA with oligonucleotides covering each of the four potential *CDX2* binding sites and nuclear extracts from Caco2 cells (Fig 3C–3F). The data revealed that all four probes were able to induce the formation of a *CDX2*/DNA complex (Fig 3C–3F, lane 1). However, two non-specific complexes for site-3 and one non-specific complex for site-4 were revealed (Fig 3E and 3F, lane 1, respectively). The binding specificity of these complexes was examined in competition assays in which excess unlabeled cold probes, specifically a sucrose isomaltase probe, a probe containing the *CDX2* consensus sequence, or probes containing *CDX2* binding site sequences (*HOXB4* site-1, -2, -3 or -4), or an unspecific oligonucleotide were utilized. Unlabeled *HOXB4* site-2 and site-4 probes completely out-competed *CDX2*/DNA complexes (Fig 3C, lane 4, and Fig 3E, lane 3). However, cold *HOXB4* site-1 and site-3 probes only partially out-competed the *CDX2*/DNA complex (Fig 3B, lane 2, and Fig 3D, lane 4). In contrast, the formation of the *CDX2*/DNA complex was completely eliminated in all EMSAs by competition with the




Fig 3. Functional and EMSA analyses of CDX2 binding sites in the *HOXB4* enhancer region. Fig 3A) Positions of the four potential CDX2 binding sites and the EMSA probes covering these sites. Fig 3B) Reporter gene assays with four different luciferase constructs containing the *HOXB4* enhancer with wild-type CDX2 binding sites or four distinct reporter constructs containing individual mutations in the four CDX2 binding sites (*HOXB4* site-1, -2, -3 or 4). The mean ($n = 4$) and SD are shown. * $P < 0.05$ and ** $P < 0.01$. Fig 3C–3F) EMSAs with

probes for the four different CDX2 binding sites (HOXB4 site-1, -2, -3 and 4). One hundred-fold unlabeled probe was added to the competition reactions as well as sucrose isomaltase (SI) or unspecific probe (unspec). Influenza hemagglutinin (HA) antibody (negative control) or CDX2 antibody (CDX2) was used in the supershift assay. Non-specific bands and supershifts indicated by black arrows.

doi:10.1371/journal.pone.0164555.g003

unlabeled sucrose isomaltase probe (Fig 3B–3D, lane 3, and Fig 3E, lane 4), whereas the unspecific probe did not affect CDX2/DNA complex formation (Fig 3C, lane 4, and Fig 3D and 3E, lane 2).

To verify that CDX2 is present in the CDX2/DNA complex, supershift assays were performed with anti-CDX2 and anti-HA (negative control) antibodies. The addition of an antibody against CDX2 effectively supershifted the CDX2/DNA complex in site-2, -3, and -4 (Fig 3C–3E, lane 5) but failed to supershift the CDX2/DNA complex in site-1 (Fig 3B, lane 6), although a partial reduction in the band intensity of the CDX/DNA complex was observed. Moreover, as expected, the CDX2/DNA complex did not supershift with the addition of an antibody against HA (Fig 3C, lane 5, and Fig 3D and 3E, lane 6).

Discussion

The regulation of *HOXB4* expression is poorly understood in the intestinal epithelium and, until now, no other transcription factors in the intestinal epithelium have been shown to directly induce *HOXB4* expression. Here, we demonstrate that CDX2 regulates *HOXB4* gene expression in the intestinal epithelium-derived cell line SW480. Induction of *HOXB4* expression has previously been studied during fetal development, and expression is induced in the hindbrain by retinoic acid. The induction of *HOXB4* expression is mediated by the retinoic acid response element located within the *HOXB4* enhancer[24]. *HOXB4* is also expressed during hematopoiesis and regulated by the transcription factor GATA-2 in hematopoietic stem cells and progenitor cells[9].

A reduction in *HOXB4* expression levels, achieved using siRNA-mediated CDX2 knock down in the COLO320 cell line, has previously been demonstrated[25]. In the present study, we made similar observations in two different cell lines, Caco2 and SW480, both derived from colorectal cancers. However, Salari et al. did not investigate the possible CDX2-mediated induction of *HOXB4*. We demonstrate CDX2 binding to the *HOXB4* enhancer region by ChIP-PCR and we have identified four CDX2 binding sites and performed a functional analysis of each site. Mutating each of the 4 sites significantly reduces enhancer activity compared to controls, but not equally. Mutating the *HOXB4* site-2 affects promoter activity to a lesser extent compared to the mutation of *HOXB4* site-1, *HOXB4* site-3 and *HOXB4* site-4.

Previous studies of CDX1, 2 and 4 have shown that they bind to the promoter of *HOXA5*, and it has been suggested that CDX4 modulates *HOXA5* expression during embryonic development[26]. CDX2 expression along the crypt-villus axis varies, with the lowest CDX2 expression at the bottom of the crypt and increased expression during cellular differentiation[27]. The expression of *HOXA5* also varies and increases during differentiation[28], and it is possible that the expression is mediated by CDX2. *HOXA5* has been suggested to be a potential tumor suppressor in breast cancer[29]. CDX has also been shown to regulate *HOXB8* expression during embryonic development. Furthermore, *in vitro* binding of CDX1, CDX2 and CDX4 has been described, and binding has been demonstrated by all three CDX transcription factors *in vivo*[30].

HOXB4 binding sites have been investigated in hematopoietic stem cells and progenitor cells, and *HOXB4* has been shown to both up- and down-regulate the expression of target genes[31]. However, the binding of *HOXB4* is dependent not only on *HOXB4* but also on several different co-factors. These co-factors form complexes with *HOB4* and are members of the

TALE family, including MEIS, PBX and PREP, which play important roles in target sequence selection and binding[32]. However, the intestinal expression levels of these co-factors remain unclear and must be investigated to identify HOXB4 target genes in the intestinal epithelium.

The function of the CDX2-mediated induction of HOXB4 in intestinal differentiation and homeostasis is not known. As *HOX* genes are transcription factors, their downstream targets must be identified to clarify the function of HOXB4 expression in the intestinal epithelium. It is possible that HOXB4 induction plays a role in cellular differentiation or intestinal homeostasis.

We show that CDX2 binds to the HOXB4 enhancer, regulates HOXB4 expression in intestinal-derived epithelial cell lines, and has a possible role in the regulation of HOXB4 expression in the intestinal epithelium and intestinal homeostasis.

Acknowledgments

We would like to thank Christa Persson and Mette Juel Riisager for excellent technical assistance

Author Contributions

Conceived and designed the experiments: SJ JT.

Performed the experiments: SJ MC.

Analyzed the data: SJ MC KH OP JT.

Contributed reagents/materials/analysis tools: SJ MC KH OP JT.

Wrote the paper: SJ MC KH OP JT.

References

1. Yu YY, Pan YS, Zhu ZG. Homeobox genes and their functions on development and neoplasm in gastrointestinal tract. *European journal of surgical oncology: the journal of the European Society of Surgical Oncology and the British Association of Surgical Oncology*. 2007; 33(2):129–32. doi: [10.1016/j.ejso.2006.09.010](https://doi.org/10.1016/j.ejso.2006.09.010) PMID: [17045774](https://pubmed.ncbi.nlm.nih.gov/17045774/).
2. Svingen T, Tonissen KF. Hox transcription factors and their elusive mammalian gene targets. *Heredity*. 2006; 97(2):88–96. doi: [10.1038/sj.hdy.6800847](https://doi.org/10.1038/sj.hdy.6800847) PMID: [16721389](https://pubmed.ncbi.nlm.nih.gov/16721389/).
3. Takahashi Y, Hamada J, Murakawa K, Takada M, Tada M, Nogami I, et al. Expression profiles of 39 HOX genes in normal human adult organs and anaplastic thyroid cancer cell lines by quantitative real-time RT-PCR system. *Experimental cell research*. 2004; 293(1):144–53. PMID: [14729064](https://pubmed.ncbi.nlm.nih.gov/14729064/).
4. Ayton PM, Cleary ML. Transformation of myeloid progenitors by MLL oncoproteins is dependent on Hoxa7 and Hoxa9. *Genes & development*. 2003; 17(18):2298–307. doi: [10.1101/gad.1111603](https://doi.org/10.1101/gad.1111603) PMID: [12952893](https://pubmed.ncbi.nlm.nih.gov/12952893/); PubMed Central PMCID: PMC196466.
5. Hamada J, Omatsu T, Okada F, Furuuchi K, Okubo Y, Takahashi Y, et al. Overexpression of homeobox gene HOXD3 induces coordinate expression of metastasis-related genes in human lung cancer cells. *International journal of cancer Journal international du cancer*. 2001; 93(4):516–25. PMID: [11477555](https://pubmed.ncbi.nlm.nih.gov/11477555/).
6. Bhatlekar S, Addya S, Salunek M, Orr CR, Surrey S, McKenzie S, et al. Identification of a developmental gene expression signature, including HOX genes, for the normal human colonic crypt stem cell niche: overexpression of the signature parallels stem cell overpopulation during colon tumorigenesis. *Stem cells and development*. 2014; 23(2):167–79. doi: [10.1089/scd.2013.0039](https://doi.org/10.1089/scd.2013.0039) PMID: [23980595](https://pubmed.ncbi.nlm.nih.gov/23980595/); PubMed Central PMCID: PMC3887463.
7. Kanai M, Hamada J, Takada M, Asano T, Murakawa K, Takahashi Y, et al. Aberrant expressions of HOX genes in colorectal and hepatocellular carcinomas. *Oncology reports*. 2010; 23(3):843–51. PMID: [20127028](https://pubmed.ncbi.nlm.nih.gov/20127028/).
8. Barba-de la Rosa AP, Briones-Cerecero E, Lugo-Melchor O, De Leon-Rodriguez A, Santos L, Castelo-Ruelas J, et al. Hox B4 as potential marker of non-differentiated cells in human cervical cancer cells. *Journal of cancer research and clinical oncology*. 2012; 138(2):293–300. doi: [10.1007/s00432-011-1081-2](https://doi.org/10.1007/s00432-011-1081-2) PMID: [22120585](https://pubmed.ncbi.nlm.nih.gov/22120585/).

9. Fujiwara T, Yokoyama H, Okitsu Y, Kamata M, Fukuhara N, Onishi Y, et al. Gene expression profiling identifies HOXB4 as a direct downstream target of GATA-2 in human CD34+ hematopoietic cells. *PloS one*. 2012; 7(9):e40959. doi: [10.1371/journal.pone.0040959](https://doi.org/10.1371/journal.pone.0040959) PMID: [23028422](https://pubmed.ncbi.nlm.nih.gov/23028422/); PubMed Central PMCID: PMC3454409.
10. Strumpf D, Mao CA, Yamanaka Y, Ralston A, Chawengsaksophak K, Beck F, et al. Cdx2 is required for correct cell fate specification and differentiation of trophectoderm in the mouse blastocyst. *Development*. 2005; 132(9):2093–102. doi: [10.1242/dev.01801](https://doi.org/10.1242/dev.01801) PMID: [15788452](https://pubmed.ncbi.nlm.nih.gov/15788452/).
11. Deschamps J, van Nes J. Developmental regulation of the Hox genes during axial morphogenesis in the mouse. *Development*. 2005; 132(13):2931–42. doi: [10.1242/dev.01897](https://doi.org/10.1242/dev.01897) PMID: [15944185](https://pubmed.ncbi.nlm.nih.gov/15944185/).
12. Hryniuk A, Grainger S, Savory JG, Lohnes D. Cdx function is required for maintenance of intestinal identity in the adult. *Developmental biology*. 2012; 363(2):426–37. doi: [10.1016/j.ydbio.2012.01.010](https://doi.org/10.1016/j.ydbio.2012.01.010) PMID: [22285812](https://pubmed.ncbi.nlm.nih.gov/22285812/).
13. Verzi MP, Shin H, Ho LL, Liu XS, Shivdasani RA. Essential and redundant functions of caudal family proteins in activating adult intestinal genes. *Molecular and cellular biology*. 2011; 31(10):2026–39. doi: [10.1128/MCB.01250-10](https://doi.org/10.1128/MCB.01250-10) PMID: [21402776](https://pubmed.ncbi.nlm.nih.gov/21402776/); PubMed Central PMCID: PMC3133364.
14. Bonhomme C, Duluc I, Martin E, Chawengsaksophak K, Chenard MP, Kedinger M, et al. The Cdx2 homeobox gene has a tumour suppressor function in the distal colon in addition to a homeotic role during gut development. *Gut*. 2003; 52(10):1465–71. PMID: [12970140](https://pubmed.ncbi.nlm.nih.gov/12970140/); PubMed Central PMCID: PMC1773830.
15. Stewart SA, Dykxhoorn DM, Palliser D, Mizuno H, Yu EY, An DS, et al. Lentivirus-delivered stable gene silencing by RNAi in primary cells. *Rna*. 2003; 9(4):493–501. PMID: [12649500](https://pubmed.ncbi.nlm.nih.gov/12649500/); PubMed Central PMCID: PMC1370415. doi: [10.1261/rna.2192803](https://doi.org/10.1261/rna.2192803)
16. Sarbassov DD, Guertin DA, Ali SM, Sabatini DM. Phosphorylation and regulation of Akt/PKB by the rictor-mTOR complex. *Science*. 2005; 307(5712):1098–101. doi: [10.1126/science.1106148](https://doi.org/10.1126/science.1106148) PMID: [15718470](https://pubmed.ncbi.nlm.nih.gov/15718470/).
17. Boyd M, Hansen M, Jensen TG, Perearnau A, Olsen AK, Bram LL, et al. Genome-wide analysis of CDX2 binding in intestinal epithelial cells (Caco-2). *The Journal of biological chemistry*. 2010; 285(33):25115–25. doi: [10.1074/jbc.M109.089516](https://doi.org/10.1074/jbc.M109.089516) PMID: [20551321](https://pubmed.ncbi.nlm.nih.gov/20551321/); PubMed Central PMCID: PMC2919073.
18. Verzi MP, Hatzis P, Sulahian R, Philips J, Schuijers J, Shin H, et al. TCF4 and CDX2, major transcription factors for intestinal function, converge on the same cis-regulatory regions. *Proceedings of the National Academy of Sciences of the United States of America*. 2010; 107(34):15157–62. doi: [10.1073/pnas.1003822107](https://doi.org/10.1073/pnas.1003822107) PMID: [20696899](https://pubmed.ncbi.nlm.nih.gov/20696899/); PubMed Central PMCID: PMC2930576.
19. Berger MF, Badis G, Gehrke AR, Talukder S, Philippakis AA, Pena-Castillo L, et al. Variation in homeodomain DNA binding revealed by high-resolution analysis of sequence preferences. *Cell*. 2008; 133(7):1266–76. doi: [10.1016/j.cell.2008.05.024](https://doi.org/10.1016/j.cell.2008.05.024) PMID: [18585359](https://pubmed.ncbi.nlm.nih.gov/18585359/); PubMed Central PMCID: PMC2531161.
20. Coskun M, Boyd M, Olsen J, Troelsen JT. Control of intestinal promoter activity of the cellular migratory regulator gene ELMO3 by CDX2 and SP1. *Journal of cellular biochemistry*. 2010; 109(6):1118–28. doi: [10.1002/jcb.22490](https://doi.org/10.1002/jcb.22490) PMID: [20127720](https://pubmed.ncbi.nlm.nih.gov/20127720/).
21. Dydensborg AB, Herring E, Auclair J, Tremblay E, Beaulieu JF. Normalizing genes for quantitative RT-PCR in differentiating human intestinal epithelial cells and adenocarcinomas of the colon. *American journal of physiology Gastrointestinal and liver physiology*. 2006; 290(5):G1067–74. doi: [10.1152/ajpgi.00234.2005](https://doi.org/10.1152/ajpgi.00234.2005) PMID: [16399877](https://pubmed.ncbi.nlm.nih.gov/16399877/).
22. Coskun M, Olsen AK, Holm TL, Kvist PH, Nielsen OH, Riis LB, et al. TNF-alpha-induced down-regulation of CDX2 suppresses MEP1A expression in colitis. *Biochimica et biophysica acta*. 2012; 1822(6):843–51. doi: [10.1016/j.bbdis.2012.01.012](https://doi.org/10.1016/j.bbdis.2012.01.012) PMID: [22326557](https://pubmed.ncbi.nlm.nih.gov/22326557/).
23. Aparicio O, Geisberg JV, Struhl K. Chromatin immunoprecipitation for determining the association of proteins with specific genomic sequences in vivo. *Current protocols in cell biology / editorial board, Juan S Bonifacino [et al]*. 2004; Chapter 17:Unit 17 7. doi: [10.1002/0471143030.cb1707s23](https://doi.org/10.1002/0471143030.cb1707s23) PMID: [18228445](https://pubmed.ncbi.nlm.nih.gov/18228445/).
24. Folberg A, Nagy Kovacs E, Luo J, Giguere V, Featherstone MS. RARbeta mediates the response of Hoxd4 and Hoxb4 to exogenous retinoic acid. *Developmental dynamics: an official publication of the American Association of Anatomists*. 1999; 215(2):96–107. doi: [10.1002/\(SICI\)1097-0177\(199906\)215:2<96::AID-DVDY2>3.0.CO;2-T](https://doi.org/10.1002/(SICI)1097-0177(199906)215:2<96::AID-DVDY2>3.0.CO;2-T) PMID: [10373014](https://pubmed.ncbi.nlm.nih.gov/10373014/).
25. A clinical evaluation of the International Lymphoma Study Group classification of non-Hodgkin's lymphoma. The Non-Hodgkin's Lymphoma Classification Project. *Blood*. 1997; 89(11):3909–18. PMID: [9166827](https://pubmed.ncbi.nlm.nih.gov/9166827/).
26. Tabaries S, Lapointe J, Besch T, Carter M, Woollard J, Tuggle CK, et al. Cdx protein interaction with Hoxa5 regulatory sequences contributes to Hoxa5 regional expression along the axial skeleton.

- Molecular and cellular biology. 2005; 25(4):1389–401. doi: [10.1128/MCB.25.4.1389–1401.2005](https://doi.org/10.1128/MCB.25.4.1389-1401.2005) PMID: [15684390](https://pubmed.ncbi.nlm.nih.gov/15684390/); PubMed Central PMCID: PMC548006.
27. Olsen AK, Boyd M, Danielsen ET, Troelsen JT. Current and emerging approaches to define intestinal epithelium-specific transcriptional networks. *American journal of physiology Gastrointestinal and liver physiology*. 2012; 302(3):G277–86. doi: [10.1152/ajpgi.00362.2011](https://doi.org/10.1152/ajpgi.00362.2011) PMID: [22094602](https://pubmed.ncbi.nlm.nih.gov/22094602/).
 28. Wang Y, Hung C, Koh D, Cheong D, Hooi SC. Differential expression of Hox A5 in human colon cancer cell differentiation: a quantitative study using real-time RT-PCR. *International journal of oncology*. 2001; 18(3):617–22. PMID: [11179495](https://pubmed.ncbi.nlm.nih.gov/11179495/).
 29. Kikuyama M, Takeshima H, Kinoshita T, Okochi-Takada E, Wakabayashi M, Akashi-Tanaka S, et al. Development of a novel approach, the epigenome-based outlier approach, to identify tumor-suppressor genes silenced by aberrant DNA methylation. *Cancer letters*. 2012; 322(2):204–12. doi: [10.1016/j.canlet.2012.03.016](https://doi.org/10.1016/j.canlet.2012.03.016) PMID: [22433712](https://pubmed.ncbi.nlm.nih.gov/22433712/).
 30. Charite J, de Graaff W, Consten D, Reijnen MJ, Korving J, Deschamps J. Transducing positional information to the Hox genes: critical interaction of cdx gene products with position-sensitive regulatory elements. *Development*. 1998; 125(22):4349–58. PMID: [9778495](https://pubmed.ncbi.nlm.nih.gov/9778495/).
 31. Oshima M, Endoh M, Endo TA, Toyoda T, Nakajima-Takagi Y, Sugiyama F, et al. Genome-wide analysis of target genes regulated by HoxB4 in hematopoietic stem and progenitor cells developing from embryonic stem cells. *Blood*. 2011; 117(15):e142–50. doi: [10.1182/blood-2010-12-323212](https://doi.org/10.1182/blood-2010-12-323212) PMID: [21343615](https://pubmed.ncbi.nlm.nih.gov/21343615/); PubMed Central PMCID: PMC3296623.
 32. Cerda-Esteban N, Spagnoli FM. Glimpse into Hox and tale regulation of cell differentiation and reprogramming. *Developmental dynamics: an official publication of the American Association of Anatomists*. 2014; 243(1):76–87. doi: [10.1002/dvdy.24075](https://doi.org/10.1002/dvdy.24075) PMID: [24123411](https://pubmed.ncbi.nlm.nih.gov/24123411/).