

Testing the yield of a pilot-scale bubble column photobioreactor for cultivation of the microalga *Rhodomonas salina* as feed for intensive calanoid copepod cultures

Thuy, Minh Vu Thi; Jepsen, Per Meyer; Jørgensen, Niels O. G.; Hansen, Benni Winding; Nielsen, Søren Laurentius

Published in:
Aquaculture Research

DOI:
[10.1111/are.13868](https://doi.org/10.1111/are.13868)

Publication date:
2018

Document Version
Peer reviewed version

Citation for published version (APA):

Thuy, M. V. T., Jepsen, P. M., Jørgensen, N. O. G., Hansen, B. W., & Nielsen, S. L. (2018). Testing the yield of a pilot-scale bubble column photobioreactor for cultivation of the microalga *Rhodomonas salina* as feed for intensive calanoid copepod cultures. *Aquaculture Research*, 50(1), 63-71. <https://doi.org/10.1111/are.13868>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

1 **Testing the yield of a pilot-scale bubble column photobioreactor for**
2 **cultivation of the microalga *Rhodomonas salina* as feed for intensive**
3 **calanoid copepod cultures**

4 Minh Thi Thuy Vu¹, Per Meyer Jepsen¹, Niels O. G. Jørgensen², Benni Winding Hansen¹,
5 Søren Laurentius Nielsen^{1*} ¹Department of Science and Environment, Roskilde University,
6 Universitetsvej 1, Postbox 260, Building 11.2, DK-4000 Roskilde, Denmark.

7 ²Department of Plant and Environmental Sciences, University of Copenhagen, Copenhagen,
8 Denmark

9 *Corresponding author: Søren Laurentius Nielsen, Tel.: +45 4674 2722, email address:
10 nielsen@ruc.dk

11

12 Running title: A PBR for cultivation of *Rhodomonas salina*

13 Keywords: Amino acids; Dilution rate; Fatty acids; Growth rate; Live feed; Upscaling

14

15 **Abstract**

16 A dual column photobioreactor (PBR) (2×47 L) with mixed CO₂/air bubbling was tested for
17 cultivation of the microalga *Rhodomonas salina* as food for live feed copepods. In the continuous
18 growth phase, the cell density was relatively stable at $2.40 \pm 0.13 \times 10^6$ cells mL⁻¹ at an average
19 dilution rate of 0.46 ± 0.02 day⁻¹ throughout the 30-day experiment. The produced algae had a high
20 content of both total fatty acids (TFA) and free amino acids (FAA). Especially, the harvested algae
21 contained a high proportion of poly-unsaturated fatty acids (PUFA) that made up 80 % of the TFA
22 and of essential amino acids (35% of all FAA), implicating desirable components as feed for
23 copepods. The current PBR was sufficient to feed a culture of the calanoid copepod *Acartia tonsa* at
24 a density of 2,500 adult L⁻¹ in ca. 500 L culture with a daily yield of approximately 17×10^6 eggs.
25 To be able to sustain the integrated copepods production, the suggested volume of the algae cultures
26 should be ca. 20% of the copepod culture volume.

27

28 **Introduction**

29 Microalgae are important live feeds for early life stages of many marine aquaculture species ranging
30 from zooplankton to fish (e.g. Støttrup, 2003 ; Muller-Feuga et al., 2003). Microalgae are attractive
31 feed organisms because of their size, rapid growth rate, potential for mass-cultivation, digestibility,
32 and particularly their high nutritional value (Brown, 2002). The nutritional quality of microalgae is
33 indicated by the abundance of highly unsaturated fatty acids (HUFA), particularly eicosapentaenoic
34 acid (EPA), 20:5n-3 and decosahexaenoic acid (DHA), 22:6n-3 (Renaud et al., 1991) and the
35 composition of amino acids (AAs) (Brown, 1991; Guisande et al., 2000). For example, HUFA are
36 essential fatty acids for various aquaculture animals (Nichols, 2003), also sustaining growth and
37 reproduction of copepods (Rasdi & Qin, 2016).

38 In aquaculture, microalgae are grown either in simple cultivation systems, such as open ponds,
39 raceways or aerated open carboys using natural sun light or in more complex closed systems such as
40 closed cylindrical tanks, vertical aerated column photobioreactors (PBRs) or tubular flat-plate PBRs
41 (reviewed in Zmora & Richmond, 2004; Ugwu et al., 2008; Carvalho et al., 2006). The closed
42 PBRs offer controllable culturing conditions such as temperature, light, pH, carbon dioxide (CO₂),
43 but they require a high initial investment cost (Singh & Sharma, 2012; Ugwu et al., 2008).

44 The present study aimed to test the culturing capacity of a pilot-scale double bubble-column PBR (2
45 × 47 L) for cultivation of *R. salina* as food for intensive copepod production. Eriksen et al. (1998)
46 succeeded in using a small-scale column PBR (1.7 L) to cultivate *Rhodomonas* sp. with a cell
47 density reaching 10⁷ cells mL⁻¹ for up to 415 days continuously with a dilution rate of 0.6 day⁻¹. For
48 large-scale cultivation of microalgae in the aquaculture industry, the production in experimental
49 small-scale PBRs must be up-scaled and the efficiency maintained. Major challenges when
50 upscaling microalgal production include increasing the total volume of the PBR, while at the same

time maintaining an optical path in the PBR that is short enough to still achieve maximal growth and output rates of the algae (Hu et al., 1996). Another challenge is to maintain a sufficient mixing in a large PBR without causing a shear stress high enough to break the algal cell membranes (Qiang & Richmond, 1996). In the present contribution, we tested capacity of a medium-sized PBR for production of microalgae with respect to both quantity and quality (fatty acids and amino acids) as feed for copepods.

57

58 **Materials and methods**

59 *Algal strain*

The cryptophyte *R. salina* was obtained as strain K-1487 from the Scandinavian Culture Collection of Algae and Protozoa (SCCAP). The stock culture of *R. salina* was grown in the laboratory at Roskilde University (RUC) at standard conditions as described by Vu et al. (2016). The cultures were maintained under a continuous irradiance of 80 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ photosynthetically active radiation (PAR) in a climate room at 20 °C. The flasks were gently aerated with atmospheric air through 0.45- μm filters.

66

67 *Design and description of bioreactors*

R. salina was cultured in the dual column PBR shown in Fig. 1. Each culture column consists of a 150 cm high glass cylinder that for practical reasons is made of two 75 cm high glass cylinders on top of each other held together by a plastic ring. The inner diameter is 20 cm, giving a volume of 47 L for each of the two columns with a volume of 94 L for the entire PBR. The glass cylinders are placed on a polypropylene base and closed with silicon O-rings at the top and bottom.

73 Seawater was filtered through a 0.2 μm filter and purified with ultra-violet light (UV) (Ultra Violet
74 filter, Water.dk WP/75/60 at a fluence of 65 - 97 $\text{mW}\cdot\text{s cm}^{-2}$) and nutrients were supplied through a
75 peristaltic pump from the bottom of the culture column.

76 The microalgae were kept in suspension by large bubbles from the air system, and atmospheric
77 airflow was adjusted by a flow meter. Using another flow meter, a lower atmospheric flow was
78 mixed with pure CO_2 from a pressure flask. The mixture of CO_2 and atmospheric airflow was
79 regulated automatically using a feedback system, where opening or closing of solenoid valves were
80 controlled by a pH probe in the column. The mixture of pure CO_2 and atmospheric air was passed
81 through a membrane to increase formation of microbubbles to improve the mass transfer rate of
82 CO_2 for algae growth.

83 A cooling coil was installed inside each of the PBR culture columns and allowed adjustment of the
84 temperature in the PBR to 20°C . A temperature sensor was used as an *in-situ* measuring probe to
85 regulate the feedback temperature system, by opening and closing solenoid valves.

86 Light was supplied by 4 pairs per PBR of LCD lamps (Aussmak ELT-S0335 T9/35W/2700K) from
87 4 sides and was adjusted to provide light intensities at 3 different levels: low, medium and high,
88 corresponding to 393 ± 44 , 608 ± 80 and $981 \pm 133 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ Photosynthetic Active Radiation
89 (PAR), respectively.

90 The microalgae were harvested from an overflow at top of each culture column of the PBR. The
91 harvested algae were stored in a tank and used for copepod cultures in $4 \times 350 \text{ L}$ tanks at Roskilde
92 University's copepod production facility (Fig. 1).

93 The PBR was connected and controlled by Programmable Logic Control (PLC). From the PLC, the
94 light mode, temperature, pH, the pump flows (for input seawater, nutrients, copepod feeding) were
95 set to required levels. Data such as pH and temperature were logged and stored every 30 minute.

96 *Environmental parameters in the photobioreactor*

97 Temperature and pH in the PBR in the three experiments were maintained at 20.01 ± 0.03 °C and
98 8.11 ± 0.03 , respectively. Nitrate concentrations were maintained at 2211 ± 374 $\mu\text{mol L}^{-1}$
99 throughout the experiment, and the phosphate concentrations at 331 ± 33 $\mu\text{mol L}^{-1}$ during the
100 experiment. Nitrate concentrations were determined by flow injection analysis using QuickChem
101 Method 31-107-04-1-A (detection limit is 0.02 $\mu\text{mol}\cdot\text{L}^{-1}$, (Diamond, 1999)), while phosphate
102 concentrations were quantified spectrophotometrically by the method described by S ndergaard and
103 Riemann (1979).

104 *Experimental design*

105 The *R. salina* was cultured under conditions with nutrients in excess (B1-medium, added daily as 1
106 mL B1 L^{-1} seawater, (Hansen, 1989), and a high light intensity ($60\text{--}140$ $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$) to
107 generate a relatively large algal biomass with desired fatty acids (FA) and free essential amino acids
108 (FAA) profiles (*sensu* Vu et al., (2016)).

109 The growth of *R. salina* was investigated in the PBR as four replicates ($2 \text{ times} \times 2$ culture column).
110 At start, each of the culture column ($V = 47$ L) was filled with UV-treated seawater and inoculated
111 with fresh algae stock cultures to a total volume of ca. 42 L with the initial cell density of $125,229 \pm$
112 $3,720$ cells mL^{-1} . The PBR was run in the initial growth phase for five days. The B1 medium was
113 added manually at the beginning of the experiment and every 1-2 days (rate at 1.2 mL L^{-1} , (Hansen,
114 1989) to ensure that nutrients were in excess in the culture. After the first five days, the algal culture

115 in the PBR was brought into the continuous phase, where steady state conditions were obtained by
116 adding UV-treated seawater by two pumps (ProMinent DULCOflex) at bottom of each of the two
117 PBRs (Fig. 1) to dilute the algal density. The B1 medium was added at a rate of 1.2-4 mL L⁻¹ of
118 newly added seawater.

119 The light was set at the low intensity for the first 2 days when the algae density was low to avoid
120 photo-inhibition. When the algae density had increased after day 3, the light was changed to the
121 highest intensity to minimize a potential light limitation.

122 The main atmospheric airflow was set at ca. 4 L min⁻¹ at the first 2-3 days of the experiment when
123 the algal cell density was low and increased to 5-6 L min⁻¹ when the algal cell density was high
124 from day 2-3 onward. The mixture of pure CO₂ (V/V = 2-5% of CO₂ in mixture) and atmospheric
125 air (V = 2-2.5 L) was added to supply CO₂ and to control the pH at 8.1 ± 0.1.

126 *Analytical procedures and measurements*

127 Samples of *R. salina* (3-4 analytical replicates, each with 50-100 mL of culture medium per sample)
128 were taken daily from each culture column to determine growth and production of the algae and
129 biochemical composition of algal cells. The cell density of *R. salina* was determined daily by
130 measuring 3 to 4 replicates in a Beckman MultisizerTM3 Coulter Counter[®] (Beckman Coulter Inc.,
131 USA). The specific growth rate (day⁻¹) of *R. salina* was calculated by fitting cell density increase in
132 each of the culture columns during the initial phase to an exponential growth function (Vu et al.,
133 2016).

134 The cell density and total volume of the harvested algae in each harvest tank (collected from the
135 culture columns) were also measured daily. The cell density of *R. salina* was converted into carbon
136 weight by assuming that one algal cell equals 47.4 pg C (Berggreen et al., 1988).

137

138 For analysis of fatty acids (FA) and free amino acids (FAA) in the microalgae, samples of about 10^6
139 algal cells were collected during both the initial (14 samples; $n=14$) and the continuous growth
140 phase (18 samples; $n=18$). The samples were filtered through a 25-mm diameter GF/C glass fiber
141 filter (Whatman) and stored in Pyrex glass vials for FA analysis, or in 1.5 mL HPLC glass vials for
142 FAA analysis, at $-80\text{ }^{\circ}\text{C}$ for later analysis FA and FAA compositions.

143 The FA composition was determined by extraction of the lipids using a HPLC-grade chloroform:
144 methanol mixture (Folch et al., 1957) followed by a trans-esterification process by acetyl chloride in
145 methanol. The detailed procedure of the FA analyses was described by Drillet et al. (2006) with
146 minor adjustments: addition of 20 μL internal standard (C-23-methylester, $1000\text{ }\mu\text{g ml}^{-1}$) and no
147 sonication.

148 For analysis of FAA, each vial containing a filter sample was added 1 mL Milli-Q water, heated to
149 95°C for 15 min and filtered through 0.2 μm cellulose nitrate syringe filters (VWR International,
150 USA) after cooling. FAA concentrations were determined by HPLC and fluorescence detection
151 after derivatization with 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate (AQC). AQC was
152 purchased from Waters as an AccQFlour kit (www.waters.com). The analysis was conducted as
153 recommended by Waters, except for adjustment of the solvent gradient, and the AA derivatives
154 were analyzed on a Waters HPLC system (Alliance 2695 solvent module, 2475 fluorescence
155 detector and a $3.9 \times 150\text{ mm}$ AccQTag column). Chromatogram peaks were identified from
156 retention times and integrated against an external amino acid standard mixture (AA-S-18 amino
157 acid mixture; www.sigma-aldrich.com).

158

159 *Calculated capacity of the photobioreactor to produce algal biomass for copepod cultures*

160 The daily production of *R. salina* served as food for the calanoid copepod *Acartia tonsa* culture at
161 Roskilde University, Denmark. The estimated production rate of the PBR for both culture columns
162 (P , g C day⁻¹) was calculated as equation (Eq. 1):

163
$$P = D \times V_A \times D_A \times W_A \quad (\text{Eq. 1})$$

164 In which

165 D : dilution rate (day⁻¹)

166 V_A : filled volume (L) of the two culture columns of the photobioreactor

167 D_A : density of the algae harvested, assuming that the density of algae harvested equals the density
168 inside the culture columns (cells·L⁻¹)

169 W_A : carbon weight of algae cells = 47.4×10^{-12} g C cell⁻¹ (Berggreen et al. 1988)

170 The algae requirement (in carbon weight) for 1 L of copepods per day (C_A , g C L⁻¹ day⁻¹) was
171 calculated as equation (Eq. 2)

172
$$C_A = D_C \times [W_F \times F + W_M(1 - F)] \times IR_{Max} \quad (\text{Eq. 2})$$

173 In which

174 C_A : algae requirement (carbon units) for 1 L of copepods.

175 D_C : density of copepods (ind L⁻¹), we used a density of 2,500 adult copepods·L⁻¹ which is reported
176 to be the optimal density of *A. tonsa* for maximal egg yield in culture tanks (Drillet et al. 2015)

177 W_F : The average carbon weight of female copepods = 3,193 ng C ind⁻¹ = 3,193×10⁻⁹ g C ind⁻¹
178 (Drillet et al. 2015)

179 W_M : The average carbon weight of male copepods = 2,282 ng C ind⁻¹ = 2,282×10⁻⁹ g C ind⁻¹ (Drillet
180 et al. 2015)

181 F : the female: male ratio = 0.45 (Drillet et al. 2015)

182 IR_{Max} : maximum specific carbon ingestion rate of *R. salina* K-1487 by the same strain of *A. tonsa* =
183 1.32 day⁻¹ (Berggreen et al. 1988)

184 The volume of copepod culture that can be sustained by feeding on the current algae production
185 (V_C , L) was estimated as equation (Eq. 3):

186
$$V_C = \frac{P}{C_A} \quad (\text{Eq. 3})$$

187 The egg production of *A. tonsa* was estimated assuming that 1/3 of carbon ingested by female
188 copepods is allocated to egg production (Kiørboe et al. 1985). Theoretical specific egg production
189 per individual (SEP , eggs female⁻¹ day⁻¹) was calculated as equation (Eq. 4)

190
$$SEP = \frac{W_F \times IR_{Max}}{3 \times W_E} \quad (\text{Eq. 4})$$

191 Whereas W_E : the carbon weight of the egg: 45.7×10⁻⁹ g C egg⁻¹ (Kiørboe et al. 1985)

192 Potential number of eggs produced with the current algal production (EP , eggs day⁻¹) was estimated
193 as equation (Eq. 5):

194
$$EP = SEP \times D_C \times F \times V_C \quad (\text{Eq. 5})$$

195 *Statistical analysis*

196 Data on fatty acid and amino acid content were subjected to one-way ANOVA. Data were tested for
197 homogeneity of variance (Cochran's test) and normal distribution (Kolmogorov-Smirnoff goodness
198 of fit test) before being analyzed by ANOVA. All tests on data were carried out using SYSTAT v.
199 13 with $\alpha = 0.05$. ANOVA was performed on data from the initial phase and the continuous phase
200 separately, and as the ANOVA showed no significant differences in data within each phase ($p >$
201 0.05), data on content of the various fatty acids and amino acids were pooled within each phase.
202 Subsequently, differences between content in the two phases were tested for each compound by
203 one-way ANOVA, using SYSTAT v. 13 with $\alpha = 0.05$.

204 **Results**

205 *Growth of R. salina in the photobioreactor*

206 In the initial phase (day 1 to 5), the specific growth rate of *R. salina* was $0.81 \pm 0.07 \text{ day}^{-1}$ (mean \pm
207 SE). The average cell density increased from Day 0 and peaked at $4.17 \pm 1.43 \times 10^6 \text{ cells mL}^{-1}$
208 (mean \pm SE) at Day 5 (Fig. 2). From Day 6, the cell density remained relative stable at 2.40 ± 0.13
209 $\times 10^6 \text{ cells mL}^{-1}$ (mean \pm SE). The dilution rate varied over time with an average of 0.46 day^{-1}
210 throughout the experiment (Fig. 2). The average production rate for one 47 L PBR column was
211 determined to $44.60 \pm 5.25 \times 10^9 \text{ cells day}^{-1}$ (mean \pm SE), equivalent to $2.11 \pm 0.25 \text{ g C day}^{-1}$.

212 *Fatty acids*

213 Data from each growth phase showed a similar content of fatty acids within the initial and
214 continuous phase respectively (one-way ANOVA, $p > 0.05$). The fatty acid data were therefore
215 pooled within each phase for further analysis. The total fatty acid (TFA) content of *R. salina*

216 increased from 28.38 ± 2.81 pg cell⁻¹ (mean \pm SE, n = 18) in the initial phase, to 41.37 ± 2.46 pg
 217 cell⁻¹ (mean \pm SE, n = 18) during the continuous phase (Fig. 3A, Table 1). This increase was
 218 statistically significant (one-way ANOVA, $p < 0.001$). The DHA content was significantly lower
 219 (2.74 ± 0.19 pg cell⁻¹, mean \pm SE) in the initial phase (one-way ANOVA, $p < 0.01$) than in the
 220 continuous phase (3.27 ± 0.16 pg cell⁻¹, mean \pm SE, Fig. 3A), but the relative abundance of DHA as
 221 percentage of TFA was higher in the initial phase (10.41 ± 0.70 % of TFA, mean \pm SE) than that in
 222 the continuous phase (8.23 ± 0.39 % (mean \pm SE) of TFA, Table 1). Similarly, the EPA content
 223 increased significantly (one-way ANOVA, $p < 0.01$) from 3.45 ± 0.35 pg cell⁻¹ (mean \pm SE) in the
 224 initial phase to 4.71 ± 0.26 pg cell⁻¹ (mean \pm SE) in the continuous phase (Fig. 3A), but the relative
 225 abundance remained at about 12 % of TFA, independent of the growth phases (Table 1). However,
 226 the DHA/EPA ratio decreased from 0.84 ± 0.04 (mean \pm SE) in the initial phase to 0.71 ± 0.02 %
 227 (mean \pm SE) of TFA in the continuous phase (Table 1).

228 The fatty acids of *R. salina* were dominated by short chain polyunsaturated fatty acids (SC-PUFA).
 229 These fatty acids evinced a statistically significant lower content (one-way ANOVA, $p < 0.01$) in
 230 the initial phase (51.99 ± 1.81 % of TFA, mean \pm SE) than in the continuous phase (57.62 ± 1.72 %
 231 of TFA, mean \pm SE, Fig. 3B, Table 1). In contrast, the relative abundance of the monounsaturated
 232 fatty acids (MUFA) was significantly lower in the continuous phase (one-way ANOVA, $p < 0.01$)
 233 (6.49 ± 0.54 % of TFA), as compared to the initial phase (11.37 ± 1.97 % of TFA, Fig. 3B, Table
 234 1). The relative abundance of highly saturated fatty acids (HUFA) and the saturated fatty acids
 235 (SFA) were not significantly different in the two growth phases (one-way ANOVA, $p > 0.05$, Fig.
 236 3B, Table 1).

237

238 *Free amino acids*

239 As was the case for the fatty acids, data from each growth phase showed a similar content of FAA
240 within the initial and continuous phase respectively (one-way ANOVA, $p > 0.05$). The FAA data
241 were thus also pooled within each phase for further analysis. The total free amino acid (TFAA)
242 content was relative stable at 3 pg cell^{-1} , regardless the growth phase (one-way ANOVA, $p > 0.05$,
243 Fig. 4A, Table 2). The essential amino acids made up around 1 pg cell^{-1} , corresponding to ca. 35%
244 of TFAA, irrespective of the growth phase (one-way ANOVA, $p > 0.05$, Fig. 4A, Table 2). No
245 significant differences in the relative abundance of the various EAA were observed between
246 different growth phases either (one-way ANOVA, $p > 0.05$, Fig. 4B, Table 2). The most abundant
247 EAA was arginine (ca. 18% of TFAA), followed by lysine (ca. 4% of TFAA, Fig. 4B).

248 *Capacity of the photobioreactor to produce algal biomass for a copepod culture*

249 Our estimation shows that the current production of the two column PBR can be 92.74×10^9 cells
250 day^{-1} which can sustain ca. 500 L of copepod *A. tonsa* culture at the density of 2500 ind. L^{-1} (Table
251 3). The estimated number of eggs produced from such *A. tonsa* cultures were 17.11×10^6 eggs day^{-1}
252 (Table 3).

253 **Discussion**

254 The purpose of this study was testing of a dual bubble-column PBR ($2 \times 47 \text{ L}$) for cultivation of the
255 microalga *R. salina* to examine if the methodology can be scaled up for commercial production in
256 the aquaculture industry. For large-scale production, a stable, continuous, and high-quality
257 production of algae is needed if the algal should serve as feed for cultures of sustain calanoid
258 copepods that later can be used as live feed in marine fish hatcheries. Thus, the overall aim was to
259 integrate PBR and copepod culture facilities at a future industrial scale.

260 *Cell density and daily production of PBR*

261 The specific growth rate obtained in this study ($0.81 \pm 0.07 \text{ day}^{-1}$) was similar to the previously
262 reported growth rates of *Rhodomonas* species ($0.3\text{-}0.8 \text{ day}^{-1}$, (Guevara et al., 2016; Vu et al., 2016;
263 Lafarga-De la Cruz et al., 2006; Eriksen et al., 1998). In contrast to these previous studies, we
264 successfully up-scaled the *Rhodomonas* cultures from low-volume cultures of 0.1-1.7 L to volumes
265 closer to a future commercial scale. In a substantially smaller PBR system (1.5 L cultures),
266 *Rhodomonas* sp. was produced and harvested continuously for 415 d at a cell density of 10^7 cells
267 mL^{-1} and with a dilution rate of 0.6 day^{-1} (Eriksen et al., 1998). In our 30 times larger PBRs, the cell
268 density was about one quarter of this during the continuous phase, but the dilution rates of algae
269 obtained were almost the same as in the study of Eriksen et al. (1998). The lower cell density
270 obtained in our PBR reflects its much larger inner diameter (20 cm vs. 7.5 cm (Eriksen et al., 1998;
271 Zhang & Richmond, 2003)), but the larger volume in our PBR more than compensates for the lower
272 volumetric productivity. This means that although the upscaling lowers the production efficiency
273 (PE) with a factor $\frac{1}{4}$, the total volumetric productivity is about 7.5 higher than in small-scale
274 facilities.

275 *Nutritional quality of algae*

276 Compared to the initial exponential phase, a higher TFA content was found in *R. salina* during the
277 continuous phase. Since microalgae for copepod feed typically are harvested in the continuous
278 phase, this observation is important. This increase in TFA is often associated with nitrogen
279 limitation (Gladue & Maxey, 1994), but this is not the case in the present study, as the nitrogen
280 concentration in the PBR is kept at a high, non-limiting, level. Our results indicate, however, that *R.*
281 *salina* accumulate more FA in steady state growth than under the initial phase. A similar pattern has
282 been observed for *Chlorella vulgaris* when growing under a high supply of CO_2 (Jose &

283 Suraishkumar, 2016) and in *Nannochloropsis oceanica* under nitrogen replete conditions as in the
284 present study (Xiao et al., 2015). Another observation from the continuous phase in our cultures
285 was an increased presence of desirable fatty acids, SC-PUFA (mainly C18:3 n-3) when the algae
286 grew in the continuous phase. In contrast, algae harvested in the continuous phase contained less
287 MUFA as compared to the algae in the initial phase. An enhanced content of SC-PUFA in
288 microalgae has previously been observed under culture conditions utilizing nitrate as the nitrogen
289 source, eliminating the pH fluctuations caused by ammonium use, and non-limiting concentrations
290 of phosphate, as many of the SC-PUFAs are found as phospholipids (Yongmanitchai & Ward,
291 1991).

292 In aquaculture, a DHA/EPA ratio of 2 is proposed as an optimum criterion for live food organisms,
293 leading to the preferred ratio in the next link in the aquaculture food chain, the fish larvae ((Reitan
294 et al., 1994)). The DHA/EPA ratio obtained in the present study (DHA/EPA ratio of 0.7 to 0.8) was
295 similar to previous studies on the same algal species (DHA/EPA ratio of 0.5 to 0.9; (Dunstan et al.,
296 2005; Mansour et al., 2005; Vu et al., 2016; Guevara et al., 2016)), indicating that the preferred
297 ratio of 2 was not reached. However, although both historical data and our data show that the
298 DHA/EPA ratio in the cultured algae is below the optimum ratio, the DHA/EPA ratio will be closer
299 to 2 by trophic upgrade when *Rhodomonas* is fed copepods that again are fed to the fish larvae.
300 Thus, Drillet et al. (2006) observed a DHA/EPA ratio of about 1.3 in fish larvae when fed *Acartia*
301 *tonsa* based on *R. salina* from our laboratory strains.

302 Regarding the general composition of FA in *R. salina* cultured in the PBR, the composition agreed
303 with previous results of especially PUFA (made up ca. 80% of TFA in this study), including SC-
304 PUFA and HUFA ((Vu et al., 2016)). PUFA are also well known to be crucial for the production

305 and hatching success of copepod eggs (Arendt et al., 2005; Broglio et al., 2003; Jónasdóttir et al.,
306 2009).

307 The increase in cell-specific content of FA in the continuous phase was not observed in TFAA that
308 remained unchanged during the entire study period. This uncoupling between FA and FAA was also
309 observed in another marine alga in which precursors for FA increased, while most of the FAA
310 declined during N starvation (Zhang et al., 2016). These authors suggested that the changes in
311 intracellular FAA pool reflected synthesis and catabolism of AA, as well as proteolytic activity, e.g.
312 degradation of Rubisco. Thus, in *R. salina*, synthesis and catabolism of FAA may have been
313 balanced with proteolytic activity throughout the study.

314 The FAA content in zooplankton is reported to be closely dependent on content and composition in
315 algae used as feed for the zooplankton (e.g., Laabir et al., 1999) and in this respect, *Rhodomonas* is
316 considered a very promising food alga for various zooplankton species, including our target species
317 (Drillet et al. 2006). For the next trophic level, the fish larvae, Rayner et al. (2017) has recently
318 published an overview of fish larval needs. Essential FAs, especially leucine and valine, as well as
319 threonine, arginine and methionine are considered most important. These amino acids were all
320 present in significant amounts in the present *R. salina* strain.

321 *Availability as food for copepod productions*

322 Based on our estimation, the current daily production of *R. salina* in our two PBRs, about 4.40 g C
323 day⁻¹, can sustain an intensive copepod production tank at a volume of ca. 500 L at an optimum
324 density of ca. 2,500 adult copepods L⁻¹, as proposed in Drillet et al. (2015). This copepod
325 production can deliver a daily egg production of ca. 17.11×10^6 eggs per culture tank. Therefore, to
326 be able to feed the copepods optimally, the required volume of algal culture at the cell density

327 achieved in this study is ca. 20% of the copepod culture. It should be noted that the estimation of
328 algae feeding to the copepods was based on the grazing rate of adult copepods. For younger life
329 stages, e.g., nauplii and copepodite stages, the demand for algae is substantially smaller.

330 *Recommendations*

331 The microalga *R. salina* was successfully cultivated in the current bubble-column PBR and
332 produced algal biomass with a high biochemical quality and that can immediately be fed out to
333 copepod cultures. Based on this, it is recommended that *R. salina* should be cultivated with the
334 addition of pure CO₂ in a mixture with atmospheric air (V/V=2-5% of CO₂ in the mixture) to
335 maintain the pH at 8.1. The harvested algae in the continuous growth phase obtained a high TFA
336 content with a relative high abundance of PUFA (including SC-PUFA and HUFA) as well as a high
337 content of FAA, which is of crucial importance for enhancing the egg production, the egg hatching
338 success, and the somatic growth of the cultured copepods. For copepod production in aquaculture
339 facilities, the estimated volume of algae cultures should be about 20% v/v of copepod culture.

340 **Acknowledgments**

341 This work was funded by the Danish National Strategic Research Council IMPAQ-IMProvement of
342 AQUaculture high quality fish fry production grant (Grant. no. 10-093522) to Benni Winding
343 Hansen and the Danish National Advanced Technology Foundation COMA-COpepod egg Mass
344 production in Aquaculture grant (Grant no. 67-2013-1) to Benni Winding Hansen and Søren
345 Laurentius Nielsen. We would like to thank Anne B. Faarborg and Rikke Guttesen (Roskilde
346 University, Denmark) for laboratory assistance.

References

- Arendt KE, Jónasdóttir SH, Hansen PJ, Gärtner S (2005) Effects of dietary fatty acids on the reproductive success of the calanoid copepod *Temora longicornis*. *Mar. Biol.*, **146**, 513-530.
- Berggreen U, Hansen B, Kiørboe T (1988) Food size spectra, ingestion and growth of the copepod *Acartia tonsa* during development: implications for determination of copepod production *Mar. Biol.*, **99**, 341-352.
- Broglio E, Jónasdóttir SH, Calbet A, Jakobsen HH, Saiz E (2003) Effect of heterotrophic versus autotrophic food on feeding and reproduction of the calanoid copepod *Acartia tonsa*: Relationship with prey fatty acid composition. *Aquat. Microb. Ecol.*, **31**, 267-278.
- Brown MR (1991) The amino-acid and sugar composition of 16 species of microalgae used in mariculture. *J. Exp. Mar. Biol. Ecol.*, **145**, 79-99.
- Brown MR (2002) Nutritional value of microalgae for aquaculture. In: *Avances en Nutrición Acuicola VI. Memorias del VI Simposium Internacional de Nutrición Acuicola*. (eds Cruz-Suárez LE, Ricque-Marie D, Tapia-Salazar M, Gaxiola-Cortés, M. G. S, N), Cancún, Quintana Roo, México.
- Carvalho AP, Meireles LA, Malcata FX (2006) Microalgal reactors: A review of enclosed system designs and performances. *Biotechnol. Prog.*, **22**, 1490-1506.
- Diamond D (1999) Determination of nitrate in brackish or seawater by flow injection analysis. QuikChem Method 31-107-04-1-A. (ed Zellweger Analytics Lachat Instruments Division WMR), Millwaukee, WI 53218-1239 USA, pp. 16.
- Drillet G, Jørgensen NOG, Sørensen TF, Ramløv H, Hansen BW (2006) Biochemical and technical observations supporting the use of copepods as live feed organisms in marine larviculture. *Aquacult. Res.*, **37**, 756-772.
- Drillet G, Rais M, Novac A, Jepsen PM, Mahjoub M-S, Hansen BW (2015) Total egg harvest by the calanoid copepod *Acartia tonsa* (Dana) in intensive culture – effects of high stocking densities on daily egg harvest and egg quality. *Aquacult. Res.*, **46**, 3028-3039.
- Dunstan GA, Brown MR, Volkman JK (2005) Cryptophyceae and rhodophyceae; chemotaxonomy, phylogeny, and application. *Phytochemistry*, **66**, 2557-2570.
- Eriksen N, Poulsen B, Lønsmann Iversen J (1998) Dual sparging laboratory-scale photobioreactor for continuous production of microalgae. *J. Appl. Phycol.*, **10**, 377-382.
- Folch J, Lees M, Stanley GHS (1957) A simple method for isolation and purification of total lipids from animal tissues. *J. Biol. Chem.*, **226**, 497-509.

380 Gladue RM, Maxey JE (1994) Microalgal feeds for aquaculture. *J. Appl. Phycol.*, **6**, 131-141.

381 Guevara M, Arredondo-Vega BO, Palacios Y, Saéz K, Gómez PI (2016) Comparison of growth and
382 biochemical parameters of two strains of *Rhodomonas salina* (Cryptophyceae) cultivated
383 under different combinations of irradiance, temperature, and nutrients. *J. Appl. Phycol.*, 1-
384 10.

385 Guisande C, Riveiro I, Maneiro I (2000) Comparisons among the amino acid composition of
386 females, eggs and food to determine the relative importance of food quantity and food
387 quality to copepod reproduction. *Mar. Ecol. Prog. Ser.*, **202**, 135-142.

388 Hansen PJ (1989) The red tide dinoflagellate *Alexandrium tamarense*: Effects on behaviour and
389 growth of a tintinnid ciliate. *Mar. Ecol. Prog. Ser.*, **53**, 105-116.

390 Hu Q, Guterman H, Richmond A (1996) A flat inclined modular photobioreactor for outdoor mass
391 cultivation of photoautotrophs. *Biotechnol. Bioeng.*, **51**, 51-60.

392 Hulatt CJ, Wijffels RH, Bolla S, Kiron V (2017) Production of Fatty Acids and Protein by
393 *Nannochloropsis* in Flat-Plate Photobioreactors. *PLOS ONE*, **12**, e0170440.

394 Jónasdóttir SH, Visser AW, Jespersen C (2009) Assessing the role of food quality in the production
395 and hatching of *Temora longicornis* eggs. *Mar. Ecol. Prog. Ser.*, **382**, 139-150.

396 Jose S, Suraishkumar GK (2016) High carbon (CO₂) supply leads to elevated intracellular acetyl
397 CoA levels and increased lipid accumulation in *Chlorella vulgaris*. *Algal Res.*, **19**, 307-315.

398 Lafarga-De la Cruz F, Valenzuela-Espinoza E, Millán-Núñez R, Trees CC, Santamaría-del-Ángel
399 E, Núñez-Cebrero F (2006) Nutrient uptake, chlorophyll a and carbon fixation by
400 *Rhodomonas sp.* (Cryptophyceae) cultured at different irradiance and nutrient
401 concentrations. *Aquacult. Eng.*, **35**, 51-60.

402 Laabir M, Poulet SA, Cueff A, Ianora A (1999) Effect of diet on levels of amino acids during
403 embryonic and naupliar development of the copepod *Calanus helgolandicus*. *Mar. Biol.*,
404 **134**, 89-98.

405 Mansour M, Frampton DF, Nichols P, Volkman J, Blackburn S (2005) Lipid and fatty acid yield of
406 nine stationary-phase microalgae: Applications and unusual C24–C28 polyunsaturated fatty
407 acids. *J. Appl. Phycol.*, **17**, 287-300.

408 Muller-Feuga A, Moal J, Kaas R (2003) The microalgae of aquaculture. In: *Live feeds in Marine*
409 *Aquaculture* (ed. by Støttrup JG, McEvoy LA). Blackwell Publising, Oxford, pp. 206-299.

410 Nichols DS (2003) Prokaryotes and the input of polyunsaturated fatty acids to the marine food web.
411 *FEMS Microbiol. Lett.*, **219**, 1-7.

412 Qiang H, Richmond A (1996) Productivity and photosynthetic efficiency of *Spirulina platensis* as
 413 affected by light intensity, algal density and rate of mixing in a flat plate photobioreactor. *J.*
 414 *Appl. Phycol.*, **8**, 139-145.

415 Rasdi NW, Qin JG (2016) Improvement of copepod nutritional quality as live food for aquaculture:
 416 a review. *Aquacult. Res.*, **47**, 1-20.

417 Rayner TA, Hwang J-S, Hansen BW (2017) Anticipating the free amino acid concentrations in
 418 newly hatched pelagic fish larvae based on recently fertilized eggs and temperature. *J.*
 419 *Plankton Res.*, **39**, 1012-1019.

420 Reitan KI, Rainuzzo JR, Olsen Y (1994) Influence of lipid composition of live feed on growth,
 421 survival and pigmentation of turbot larvae. *Aquacult. Int.*, **2**, 33-48.

422 Renaud SM, Parry DL, Thinh L-V, Kuo C, Padovan A, Sammy N (1991) Effect of light intensity on
 423 the proximate biochemical and fatty acid composition of *Isochrysis* sp. and *Nannochloropsis*
 424 *oculata* for use in tropical aquaculture. *J. Appl. Phycol.*, **3**, 43-53.

425 Singh RN, Sharma S (2012) Development of suitable photobioreactor for algae production – A
 426 review. *Renew. Sust. Energ. Rev.*, **16**, 2347-2353.

427 Støttrup JG (2003) Production and nutritional value of copepods. In: *Live feeds in Marine*
 428 *Aquaculture* (ed. by Støttrup JG, McEvoy LA). Blackwell Publishing, Oxford, pp. 145-205.

429 Søndergaard M, Riemann B (1979) *Ferskvandsbiologiske analysemetoder*, Akademisk Forlag,
 430 Copenhagen.

431 Ugwu CU, Aoyagi H, Uchiyama H (2008) Photobioreactors for mass cultivation of algae.
 432 *Bioresour. Technol.*, **99**, 4021-4028.

433 Vu MTT, Douët C, Rayner TA, Thoisen C, Nielsen SL, Hansen BW (2016) Optimization of
 434 photosynthesis, growth, and biochemical composition of the microalga *Rhodomonas*
 435 *salina*—an established diet for live feed copepods in aquaculture. *J. Appl. Phycol.*, **28**, 1485-
 436 1500.

437 Xiao Y, Zhang JT, Cui JT, Yao XZ, Sun ZJ, Feng YG, Cui Q (2015) Simultaneous accumulation of
 438 neutral lipids and biomass in *Nannochloropsis oceanica* IMET1 under high light intensity
 439 and nitrogen replete conditions. *Algal Res.*, **11**, 55-62.

440 Yongmanitchai W, Ward OP (1991) Growth of and Omega-3-fatty-acid production by
 441 *Phaeodactylum tricornutum* under different culture conditions. *Appl. Env. Microbiol.*, **57**,
 442 419-425.

443 Zhang CW, Richmond A (2003) Sustainable, high-yielding outdoor mass cultures of *Chaetoceros*
 444 *muelleri* var. *subsalsum* and *Isochrysis galbana* in vertical plate reactors. *Mar. Biotechnol.*,
 445 **5**, 302-310.
 446 Zhang YS, Liu YT, Cao XP, Gao P, Liu XY, Wang XY, Zhang JJ, Zhou JN, Xue S, Xu GW, Tian J
 447 (2016) Free amino acids and small molecular acids profiling of marine microalga *Isochrysis*
 448 *zhangjiangensis* under nitrogen deficiency. *Algal Res.*, **13**, 207-217.
 449 Zmora O, Richmond A (2004) Microalgae production for aquaculture. In: *Handbook of microalgal*
 450 *culture: biotechnology and applied phycology* (ed by Richmond A). Blackwell Science,
 451 Oxford, pp. 365-379.
 452
 453

454 **Captions for figures**

455 **Fig. 1** The pilot-scale bubble column photobioreactor for cultivation of *Rhodomonas salina*
456 connecting with the copepod cultures. Note: figure not drawn to scale

457 **Fig. 2** Cell density of *Rhodomonas salina* cultured in the PBR. The data indicated by means (\pm SE) of
458 the cell density (●) and dilution rate (Δ) of algae cultured in the PBR with 4 replicates from day 0 to
459 day 16 and only 1 replicate from day 17 onwards.

460 **Fig. 3** The total fatty acids, absolute DHA and EPA content (A) and the fatty acids main group
461 composition (B) of *Rhodomonas salina* during initial (black bar) and continuous phase (grey bar).
462 Data presented by means \pm SE. Different letters represent significant differences of a specific fatty
463 acid or main group of fatty acids at $p = 0.05$ between the two growth phases.

464 **Fig. 4** The total free amino acids (TFAA), absolute total essential amino acids (TEAA) content (A)
465 and the relative essential amino acids composition (B) of *Rhodomonas salina* during the initial
466 (black bar) and continuous phase (grey bar) growth phase. Data presented by means \pm SE.

467

468

469 **Captions for tables**

470 **Table 1** Total fatty acids (TFA) and FA compositions of *Rhodomonas salina* cultured in the
471 photobioreactors at different growth phases (initial vs continuous phase)

472 **Table 2** Results of one-way ANOVA testing the effects of different growth phases (initial vs
473 continuous phase) on total free amino acids, total essential amino acids content and specific amino
474 acids of *Rhodomonas salina* cultured in the photobioreactors

475 **Table 3** Estimated ability to support the copepod *Acartia tonsa* production (2500 ind L⁻¹) by
476 *Rhodomonas salina* algae produced in the current bubble column photobioreactors

477

478 **Table 1**

Variables	Initial phase	Continuous phase	df1	df2	SS	MS	F	p
C14:0	1.74±0.33 ^a	1.65±0.21 ^a	1	31	0.065	0.065	0.0585	0.81
C16:0	9.56±0.91 ^a	11.42±0.57 ^b	1	31	3.011	3.011	10.655	0.003
C18:0	1.18±0.19 ^a	0.56±0.07 ^b	1	31	3.011	3.011	10.655	0.003
SFA	12.68±1.02 ^a	13.78±0.65 ^a	1	31	9.456	9.456	0.891	0.353
C16:1	0.88±0.23 ^a	1.23±0.24 ^a	1	31	0.967	0.967	1.032	0.318
C18:1	3.45±0.57 ^a	3.59±0.37 ^a	1	31	0.159	0.159	0.0473	0.829
C22:1	6.71±2.48 ^a	1.26±0.30 ^b	1	31	234.318	234.318	6.137	0.019
MUFA	11.37±1.97 ^a	6.49±0.54 ^b	1	31	187.522	187.522	7.056	0.013
C18:2	10.48±1.55 ^a	15.25±1.39 ^b	1	31	179.275	179.275	5.244	0.029
C18:3 n-6	1.80±0.26 ^a	2.15±0.08 ^b	1	31	0.964	0.964	2.042	0.163
C18:3 n-3	18.26±0.93 ^a	22.46±0.84 ^b	1	31	139.059	139.059	11.203	0.002
C18:4	21.43±1.07 ^a	17.74±1.72 ^a	1	31	106.342	106.342	2.863	0.101
SC-PUFA	51.99±1.81 ^a	57.62±1.72 ^b	1	31	249.151	249.151	7.982	0.008
C20:4 n-6	1.06±0.18 ^a	2.00±0.14 ^b	1	31	6.905	6.905	17.639	<0.001
C20:5 n-3 (EPA)	12.47±0.57 ^a	11.86±0.73 ^a	1	31	2.936	2.936	0.394	0.535
C22:6 n-3 (DHA)	10.41±0.70 ^a	8.23±0.39 ^b	1	31	37.347	37.347	8.224	0.007
HUFA	23.95±1.06 ^a	22.11±1.08 ^a	1	31	26.684	26.684	1.428	0.241
DHA/EPA	0.84±0.04 ^a	0.71±0.02 ^b	1	31	0.138	0.138	10.136	0.003
TFA	28.38±2.81 ^a	41.37±2.46 ^b	1	31	1328.75	1328.75	12.118	0.002

479 Note: SFA: saturated fatty acids, MUFA: mono unsaturated fatty acids, SC-PUFA: short chain-poly
480 unsaturated fatty acids, HUFA: highly unsaturated fatty acids. Units of TFA: pg cell⁻¹, specific
481 FA/FA group: % of TFA. Values for limited irradiance = mean ± SEs of TFA, specific FA or FA
482 groups at different growth phases (with n = 18 replicates for initial phase and n = 14 replicates for
483 the continuous phase). Different letters in the same row denote the significant differences (Tukey
484 test) in the TFA, or the same specific FA or FA groups between the different growth phases.

485

486 **Table 2**

Variables	df1	df2	SS	MS	F	<i>p</i>
TFAA	1	31	0.0043	0.0043	0.003	0.958
TEAA	1	31	0.149	0.149	0.325	0.573
His	1	31	0.0059	0.0059	0.0176	0.895
Thr	1	31	1.92	1.92	1.193	0.283
Arg	1	31	34.195	34.195	1.017	0.321
Val	1	31	2.257	2.257	0.319	0.576
Met	1	31	1.699	1.699	0.876	0.357
Ile	1	31	0.0421	0.0421	0.0317	0.86
Leu	1	31	1.878	1.878	2.084	0.159
Lys	1	31	0.0205	0.0205	0.00494	0.944
Phe	1	31	3.502	3.502	2.258	0.143

487

488


489 **Table 3**

Estimated parameters	
Dilution rate, D (day^{-1})	0.46
Total volume of the two culture columns (L)	94.0
Actual filled volume of the two culture columns, V_A (L)	84.0
Density of the produced algae (DA , 10^9 cells L^{-1})	2.40
Algae production (P , 10^9 cells day^{-1})	92.74
Algae production (P , g C day^{-1})	4.40
The algae requirement for 1 L of copepods ($2500 \text{ ind } \text{L}^{-1}$) per day (CA , g C $\text{L}^{-1} \text{ day}^{-1}$)	0.0089
The volume of copepod culture that can be sustained by feeding on the current algae production (V_C , L)	495
Specific egg production (SEP , egg $\text{female}^{-1} \cdot \text{day}^{-1}$)	30.70
Potential number of eggs produced with the current algal production (EP , $\times 10^6$ eggs day^{-1})	17.11


490

491

492



495 **Figure 2**




496

497

498 **Figure 3**


499



500

501

502 **Figure 4**



503