Effects of combined exposures of fluoranthene and polyethylene or polyhydroxybutyrate microplastics on oxidative stress biomarkers in the blue mussel (*Mytilus edulis*)

Magara, Gabriele; Khan, Farhan; Pinti, Marika ; Syberg, Kristian; Inzirillo, Angelo ; Elia, Antonia Concetta

Published in:
Journal of Toxicology and Environmental Health. Part A

DOI:
[10.1080/15287394.2019.1633451](https://doi.org/10.1080/15287394.2019.1633451)

Publication date:
2019

Document Version
Peer reviewed version

Citation for published version (APA):

Magara, G., Khan, F., Pinti, M., Syberg, K., Inzirillo, A., & Elia, A. C. (2019). Effects of combined exposures of fluoranthene and polyethylene or polyhydroxybutyrate microplastics on oxidative stress biomarkers in the blue mussel (*Mytilus edulis*). *Journal of Toxicology and Environmental Health. Part A*, 82(10), 616-625.
<https://doi.org/10.1080/15287394.2019.1633451>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

Effects of combined exposures of fluoranthene and polyethylene or polyhydroxybutyrate microplastics on oxidative stress biomarkers in the blue mussel (*Mytilus edulis*)

Journal:	<i>Journal of Toxicology and Environmental Health, Part A: Current Issues</i>
Manuscript ID	UTEH-2019-0151.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Magara, Gabriele; Universita degli Studi di Perugia Khan, Farhan; Roskilde University, Department of Science and Environment Pinti, Marika; Universita degli Studi di Perugia Syberg, Kristian; Roskilde University, Department of Science and Environment Inzirillo, Angelo; Universita degli Studi di Perugia Elia, Antonia; Universita degli Studi di Perugia,
Keywords:	biodegradable polymer, biomarker profile, bioplastics, blu mussels, microplastic

SCHOLARONE™
Manuscripts

1

2 1 **Effects of combined exposures of fluoranthene and polyethylene or**

3

4 2 **polyhydroxybutyrate microplastics on oxidative stress biomarkers in the blue**

5

6 3 **mussel (*Mytilus edulis*)**

7

8

9 4 *Gabriele Magara^a, Farhan R. Khan^{b*}, Marika Pinti^a, Kristian Syberg^b, Angelo Inzirillo^a*

10

11 5 *and Antonia Concetta Elia^{a*}*

12

13

14 6 *a Department of Chemistry, Biology and Biotechnology, via Elce di sotto 8, University of Perugia,*

15

16 7 *06123 Perugia, Italy*

17

18 8 *b Department of Science and Environment, Universitetsvej 1, P.O. Box 260, Roskilde University, DK-*

19

20 9 *4000 Roskilde, Denmark*

21

22

23 10 Running title: Bioplastics as a vector for environmental pollutants

24

25

26 11

27

28 12 ** Corresponding author: Dr. Antonia Concetta Elia*

29

30

31 13 *Department of Chemistry,*

32

33

34 14 *Biology and Biotechnology,*

35

36

37 15 *University of Perugia, Italy*

38

39 16 *Email: antonia.elia@unipg.it*

40

41

42 17 *Dr. Farhan R. Khan*

43

44

45 18 *Department of Science and Environment,*

46

47

48 19 *Roskilde University, Denmark*

49

50 20 *Email: frkhan@ruc.dk, farhan.khan@gmx.com*

51

52

53 21

54

55 22

56

57 23

58

59

60

ABSTRACT

A growing interest in developing and commercialization of new ecofriendly plastic polymers is occurring attributed to the impact of marine plastics debris and microplastics that result from the degradation of oil-based polymers as these substances adversely affect ecosystem health. Recently, polyhydroxybutyrate (PHB) has become of interest due its biodegradability and physicochemical properties. However, biological consequences resulting from bioplastics exposure remains to be determined. Further, few data are apparently available regarding the potential for bioplastics to act as a vector for exogenous chemicals in the environment. The aim of the study was to compare the effects of polyethylene (PE MPs) and polyhydroxybutyrate (PHB MPs) microplastics administered alone or in combination with fluoranthene (Flu) on detoxifying enzymes in digestive glands and gills of *Mytilus edulis*. Blue mussels were exposed for 96hr to 8 experimental groups: control, Flu-only, PE MPs-only, PHB MPs-only, PE MPs-Flu co-exposure, PHB MPs-Flu co-exposure, Flu-incubated PE MPs and Flu-incubated PHB MPs. Superoxide dismutase (SOD), catalase (CAT), glutathione peroxidases (GPx), glutathione S-transferase (GST) and glutathione reductase (GR) were found to be significantly susceptible to Flu and plastics in both tissues. Interestingly, single exposure to PHB MPs led to decreased levels of CAT and GST in gills, SOD in digestive glands and SeGPx in both tissues. In co-exposure and incubation treatments, biochemical responses were generally comparable with those exerted by PE MPs or PHB MPs only, suggesting an apparent absence of combined effects with pollutant. Data demonstrate the ecotoxicological impact of bioplastics materials on digestive glands and gills of *Mytilus edulis*.

Keywords: biodegradable polymer, biomarker profile, bioplastics, blue mussels, microplastic

1

2 48 **Introduction**

3

4 49 Oil-based plastic debris and subsequently the microplastics (MPs) formed by their degradation are

5

6 50 one of the current major environmental concerns. These plastics are defined chemically as organic or

7

8 51 semi-organic materials with long chains (macromolecules) and high molecular weight, generated by the

9

10 52 polymerization of monomers extracted from oil or gases (Cole et al., 2011). Once in seawater,

11

12 53 microplastics were shown to be ingested by aquatic species which resulted in a number of deleterious

13

14 54 effects (Guzzetti et al., 2018; Alimba and Faggio, 2019) . The color, density, shape and size of MPs

15

16 55 often leads marine organisms to mistake MPs for food items and the consequences of this includes

17

18 56 blockage of the digestive system and false feeling of satiation (Wright et al., 2013). In addition, MPs

19

20 57 may act as a vehicle for transport of several contaminants dissolved in the aquatic environment

21

22 58 (Andrady, 2011). This ‘vector effect’ (Syberg et al., 2015), suggests that oil-based MPs may trap and

23

24 59 transport environmental pollutants including persistent organic pollutants (POPs) or metals (Oliveira et

25

26 60 al., 2013; Avio et al., 2015; Khan et al., 2015; Magara et al., 2018). The possibility that plastics might

27

28 61 then transport the adhered contaminants into aquatic biota is currently being debated (Koelmans et al.,

29

30 62 2016), and lab investigations into the vector effect remain ongoing.

31

32

33

34

35

36 63 There is a mounting interest in “green materials” and a focus on development and production of

37

38 64 biodegradable polymers (Mohanty et al., 2002). In contrast to oil-based polymers, these biopolymers

39

40 65 are derived from bioprocesses, using renewable resource in bio-refineries. Due to their mechanical and

41

42 66 physicochemical characteristics, these biodegradable polymers may be considered as an eco-friendly

43

44 67 substitutes to plastic (Anjum et al., 2016). In recent years, polyhydroxybutyrate (PHB) has received

45

46 68 growing interest for its characteristics similar to oil-based plastic (Dacosta et al., 2016).

47

48 69 Polyhydroxybutyrate is water insoluble, but highly biodegradable biopolymer, relatively resistant to

49

50 70 hydrolytic degradation which is extracted with chloroform from bacterial cultures grown on

51

52 71 carbohydrates. PHB (1) displays low oxygen permeability and reliable thermoplasticity with poor

53

54 72 mechanical properties, (2) high crystallinity degree and (3) is optically active with a mass about

55

56

57

58

59

60

approximately 0.5×10^6 Da (Anjum et al., 2016). However, although PHB is currently available commercially, especially as main component of biodegradable market bags, to the best of our knowledge little is known regarding the environmental impact or effects on aquatic organisms. By sharing many characteristics with oil-based plastics, it is feasible that in the aquatic environment, bioplastic undergoes modifications, resulting in generation of MPs as is the case for oil-based polymers (Andrady, 2011). If due to environmental and societal pressures a move is made towards use of bioplastics, then it is vital to gather information on the influence of these bioplastics on the ecosystem health.

Oxidative stress results from an imbalance between pro-oxidants such as reactive oxygen species (ROS), and the protective antioxidant system. Mechanisms that involve glutathione (GSH), and the related antioxidant enzymes superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidases (SeGPx and GPx) represent important protective metabolic pathways which are used as biomarkers related to pollutant induced oxidative stress. Glutathione (GSH) in its reduced form is an important non-enzymatic scavenger of oxyradicals, involved in the metabolism of toxic compounds and endogenous substances (Meister and Anderson, 1983). Both SOD and CAT enzymes catalyze the breakdown of ROS-generating O_2^- and H_2O_2 , respectively and are key components within the primary defense system against oxidative stress-induced damage. SeGPx enzyme reduces either hydrogen peroxide or organic peroxides and GPx enzyme affects organic hydroperoxides. Polycyclic aromatic hydrocarbons (PAHs), of which fluoranthene (Flu) is often used as a model contaminant (and in addition a priority aquatic pollutant (Directive 2008/105/EC)), are known to initiate oxidative stress which was assessed by these aforementioned biomarkers (Cheung et al., 2001). The oxidative stress biomarkers employed in the present study were utilized to determine the impact of a range of xenobiotics on aquatic organisms (Al kaddissi et al., 2012; 2016; Cozzari et al., 2015; Elia et al., 2017a; 2017 b; 2018; Magara et al., 2018). Previous studies utilizing these biomarkers to elucidate the

MP vector effect demonstrated a perturbation of intracellular antioxidant defenses (Oliveira et al., 2013; Avio et al., 2015; Magara et al., 2018).

Previously Magara et al. (2018) investigated the influence of polyethylene microplastic beads (PE MPs) on the accumulation and associated oxidative stress responses attributed to Flu in blue mussels, *Mytilus edulis*. The mussels were exposed to 4 treatment groups: Flu-only, MP-only, Flu and MP co-exposure, and Flu-incubated MP. Individual contaminant exposures to Flu or MP resulted in varying responses, but co-exposures and incubated treatments (i.e. Flu was sorbed to plastic surfaces prior to exposure) did not induce additive or synergistic responses. Further, MP-only exposure appeared to be capable of eliciting direct effects on the oxidative stress system as evidenced by enhanced activities of CAT and GPx (Magara et al., 2018).

Thus, the present study was designed to investigate the potential impact of PHB microplastics as a single contaminant and in combination with Flu on the oxidative stress system of blue mussel (*Mytilus edulis*), a well-regarded test species for MPs (Avio et al., 2015; Paul-Pont et al., 2016; Sahlmann et al. 2017; Von Moos, 2012). PE MPs, as an example of oil-based MPs, were employed as a comparison to PHB MPs. The aim of the study was to examine the comparative changes on oxidative stress biomarkers levels induced by oil based (PE) MPs and ‘new’ bioplastic PHB MPs as both single contaminants and as potential vectors for PAHs.

Materials and methods

Mussels collection and maintenance

Blue mussel (*Mytilus edulis*) were collected in June, 2016, in Vellerup Vig – Isefjord, N 55°40.6'; E 11°48.7', in an non-polluted deep inlet of Baltic Sea near Roskilde (Denmark). The mussels were maintained in a temperature-controlled room (10 ± 1°C). All specimens were placed in tanks (size 39 x 21 x 25 cm), filled with 10 L of field water. Field water was gradually replaced with lab water

maintaining similar salinity (20 salinity). Mussels were fed for the following three weeks twice weekly. A 16:8 hr light/dark photoperiod was maintained throughout.

Chemicals and preparation

A primary stock solution of Flu (Fluka Chemika, Steinheim, Switzerland) was prepared at 1 mg/ml in acetone. PE MPs were purchased from Cospheric LLC (Santa, Barbra, CA, USA) and PHB beads provided by Prodotti Gianni S.R.L. (Milano, Italy). Both polymers were white in color and similar in size (10–90 μm). Both MPs were treated with 0.1% Tween80 to enable dispersion in water column (Khan et al., 2015).

To prepare each exposure treatment accurately a weight to MP number ratio was determined for both PE and PHB particles. The weight associated to 1×10^6 MPs was weighed out for each treatment group containing MPs. The MPs were dispersed in 5 ml of 0.1% Tween80 solution and shaken at 150 rpm overnight. The MPs were filtered through 1 μm nylon mesh and resuspended in 200 ml 20 salinity water and kept shaking until the start of the experiment (150 rpm). MPs that required incubation with Flu were resuspended into 20 ml acetone to which the appropriate amount of Flu stock was added to yield the correct final concentration of 100 $\mu\text{g/L}$ when made up to the final exposure solution. After an initial mixing, dispersions were left under a fume hood for the acetone to evaporate and the Flu sorb to the PE MPs. This method was adopted from use with other particulate contaminants (Al-Subiai et al., 2012). When complete dryness was achieved overnight the Flu-incubated MPs were resuspended in 200 ml 20 salinity water and placed with the other dispersions on the shaking table overnight.

Mussels exposure

Mussels were exposed for 96 hr to 8 treatment groups as follows, (A) Control (no added contaminants), (B) Flu only, (C) PE MPs only, (D) PHB MPs only, (E) PE MPs-Flu co-exposure, (F) PHB MPs-Flu co-exposure, (G) Flu-incubated PE MPs and (H) Flu-incubated PHB MPs (n=4 per treatment). The Flu and MP concentrations across all treatments were 100 $\mu\text{g/L}$ and 1000 MPs/ml,

1
2 145 respectively. Although the concentrations exceeded environmental realistic conditions , the aim of this
3
4 146 study was to understand mechanistic responses that are more easily captured at higher exposure levels.
5
6 147 The concentrations used are consistent with previous studies (Al-Subiai et al.,2012; Cole et al., 2016;
7
8
9 148 Magara et al., 2018). Acetone and Tween80 were consistently added across treatments to prevent the
10
11 149 confounding influence of solvent and detergent. Ninety-six hr mussel exposures were conducted at
12
13 150 10±1°C and a 16/8 hr light/dark photoperiod. Mussels were individually exposed in aerated beakers
14
15
16 151 (250 ml) filled with 200 ml treatment solution. Four samples of two pooled mussels were exposed per
17
18 152 treatment group.

19
20 153 To start the experiment, each 200 ml dispersion was made up to 1 L with 20 salinity water and
21
22
23 154 vigorously stirred to ensure homogenous distribution of MPs in the water column. The 1 L dispersion
24
25 155 was then divided into 5 200 ml units again ensuring a homogenous distribution of MPs between
26
27 156 beakers. The appropriate amount of Flu stock was added to the Flu-only and co-exposures groups prior
28
29
30 157 to the start of the experiment. Post exposure (96 hr), mussels were removed and rinsed thoroughly
31
32 158 under 20 salinity water. Gill and digestive gland tissues were extracted from each individual and
33
34 159 placed in separate pre-weighed and pre-labelled vials. Tissues for biomarkers and Flu concentration
35
36 160 analysis were stored at -80°C and -20° C, respectively. No mortalities were recorded during the
37
38
39 161 experiments. .

40
41
42 162 **Biochemical analysis**

43
44 163 The analysis of biomarkers is described in detail elsewhere (Cozzari et al., 2015; Magara et al.,
45
46 164 2018). Briefly, total GSH levels as well as activities of SOD, CAT, GPx and SeGPx were measured in
47
48
49 165 cytosolic fraction of gills and digestive gland spectrophotometrically (Varian Cary 50
50
51 166 spectrophotometer at 25°C). Protein concentrations in the cytosol were determined according to Lowry
52
53 167 et al. (1951) and used to normalize enzyme levels. All analyses were performed in triplicate along with
54
55
56 168 blank samples (buffer and reagents only). These absorbance values were subtracted from those of the
57
58
59

sample. SOD activity was determined according to an established method (McCord and Fridovich, 1969). Catalase levels were measured following the decrease in absorbance due to H_2O_2 consumption at 240 nm (Greenwald, 1985). GPx activities (SeGPx and total GPx) were determined following the oxidation of nicotinamide adenine dinucleotide phosphate reduced form (NADPH) at 340 nm (Lawrence and Burk, 1976). GR activity was measured in 100 mM NaH_2PO_4 + Na_2HPO_4 buffer, pH 7, 1 mM oxidized glutathione (GSSG), and 60 μM NADPH; the decrease in absorbance due to the oxidation of NADPH was measured at 340 nm (Chung et al. 1991). GST activity was determined in 100 mM NaH_2PO_4 + Na_2HPO_4 buffer, pH 6.5, 1 mM GSH, 1 mM 1-chloro-2,4 dinitrobenzene (CDNB) as substrates and sample. Formation of the conjugate with GSH was read at 340 nm (Habig et al. 1974). Total glutathione (GSH+2GSSG) was measured following homogenization in 5% sulphosalicylic acid with 4mM EDTA by the GR recycling assay at 412 nm (Akerboom and Sies, 1981).

Statistical analysis

Data are reported as mean values $\pm$ the standard deviation (SD). Levene's test for normal distribution was performed on datasets prior to statistical analysis. A one-way ANOVA with Bonferroni's multiple comparison test was used to investigate differences between treatment groups (GraphPad Prism software). Statistical significance was set at $P < 0.05$.

Results

In the gills, CAT activity levels significantly decreased in all treatment groups (up to 80%), except for PE MPs only exposure compared to control (Figure 1C). The activity of SeGPx was markedly lower (up to 90%) through the treatments (Figure 2C). GST levels in the treated groups were significantly lower than control (up to 50%), except for Flu-incubated PHB MPs (Figure 3A). No

marked changes of SOD, GPx and GR activities were noted for all treatment compared to respective controls (Figures 1A, 2A, 3C).

In the digestive glands, SOD activity was significantly reduced (up to 50%) in all treatments when compared to control with the exception of the group incubated with to PE MPs only (Figure 1B). CAT activity was markedly decreased similar to SOD (up to 60%)., However, no significant differences were found in mussels exposed to PHB MPs only and Flu-incubated PHB MPs (Figure 1D). Total GPx markedly fell (60%) following Flu exposure (Figure 2B). Levels of SeGPx activity were significantly reduced compared to control (up to 50%), except for PE MPs only and Flu-incubated PHB MPs groups (Figure 2D). GST activity peaked (1.5 fold higher) in Flu exposed group (Figure 3B), whereas GR activity levels showed increasing trends in all treated groups. The GR activity was markedly higher following PE MPs only exposure (1 fold), PE MPs-Flu co-exposure (1.2 fold) and Flu-incubated PHB MPs treatment (1 fold, Figure 3D).

Discussion

The present study provides the first experimental evidence of stress-related effects initiated by novel bio-microplastic exposure, either as a single contaminant or in combination with Flu, in the blue mussel *Mytilus edulis*. The comparison was conducted with the more traditional oil-based plastic polyethylene. Flu and both PE and PHB MPs modified the antioxidant responses in both gills and digestive glands. The present results are largely in agreement with those previously reported for Flu and PE MPs, both in single or combined exposures (Magara et al., 2018).

PE MPs altered the levels of several of the tested biomarkers. Although previous studies showed that oil-based plastic led to enhanced antioxidant activities in copepods (Jeong et al., 2017) and fish (Barboza et al., 2018), our results are in agreement with studies on mussels, showing a significant inhibition of CAT, SeGPx and GST activity following oil-based MPs mussel exposure (Avio et al.,

2015; Paul-Pont et al., 2016). A decrease of CAT and GPx activities in digestive glands and of SeGPx and GST in gills exposed to PE MPs may be related to the size and shape of microplastics, that play a key role in initiation of biological changes (Browne et al., 2008; Von Moos, 2012; Avio et al., 2015). Small polystyrene plastic particles may be translocated from the gut to the circulatory system of *Mytilus edulis* and subsequently retained suggesting that these substances might reach crucial organs such as the heart or hepatic tissues to produce adverse effects (Browne et al., 2008). It is well known that oil-based microplastics may accumulate within organisms and induce tissue abrasions, as evidenced by histological changes in digestive cells and triggered inflammatory responses, formation of granulocytomas and lysosomal destabilization (Avio et al., 2015; Von Moos, 2012). Therefore, in this scenario, the imbalance of antioxidant defense markers might be expected and might be attributed to physical damage induced by MPs enabling further accumulation of reactive products. Further, in the present study increased GR activity in digestive glands might reflect the demand for GSH as defense mechanism against oxidative stress induced by oil-based MPs (Jeong et al., 2016).

The present study also carried out a co-exposure and incubation scenario of both PE and PHB MPs with Flu and reported information on oxidative stress related to the simultaneous exposure to compounds. In this scenario, a mussel experimental group exposed to Flu-only was used as positive control. Previously Pan et al (2005) found that at high benzo(k)fluoranthene, a PAH, concentrations the activity of antioxidant enzymes was diminished in the scallop *Chlamys farreri* in a time-dependent manner. Our findings that the marked decrease of SOD activity levels and both GPx in digestive glands, as well as SeGPx in gills and CAT activity in both tissues indicate that this Flu may exert detrimental effects when administered alone resulting in a severe oxidative stress in *Mytilus edulis*. The impairment of antioxidant biomarkers levels may be attributed to a chemical-mediated damage. Indeed, the enhanced GST activity measured in digestive glands might represent an important outcome, since this indicate elevated of phase-II biotransformation metabolism. The significant induction of GST activity in mussels exposed to Flu-only may be attributed to production of specific

1
2 239 substrates for GST by phase I enzymes, representing an effective defense line against Flu. Babson et al
3
4 240 (1986) showed that rat liver microsomes converted Flu to trans-2,3-dihydrodiol as the major
5
6 241 metabolite. Although data regarding Flu metabolites in blue mussels are lacking, it is suggested that
7
8
9 242 GST induction may be related to production of this intermediate.

10
11 243 Biomarkers responses in digestive glands and gills of mussels exposed to both Flu and PE or PHB
12
13 244 MPs (co-exposure and incubated treatment) generally followed a trend similar to PAH alone or the
14
15
16 245 new plastic alone and no combined effects of these agents was apparent. The absence of combined
17
18 246 effects in *Mytilus edulis* exposed to PE MPs and Flu is in agreement with our previous results
19
20 247 (Magara et al., 2018), suggesting that MPs may act as a “sink” of environmental pollutants (Chua et al.
21
22
23 248 2014; Khan et al. 2015). Chua et al. (2014), showed that the concentration of polybrominated diphenyl
24
25 249 ethers (PBDE) was lower in *Allorchestes compressa* co-exposed to microplastics then specimens
26
27 250 treated with PBDE alone. Chua et al (2014) concluded that the presence of MPs may inhibit the
28
29
30 251 uptake of PBDE, perhaps because this contaminant is strongly absorbed onto microplastics surface
31
32 252 making it less bioavailable. It is already known that oil-based micropolymers have the propensity to
33
34 253 aggregate in water (Alimi et al., 2018) . Previously Khan et al (2015) found in zebrafish a decrease in
35
36
37 254 silver (Ag) uptake by fish attributed to diminished contact between Ag and tissues following oil-based
38
39 255 MP aggregation. Further, Magara et al. (2018) reported that Flu tissue concentrations were lower in
40
41 256 *Mytilus edulis* exposed to both co-exposure with polyethylene and PAH compared to specimens
42
43 257 treated with Flu alone. The absence of combined effects of PE MPs and Flu in *Mytilus edulis* and
44
45
46 258 previous results on Flu uptake in blue mussel (Magara et al., 2018), suggests that the interaction of
47
48 259 tissues with Flu might be delayed by this aggregation mechanism exerted by microplastics.

49
50 260
51
52 261 Bioplastics are currently becoming the leading material for replacing oil-based polymers since it is
53
54
55 262 more desirable than traditional plastic due to a propensity to biodegrade in the environment (Anjum et
56
57 263 al., 2016). Recently Napper and Thompson (2019) demonstrated that biodegradable polymers may

not undergo any substantial deterioration over a 3 year period in marine environment, but may be reduced in small fragments similar to oil-based plastics. Our data suggest that PHB MPs results in altered levels of some oxidative stress biomarkers similar to oil-based MPs. The fact that bioplastics may exert stress effects and the potential low biodegradability rate indicate a novel threat for the marine environment. Therefore, it is crucial to understand degradation processes of bioplastics in the environment, but also gathering knowledge regarding potential ecotoxicological impacts. Although the damage exerted by oil-based polymers is becoming increasingly understood, it is still necessary to investigate the possible consequences of bioplastics exposure both alone and in combination with other environmental pollutants present in the aquatic ecosystem.

Conclusions

The results of the present study demonstrated that PHB MPs modify the baseline levels of biomarkers related to oxidative stress in *Mytilus edulis*. These alterations were similar to those exerted by PE MPs treatments for some of the antioxidant biomarkers. Further, the responses measured in both co-treatments were similar to those noted with MP-alone exposures. Data indicated the absence of any combined effects produced by oil-based MPs or bioplastics MPs and Flu.

Acknowledgements

The visit of A. Inzirillo to Department of Science and Environment, Roskilde University, was funded by a LLP/Erasmus Placement program (2015/2016). The Authors thank Prodotti Gianni S.R.L. (Milano, Italy) company kindly providing the PHB beads.

References

- Al Kaddissi, S., O. Simon, A.C. Elia, P. Gonzalez, M. Floriani, I. Cavalie, V. Camilleri, S. Frelon, and A. Legeay. 2016. How toxic is the depleted uranium to crayfish *Procambarus clarkii* compared

- with cadmium? *Environmental Toxicology* 31:211–223.
- Al Kaddissi S., S. Frelon, A.C. Elia, A. Legeay, P. Gonzalez, F. Coppin, D. Orjollet, V. Camilleri, K. Beaugelin-Seiller, R. Gilbin, and O. Simon. 2012. Are antioxidant and transcriptional responses useful for discriminating between chemo- and radiotoxicity of uranium in the crayfish *Procambarus clarkii*? *Ecotoxicology and Environmental Safety* 80: 266-272.
- Alimba, C., and C. Faggio. 2019. Microplastics in the marine environment: current trends in environmental pollution and mechanisms of toxicological profile. *Science of the Total Environment* 68: 61-74.
- Alimi, O. S., J.Farner Budarz, L. M.,Hernandez, and N. Tufenkji. 2018. Microplastics and nanoplastics in aquatic environments: aggregation, deposition, and enhanced contaminant transport. *Environmental Science and Technology* 52: 1704-1724.
- Al-Subiai, S. N., V. M. Arlt, P. E. Frickers, J. W. Readman, B. Stolpe, J. R. Lead, A. J. Moody, and A. N. Jha. 2012. Merging nano-genotoxicology with eco-genotoxicology: An integrated approach to determine interactive genotoxic and sub-lethal toxic effects of C 60 fullerenes and fluoranthene in marine mussels, *Mytilus* sp. *Mutation Researcher Genetic Toxicology and Environment Mutagenesis* 745: 92–103.
- Akerboom, T.P.M., and H. Sies. 1981. Assay of glutathione disulfide and glutathione mixed disulfide in biological samples. *Methods in Enzymology* 71: 373–382.
- Andrady, A.L. 2011. Microplastics in the marine environment. *Marine Pollution Bulletin* 62: 1596–1605.
- Anjum, A., M. Zuber, K.M. Zia, A. Noreen, M.N. Anjum, and S. Tabasum. 2016. International Journal of Biological Macromolecules Microbial production of polyhydroxyalkanoates (PHAs) and its copolymers: A review of recent advancements. *International Journal of Biological Macromolecules* 89:161–174.
- Avio, C.G., S. Gorbi, M. Milan, M. Benedetti, D. Fattorini, G. d'Errico, M. Pauletto, L. Bargelloni, and

- F. Regoli. 2015. Pollutants bioavailability and toxicological risk from microplastics to marine mussels. *Environmental Pollution* 198: 211–222.
- Babson, J.R., S.E. Russo-Rodriguez, R.V. Wattley, P.L. Bergstein, W.H. Rastetter, H.L. Liber, B.M. Andon, W.G. Thilly, and G.N. Wogan. 1986. Microsomal activation of fluoranthene to mutagenic metabolites. *Toxicology and Applied Pharmacology* 85: 355–366.
- Barboza, L.G.A., L. R. Vieira, V. Branco, C. Carvalho, and L. Guilhermino. 2018. Microplastics increase mercury bioconcentration in gills and bioaccumulation in the liver, and cause oxidative stress and damage in *Dicentrarchus labrax* juveniles. *Scientific Reports* 8: 15655
- Browne, M.A.D., T.S. Galloway, D.M. Lowe, and R.C. Thompson. 2008. Ingested microscopic plastic translocates to the circulatory system of the mussel, *Mytilus edulis* (L). *Environmental Science and Technology* 42: 5026–5031.
- Cheung, C.C.C., G.J. Zheng, A.M.Y. Li, B.J. Richardson, and P.K.S. Lam. 2001. Relationships between tissue concentrations of polycyclic aromatic hydrocarbons and antioxidative responses of marine mussels, *Perna viridis*. *Aquatic Toxicology* 52:189–203.
- Chua, E.M., J. Shimeta, D. Nugegoda, P.D. Morrison, and B.O. Clarke. 2014. Assimilation of polybrominated diphenyl ethers from microplastics by the marine amphipod, *Allorchestes compressa*. *Environmental Science and Technology* 48: 8127–8134.
- Chung, P.M., R.E. Cappel, and H.F. Gilbert. 1991. Inhibition of glutathione disulfide reductase by glutathione. *Archives of Biochemistry and Biophysics* 288: 48–53.
- Cole, M., P. Lindeque, C. Halsband, and T.S. Galloway. 2011. Microplastics as contaminants in the marine environment: A review. *Marine Pollution Bulletin* 62: 2588–2597.
- Cole, M., P.K. Lindeque, E. Fileman, J. Clark, C. Lewis, C. Halsband, and T.S. Galloway. 2016. Microplastics alter the properties and sinking rates of zooplankton faecal pellets. *Environmental Science and Technology* 50: 3239–3246.
- Cozzari, M., A.C. Elia, N. Pacini, B.D. Smith, D. Boyle, P.S. Rainbow, and F.R. Khan. 2015.

1
2 339 Bioaccumulation and oxidative stress responses measured in the estuarine ragworm (*Nereis*
3
4 340 *diversicolor*) exposed to dissolved, nano- and bulk-sized silver. *Environmental Pollution* 198:
5
6 341 32-40.
7
8
9 342 Dacosta, C.F., J.A. Posada, and A. Ramirez. 2016. Techno-economic and carbon footprint assessment
10
11 343 of methyl crotonate and methyl acrylate production from wastewater-based polyhydroxybutyrate
12
13 344 (PHB). *Journal of Cleaner Production* 137: 942-952.
14
15
16 345 Directive 2008/105/EC of the European Parliament and of the Council. Official Journal of the
17
18 346 European Union.
19
20 347 Elia, A.C., G. Magara, C. Caruso, L. Masoero, M. Prearo, P. Arsieni, B. Caldaroni, M. Scoparo, A.J.M.
21
22
23 348 Dörr, S. Salvati, P. Brizio, S. Squadrone, and M.C. Abete. 2018. A comparative study on
24
25 349 subacute toxicity of arsenic trioxide and dimethylarsinic acid on antioxidants and antioxidant-
26
27 350 related enzymes in Crandell Rees feline kidney (CRFK), human hepatocellular carcinoma
28
29 351 (PCL/PRF/5) and epithelioma *Papulosum cyprini* (EPC) cell lines. *Journal of Toxicology and*
30
31 *Environmental Health A* 81: 333-348.
32 352
33
34 353 Elia, A. C., G. Magara, M. Righetti, A. J. M. Dörr, T. Scanzio, N. Pacini, M. C. Abete, and M. Prearo.
35
36 354 2017a. Oxidative stress and related biomarkers in cupric and cuprous chloride-treated rainbow
37
38
39 355 trout. *Environmental Science and Pollution Research* 24:10205–19.
40
41 356 Elia, A.C., F. Giorda, N. Pacini, A.J.M. Dörr, T. Scanzio, and M. Prearo. 2017b. Subacute toxicity
42
43 357 effects of deltamethrin on oxidative stress markers in rainbow trout.
44
45
46 358 *Journal of Aquatic Animal Health* 29:165-172.
47
48 359 Greenwald, R.A. 1985. *Handbook of Methods for Oxygen Radicals Research*. CRC Press, Boca
49
50 360 Raton, FL. Ed. by R. A. Greenwald, pp 447.
51
52
53 361 Guzzetti, E., A. Sureda, S.Tejada, and C. Faggio. 2018. Microplastic in marine organism:
54
55 362 environmental and toxicological effects. *Environmental Toxicology and Pharmacology* 64: 164-
56
57 363 171.
58
59
60

- Habig, W.H., M.J. Pabst, and W.B. Jakoby. 1974. Glutathione S transferases. The first enzymatic step in mercapturic acid formation. *Journal of Biological Chemistry* 249: 7130–7139.
- Jeong, C.B., H.M. Kang, M.C. Lee, D.H. Kim, J. Han, D.S. Hwang, S. Souissi, S.J. Lee, K.H. Shin, H.G. Park, and J.S. Lee. 2017. Adverse effects of microplastics and oxidative stress-induced MAPK/Nrf2 pathway-mediated defense mechanisms in the marine copepod *Paracyclops nana*". *Scientific Reports* 7: 41323.
- Jeong, C., E. Won, H. Kang, M. Lee, D. Hwang, and U. Hwang. 2016. Microplastic size-dependent toxicity, oxidative stress induction, and p-JNK and p-p38 activation in the monogonont rotifer (*Brachionus koreanus*). *Environmental Science and Technology* 50: 8849-8857.
- Khan, F.R., K. Syberg, Y. Shashoua, and N.R. Bury. 2015. Influence of polyethylene microplastic beads on the uptake and localization of silver in zebra fish (*Danio rerio*). *Environmental Pollution* 206: 73–79.
- Koelmans, A. A., A. Bakir, G. A. Burton, and C.R. Janssen. 2016. Microplastic as a vector for chemicals in the aquatic environment: Critical review and model-supported reinterpretation of empirical studies. *Environmental Science and Technology* 50: 3315–3326.
- Lawrence, R.A., and R.F. Burk. 1976. Glutathione peroxidase activity in selenium-deficient rat liver. *Biochemical and Biophysical Research Communications* 71: 592–598.
- Lowry, O.H., N.J. Rosebrough, A.L. Farr, and R.J. Randall 1951. Protein measurement with the Folin phenol reagent. *Journal of Biological Chemistry* 193: 265–275.
- Madison, L.L., and G.W. Huisman. 1999. Metabolic engineering of poly (3-hydroxyalkanoates): From DNA to plastic. *Microbiology and Molecular Biology Review* 63: 21–53.
- Magara, G., A.C. Elia, K. Syberg, and F.R. Khan. 2018. Single contaminant and combined exposures of polyethylene microplastics and fluoranthene: accumulation and oxidative stress response in the blue mussel, *Mytilus edulis*. *Journal of Toxicology and Environmental Health A* 81: 761–773.

- McCord, J.M., and I. Fridovich. 1969. Superoxide dismutase: an enzymatic function for erythrocuprein (hemocuprein). *Journal of Biological Chemistry* 244:6049–6055.
- Meister, A., and M.E. Anderson. 1983. Glutathione. *Annual Review of Biochemistry* 52:711–60.
- Mohanty, A.K., M. Misra, and L.T. Drzal. 2002. Sustainable bio-composites from renewable resources: Opportunities and challenges in the green materials world. *Polish Journal of Environmental Studies* 10: 19–26.
- Napper, I.E., and R.C. Thompson. 2019. Environmental deterioration of biodegradable, oxo-biodegradable, compostable, and conventional plastic carrier bags in the sea, soil, and open-air over a 3-year period. *Environmental Science and Technology* 53: 4775-4783.
- Oliveira, M., A. Ribeiro, K. Hylland, and L. Guilhermino. 2013. Single and combined effects of microplastics and pyrene on juveniles (0+ group) of the common goby *Pomatoschistus microps* (Teleostei, Gobiidae). *Ecological Indicators* 34: 641–647.
- Pan, L., J. Ren, and J. Liu. 2005. Effects of benzo(k)fluoranthene exposure on the biomarkers of scallop *Chlamys farreri*. *Comparative Biochemistry and Physiology C* 141:248–256.
- Paul-Pont, I., C. Lacroix, C. Gonzalez Fernandez, H. Helene, C. Lambert, N. Le Goic, L. Frère, A.L. Cassone, R. Sussarellu, C. Fabioux, J. Guyomarch, M. Albentosa, A. Huvet, and P. Soudant. 2016. Exposure of marine mussels *Mytilus* spp. to polystyrene microplastics: Toxicity and influence on fluoranthene bioaccumulation. *Environmental Pollution* 216:724–737.
- Sahlmann, A., R. Wolf, T. F. Holth, J. Titelman, and K. Hylland. 2017. Baseline and oxidative DNA damage in marine invertebrates. *Journal of Toxicology and Environmental Health A* 80: 807-819.
- Syberg, K., F. R. Khan, H. Selck, A. Palmqvist, G.T. Banta, J. Daley, L. Sano, and M.B. Duhaime. 2015. Microplastics: Addressing ecological risk through lessons learned. *Environmental Toxicology and Chemistry* 34: 945–953.
- Verlinden, R.A.J., D.J. Hill, M.A. Kenward, C.D. Williams, and I. Radecka. 2007. Bacterial synthesis of biodegradable polyhydroxyalkanoates. *Journal of Applied Microbiology* 102: 1437–1449.

- Von Moos, N.B.P. 2012. Uptake and effects of microplastics on cells and tissue of the blue mussel *Mytilus edulis* L. after an experimental exposure. *Environmental Science and Technology* 46: 11327-11335.
- Wright, S.L., R.C. Thompson, and T.S. Galloway. 2013. The physical impacts of microplastics on marine organisms: A review. *Environmental Pollution* 178: 483–492.
- Zinn, M., B. Witholt, and T. Egli. 2001. Occurrence, synthesis and medical application of bacterial polyhydroxyalkanoate. *Advanced Drug Delivery Reviews* 53: 5–21.

1
2 423 **Legend of Figures**

3
4 424
5
6
7 425 **Figure 1.** SOD (A and B) and CAT (C and D) activity in gills (A and C) and digestive gland (B and D)
8
9 426 of *Mytilus edulis* exposed to different treatment groups A-H as follows: (A) Control (no added
10
11 427 contaminants), (B) Flu only, (C) PE MPs only, (D) PHB MPs only, (E) PE MPs-Flu co-exposure, (F)
12
13 428 PHB MPs-Flu co-exposure, (G) Flu-incubated PE MPs and (H) Flu-incubated PHB MPs. Bars show
14
15
16 429 mean values $\pm$ S.D. Different letters (a, b, c) indicate statistical significant differences ($P < 0.05$, one-
17
18 430 way ANOVA with post-hoc Bonferroni test). Note different scales on different x-axes.

19
20 431
21
22
23 432 **Figure 2.** GPx (A and B) and SeGPx (C and D) activity in gills (A and C) and digestive gland (B and
24
25 433 D) of *Mytilus edulis* exposed to different treatment groups A-H as follows: (A) Control (no added
26
27 434 contaminants), (B) Flu only, (C) PE MPs only, (D) PHB MPs only, (E) PE MPs-Flu co-exposure, (F)
28
29
30 435 PHB MPs-Flu co-exposure, (G) Flu-incubated PE MPs and (H) Flu-incubated PHB MPs. Bars show
31
32 436 mean values $\pm$ S.D. Different letters (a, b, c) indicate statistical significant differences ($P < 0.05$, one-
33
34 437 way ANOVA with post-hoc Bonferroni test). Note different scales on different x-axes.

35
36 438
37
38
39
40 439 **Figure 3.** GST (A and B) and GR (C and D) activity in gills (A and C) and digestive gland (B and D)
41
42 440 of *Mytilus edulis* exposed to different treatment groups A-H as follows: (A) Control (no added
43
44 441 contaminants), (B) Flu only, (C) PE MPs only, (D) PHB MPs only, (E) PE MPs-Flu co-exposure, (F)
45
46 442 PHB MPs-Flu co-exposure, (G) Flu-incubated PE MPs and (H) Flu-incubated PHB MPs.
47
48
49 443 Bars show mean values $\pm$ S.D. Different letters (a, b, c) indicate statistical significant differences ($P <$
50
51 444 0.05 , one-way ANOVA with post-hoc Bonferroni test).

52
53 445
54
55
56
57
58
59
60

~~Declarations of interest: none.~~

Effects of combined exposures of fluoranthene and polyethylene or polyhydroxybutyrate microplastics on oxidative stress biomarkers in the blue mussel (*Mytilus edulis*)

Gabriele Magara^a, Farhan R. Khan^{b*}, Marika Pinti^a, Kristian Syberg^b, Angelo Inzirillo^a and Antonia Concetta Elia^{a*}

^a Department of Chemistry, Biology and Biotechnology, via Elce di sotto 8, University of Perugia, 06123 Perugia, Italy

^b Department of Science and Environment, Universitetsvej 1, P.O. Box 260, Roskilde University, DK-4000 Roskilde, Denmark

Running title: Bioplastics as a vector for environmental pollutants

* Corresponding author: Dr. Antonia Concetta Elia

Department of Chemistry,

Biology and Biotechnology,

University of Perugia, Italy

Email: ~~antonia.elia~~@unipg.it

Dr. Farhan R. Khan

Department of Science and Environment,

Roskilde University, Denmark

Email: frkhan@ruc.dk, farhan.khan@gmx.com

1
2 24
3
4 25
5
6 26
7
8
9 27
10
11 28
12
13 29
14
15 30
16
17 31
18
19 32
20
21 33
22
23 34
24
25 35
26
27 36
28
29 37
30
31 38
32
33 39
34
35 40
36
37 41
38
39 42
40
41 43
42
43 44
44
45 45
46
47 46
48
49 47
50
51
52
53
54
55 47
56
57
58
59
60

ABSTRACT

A growing interest in developing and commercialization of new ecofriendly plastic polymers is occurring ~~attributed~~ ~~owing~~ to the impact of marine plastics debris and microplastics that result from the degradation of oil-based polymers ~~as these substances adversely affect ecosystem health~~. Recently, polyhydroxybutyrate (PHB) has become of interest due its biodegradability and physicochemical properties. However, biological ~~consequences~~ ~~impacts~~ resulting from bioplastics exposure ~~remains to be determined~~ ~~is yet unknown~~. Moreover ~~Further~~, few data are apparently available regarding ~~nothing is known about~~ the potential for bioplastics to act as a vector for exogenous chemicals in the environment. The aim of the study was to compare the effects of polyethylene (PE MPs) and polyhydroxybutyrate (PHB MPs) microplastics administered ~~alones~~ ~~singularly~~ or in combination with fluoranthene (Flu) on detoxifying enzymes in digestive glands and gills of *Mytilus edulis*. Blue mussels were exposed for 96hr to ~~8~~ ~~eight~~ experimental groups: control, Flu-only, PE MPs-only, PHB MPs-only, PE MPs-Flu co-exposure, PHB MPs-Flu co-exposure, Flu-incubated PE MPs and Flu-incubated PHB MPs. Superoxide dismutase (~~SOD~~), catalase (~~CAT~~), glutathione peroxidases (~~GPx~~), glutathione S-transferase (~~GST~~) and glutathione reductase (~~GR~~) ~~were found~~ ~~have proved~~ to be ~~significantly~~ susceptible to Flu and plastics in both tissues. Interestingly, ~~single exposure to~~ PHB MPs ~~single exposure~~ led to ~~decreased levels of~~ CAT and GST in gills, SOD in digestive glands and SeGPx in both ~~tissues~~ ~~biomarkers~~ ~~biomarkers~~ ~~STATE INCREASE OR DECREASE?~~ ~~in SOD, CAT, etc.~~ ~~impairments~~. In co-exposure and incubation treatments, biochemical responses were generally comparable with those exerted by PE MPs or PHB MPs only, suggesting ~~an apparent absence of~~ ~~no~~ combined effects with pollutant. ~~The present findings are amongst the first to describe~~ ~~Data demonstrate~~ the ecotoxicological ~~potential~~ impact of bioplastics materials ~~on digestive glands and gills of Mytilus edulis~~.

Keywords: biodegradable polymer, biomarker profile, -bioplastics, [blue mussels](#), microplastic

Introduction

Oil-based plastic debris and subsequently the microplastics (MPs) formed by their degradation are one of the ~~current major biggest~~ environmental ~~concerns. issues of the day.~~ These plastics are defined ~~chemically~~ as organic or semi-organic materials with long chains (macromolecules) and high molecular weight, generated by the polymerization of monomers extracted from oil or gases (Cole et al., 2011). Once in seawater, microplastics ~~were have been~~ shown to be ingested by aquatic species which ~~resulted in can lead to~~ a number of deleterious effects ([Guzzetti et al., 2018](#); [Alimba and Faggio, 2019](#)) (~~NEED REF??~~). The color, density, shape and size of MPs often leads marine organisms to mistake MPs for food items and the consequences of this includes, blockage of the digestive system and false feeling of satiation (Wright et al., 2013). ~~In addition, -and also~~ MPs may act as a vehicle for ~~the~~ transport of several contaminants dissolved in ~~the aquatic~~ environment (Andrady, 2011). This ‘vector effect’ (Syberg et al., 2015), suggests that oil-based MPs may trap and transport environmental pollutants ~~including, such as~~ persistent organic pollutants (POPs) or metals (Oliveira et al., 2013; Avio et al., 2015; Khan et al., 2015; Magara et al., 2018). The possibility that plastics ~~might can~~ then transport the adhered contaminants into aquatic biota is ~~currently being still~~ debated (Koelmans et al., 2016), ~~and but~~ laboratory investigations into the vector effect remain ongoing.

There is a mounting interest in “green materials” and a focus on ~~the~~ development and production of biodegradable polymers (Mohanty et al., 2002). ~~In contrast to Unlike~~ oil-based polymers, these biopolymers ~~are derived~~ from bioprocesses, using renewable resource in bio-refineries, ~~-and d~~ Due to their mechanical and ~~physicochemical -physieal~~ characteristics, ~~these biodegradable polymers~~ may be considered as an eco-friendly substitutes to plastic (Anjum et al., 2016). In recent years,

polyhydroxybutyrate (PHB) has received ~~ing~~ growing interest for its characteristics similar to oil-based plastic ~~(NEED-REF??)~~(Dacosta et al., 2016). ~~It~~ Polyhydroxybutyrate is a water insoluble, but highly biodegradable biopolymer, relatively resistant to hydrolytic degradation ~~which~~. ~~It~~ is extracted with chloroform from bacterial cultures ~~which is~~ grown on carbohydrates. PHB (1) displays ~~has~~ low oxygen permeability and ~~reliable good~~ thermoplasticity with poor mechanical properties, (2) high crystallinity degree and (3) is optically active, with a mass about approximately 0.5×10^6 Da (Anjum et al., 2016). However, although PHB is currently available commercially, especially as main component of biodegradable market bags, to ~~the our~~ best ~~of our~~ knowledge little is known ~~regarding about it is the~~ environmental impact or effects on aquatic organisms. By sharing many characteristics with oil-based plastics, it is feasible that in the aquatic environment, bioplastic undergoes modifications, ~~resulting in~~ ~~leading to~~ generation of MPs as is the case for oil-based polymers (Andrady, 2011). ~~If due to~~ ~~environmental and societal pressures a move is made towards use of bioplastics, then it is vital to~~ ~~gather information on the influence of these bioplastics on the ecosystem health.~~

Oxidative stress results from an imbalance between pro-oxidants, such as reactive oxygen species (ROS), and the protective antioxidant system. Mechanisms that involve glutathione (GSH), and the related antioxidant ~~enzymes s~~ superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidases (SeGPx and GPx) represent important protective metabolic pathways ~~which and~~ are used as biomarkers related to pollutant induced oxidative stress. Glutathione (GSH) in its reduced form is an important non-enzymatic scavenger of oxyradicals, ~~and is~~ involved in the metabolism of toxic compounds and endogenous substances (Meister and Anderson, 1983). Both SOD and CAT enzymes catalyze the breakdown of ROS-~~generating causing~~ O_2^- and H_2O_2 , respectively and are key components within the primary defense system against oxidative stress-induced damage. SeGPx enzyme ~~can~~ reduces either hydrogen peroxide or organic peroxides and GPx enzyme affects organic hydroperoxides. ~~Polycyclic aromatic hydrocarbons (PAHs)~~, of which fluoranthene (Flu) is often used as a model contaminant ~~(and in addition a priority aquatic pollutant (Directive 2008/105/EC))~~, are

known to ~~initiate~~~~generate~~ oxidative stress which ~~was~~ ~~has been~~ assessed by these aforementioned biomarkers (Cheung et al., 2001). The oxidative stress biomarkers ~~employed~~ ~~used~~ in the present study; ~~were~~~~have been~~ utilized to ~~determine~~ ~~assess~~ the impact of a range of xenobiotics on aquatic organisms (Al kaddissi et al., 2012; 2016; Cozzari et al., 2015; Elia et al., 2017a; 2017 b; 2018; Magara et al., 2018). Previous studies utilizing these biomarkers to elucidate the MP vector effect ~~have~~ demonstrated a perturbation of intracellular antioxidant defenses (Oliveira et al., 2013; Avio et al., 2015; Magara et al., 2018).

~~Our p~~ ~~Previously~~ ~~study~~ (Magara et al., 2018) investigated the influence of polyethylene microplastic beads (PE MPs) on the accumulation and associated oxidative stress responses attributed to ~~fluoranthene~~ (Flu) in blue mussels, *Mytilus edulis*. The mussels were exposed to 4 ~~four~~ treatment groups: Flu-only, MP-only, Flu and MP co-exposure, and Flu-incubated MP. Individual contaminant exposures to Flu or MP ~~resulted~~ ~~in~~ ~~to~~ varying responses, but co-exposures and

~~WHAT IS THIS~~ incubated ~~???DO YOU MEAN IN VITRO?~~ ~~treatments~~ (i.e. Flu was sorbed to plastic surfaces prior to exposure) ~~exposures~~ did not ~~induce~~ ~~result in~~ additive or synergistic ~~responses~~ ~~effects~~.

~~Moreover~~ ~~Further~~, MP-only exposure appeared to be capable of eliciting direct effects on the oxidative stress system as ~~evidenced~~ ~~demonstrated~~ by ~~the~~ ~~enhanced~~ activities of CAT and GPx (Magara et al., 2018).

~~If due to environmental and societal concerns a move is made towards bioplastics, then it is vital to gather information on the impacts of the bioplastics. This includes the possibility of bioplastics as vectors of other pollutants.~~

Thus, the present study was designed to investigate the potential impact of PHB microplastics as a single contaminant and in combination with ~~fluoranthene~~ (Flu), ~~a representative PAH~~, on the oxidative stress system of blue mussel (*Mytilus edulis*), a well-regarded test species for MPs (Avio et al., 2015; Paul-Pont et al., 2016; Sahlmann et al 2017; Von Moos, 2012). PE MPs, as an example of oil-based MPs, were ~~employed~~ ~~as~~ ~~used~~ a comparison to ~~the~~ PHB MPs. The aim of the study was to examine the

comparative changes on oxidative stress biomarkers levels induced by oil based (PE) MPs and ‘new’ bioplastic PHB MPs as both single contaminants and as potential vectors for PAHs.

~~SHIFT TO METHODS~~Mussels were exposed for 96 h to seven treatment groups; Control (no added contaminants), Flu only, PE MPs only, PHB MPs only, PE MPs-Flu co-exposure, PHB MPs-Flu co-exposure, Flu-incubated PE MPs and Flu-incubated PHB MPs. !!

~~The aim of the study was to investigate the comparative changes on oxidative stress biomarkers levels induced by oil based (PE) MPs and ‘new’ bioplastic PHB MPs as both single contaminants and as potential vectors for PAHs.~~

Materials and methods

Mussels collection and maintenance

Blue mussel (*Mytilus edulis*) were collected in June, 2016, in Vellerup Vig – Isefjord, N 55°40.6'; E 11°48.7', in an ~~un~~ non-polluted deep inlet of Baltic Sea near Roskilde (Denmark). The mussels were ~~maintained~~ located in a temperature-controlled room (10 ± 1°C). All specimens were placed in tanks (size 39 x 21 x 25 cm), filled with 10 L of field water. Field water was gradually replaced with ~~laboratory~~ water maintaining similar salinity (20 salinity). Mussels were fed for the following three weeks twice weekly. A 16:8 h ~~l~~ light/dark photoperiod was maintained throughout.

Chemicals and preparation

~~___~~ A primary stock solution of Flu (Fluka Chemika, Steinheim, Switzerland) was prepared at 1 mg/ml ~~L~~ in acetone. PE MPs were purchased from Cospheric LLC (Santa, Barbra, CA, USA) and PHB beads ~~were~~ provided by Prodotti Gianni S.R.L. (Milano, Italy). Both polymers were white in color and similar in size (10–90 µm). Both MPs were treated with 0.1% Tween80 to enable ~~allow~~ dispersion in water column (Khan et al., 2015).

To prepare each exposure treatment ~~correct~~ accurately a weight to MP number ratio was determined for both PE and PHB particles. The weight associated to 1×10^6 MPs was weighed out for each treatment group containing MPs. The MPs were dispersed in 5 ml of 0.1% Tween80 solution and shaken at 150 rpm overnight. The MPs were filtered through 1 μ m nylon mesh and resuspended in 200 ml ~~L-of~~ 20 salinity water and kept shaking until the start of the experiment (150 rpm). MPs that required incubation with Flu were resuspended into 20 ml ~~L-of~~ acetone to which the appropriate amount of Flu stock was added to ~~yield~~ give the correct final concentration of $100 \mu\text{g}/\text{L}^{-1}$ when made up to the final exposure solution. After an initial mixing, dispersions were left under a fume hood for the acetone to evaporate and the Flu sorb to the PE MPs. This method was adopted from use with other particulate contaminants (Al-Subiai et al., 2012). ~~Having achieved-When~~ complete dryness ~~was~~ achieved overnight the Flu-incubated MPs were resuspended in 200 ml ~~L-of~~ 20 salinity water and ~~were~~ placed with the other dispersions on the shaking table overnight.

Mussels exposure

Mussels were exposed for 96 hr to ~~8 eight~~ treatment groups as follows, (A) Control (no added contaminants), (B) Flu only, (C) PE MPs only, (D) PHB MPs only, (E) PE MPs-Flu co-exposure, (F) PHB MPs-Flu co-exposure, (G) Flu-incubated PE MPs and (H) Flu-incubated PHB MPs (n=4 per treatment). ~~(1) Control (no added contaminants), (2) Flu only, (3) PE MPs only, (4) PHB MPs only, (5) PE MPs-Flu co-exposure, (6) PHB MPs-Flu co-exposure, (7) Flu-incubated PE MPs and (8) Flu-incubated PHB MPs.~~ The Flu and MP concentrations across all treatments ~~were as~~ 100 $\mu\text{g}/\text{L}^{-1}$ and 1000 $\text{MPs}/\text{mL}^{-1}$, respectively. Although the concentrations ~~exceeded~~ exceeded environmental realistic ~~conditions~~ conditions, the aim of this study was to understand mechanistic responses that are more easily captured at higher exposure ~~concentration levels~~ concentration levels. The concentrations used are consistent with previous studies (Al-Subiai et al., 2012; Cole et al., 2016; Magara et al., 2018). Acetone and ~~t~~ Tween80 were consistently added across treatments to prevent the confounding ~~influence~~ influence ~~mpaet~~ of solvent and

detergent. Ninety-six ~~hr~~ mussel exposures were conducted at $10\pm 1^{\circ}\text{C}$ and a 16/8 hr light/dark photoperiod. Mussels were individually exposed in aerated beakers (250 mL) filled with 200 mL of ~~exposure~~ ~~treatment~~ solution. ~~Five~~ Four samples of two pooled mussels were exposed per treatment group.

To start the experiment, each 200 mL dispersion was made up to 1 L with 20 salinity water and vigorously stirred to ensure ~~the~~ homogenous distribution of MPs in the water column. The 1 L dispersion was then divided ~~split~~ into ~~5~~ five 200 mL units ~~L exposures~~, again ensuring a homogenous distribution of MPs between beakers. The appropriate amount of Flu stock was added to the Flu-only and co-exposures groups prior to ~~before~~ the start of the experiment. Post exposure (96 hr), mussels were removed and rinsed thoroughly under 20 salinity water. Gill and digestive gland tissues were extracted ~~dissected~~ from each individual and placed in separate pre-weighed and pre-labelled vials. Tissues for biomarkers and Flu concentration analysis were stored at -80°C and -20°C , respectively. No mortalities were recorded during the experiments ~~posure~~.

Biochemical analysis

The analysis of biomarkers is described in detail elsewhere (Cozzari et al., 2015; Magara et al., 2018). Briefly, total GSH levels ~~glutathione, as well as activities of~~ SOD, CAT, GPx and SeGPx were measured in cytosolic fraction of gills and digestive gland ~~by~~ spectrophotometrically ~~analysis~~ (Varian Cary 50 spectrophotometer at 25°C). Protein concentrations in the cytosol were determined according to Lowry et al. (1951) and ~~were~~ used to normalize enzyme levels. All analyses were performed in triplicate along with blank samples (buffer and reagents only). These absorbance values were subtracted from those of the sample. SOD activity was determined according to an established method (McCord and Fridovich, 1969). Catalase levels were measured following the decrease in absorbance due to H_2O_2 consumption at 240 nm (Greenwald, 1985). GPx activities (SeGPx and total GPx) were determined following the oxidation of nicotinamide adenine dinucleotide phosphate reduced form

(NADPH) at 340 nm (Lawrence and Burk, 1976). GR activity was measured in 100 mM NaH₂PO₄ + Na₂HPO₄ buffer, pH 7, 1 mM oxidized glutathione (GSSG), and 60 µM NADPH; the decrease in absorbance due to the oxidation of NADPH was measured at 340 nm (Chung et al. 1991). GST activity was determined in 100 mM NaH₂PO₄ + Na₂HPO₄ buffer, pH 6.5, 1 mM GSH, 1 mM 1-chloro-2,4-dinitrobenzene (CDNB) as substrates and sample. Formation of the conjugate with GSH was read at 340 nm (Habig et al. 1974). Total glutathione (GSH+2GSSG) was measured following homogenization in 5% sulphosalicylic acid with 4mM EDTA by the GR recycling assay at 412 nm (Akerboom and Sies, 1981).

Statistical analysis

Data are reported as mean values ± the standard deviation (SD). Levene's test for normal distribution was performed on datasets prior to statistical analysis. A one-way ANOVA with Bonferroni's multiple comparison test was used to investigate differences between treatment groups ([GraphPad Prism software](#)). Statistical significance was set at P < 0.05.

Results

In the gills, CAT activity levels significantly decreased in all treatment groups (up to 80%), except for PE MPs only exposure ~~when~~ compared to control (Figure 1C). The activity of SeGPx was markedly consistently lower (up to 90%) through the treatments (Figure 2C). GST levels in the treated groups were significantly lower than control (up to 50%), ~~unless exposed to~~ except for -Flu-incubated PHB MPs (Figure 3A). No marked changes of SOD, GPx and GR activities ~~y~~ were ~~noted~~ recorded for all ~~the~~ treatment compared to ~~respective~~ the own controls (Figures 1A, 2A, 3C).

In the digestive glands, SOD ~~activity concentrations~~ were significantly reduced consistently ~~lowered~~ (up to 50%) in all treatments when compared to control with the exception of the group incubated with ~~exposed~~ to PE MPs only (Figure 1B). CAT activity was markedly decreased ~~lowered~~

similar~~ly~~ to SOD (up to 60%), ~~but no statistically~~ However, no significant differences were ~~found~~ recorded in mussels exposed to PHB MPs only and Flu-incubated PHB MPs (Figure 1D). Total GPx markedly ~~fell dropped~~ (60%) following Flu exposure (Figure 2B). Levels of SeGPx ~~activity~~ were ~~reliably lower significantly reduced compared to than~~ control (up to 50%), ~~except unless~~ for PE MPs only and Flu-incubated PHB MPs groups (Figure 2D). GST activity peaked (1.5 fold ~~higher~~) in Flu exposed group (Figure 3B), whereas GR ~~activity~~ levels showed increasing trends in all treated groups. ~~‡~~The GR activity was markedly higher following PE MPs only exposure (1 fold), PE MPs-Flu co-exposure (1.2 fold) and Flu-incubated PHB MPs treatment (1 fold, Figure 3D).

Discussion

The present study provides the first experimental evidences of ~~the~~ stress-related effects initiated~~caused~~ by novel bio-microplastic exposure, either as a single contaminant or in combination with Flu, ~~in to~~ the blue mussel *Mytilus edulis*. The comparison was ~~conductedmade~~ with the more traditional oil-based plastic polyethylene. Flu and both PE and PHB MPs modified the antioxidant responses in both gills and digestive glands. The present results are largely in ~~agreement line~~ with those previously reported for Flu and PE MPs, both in single or combined exposures (Magara et al., 2018).

PE MPs altered the levels of several of the tested biomarkers. ~~Although previous studies showed that oil-based plastic led to enhanced antioxidant activities in copepods (Jeong et al., 2017) and fish (Barboza et al., 2018), our results are in agreement with studies on mussels~~ This outcome is in ~~agreement line with previous studies~~, showing a significant inhibition of CAT, SeGPx and GST activity following oil-based MPs mussel exposure (Avio et al., 2015; Paul-Pont et al., 2016). ~~A D~~ decrease of CAT and GPx ~~activities~~ in digestive glands and of SeGPx and GST in gills exposed to PE MPs may be related to the size and shape of microplastics, that ~~can~~ play a key role ~~in initiation of on-promote~~ biological changes (Browne et al., 2008; Von Moos, 2012; Avio et al., 2015). Small polystyrene plastic particles ~~may can~~ be translocated from the gut to the circulatory system of *Mytilus edulis* and

~~subsequently be~~ retained, suggesting that ~~these substances~~ y ~~might can~~ reach crucial organs such as the heart or ~~the~~ hepatic tissues ~~to produce adverse effects and then hurt~~ (Browne et al., 2008). It is well known that oil-based microplastics may accumulate within organisms and ~~induce cause~~ tissue abrasions, as ~~evidenced~~ by histological changes in digestive cells and triggered inflammatory responses, formation of granulocytomas and lysosomal destabilization (Avio et al., 2015; Von Moos, 2012). Therefore, in this scenario, the imbalance of antioxidant defense markers ~~might be expected is not surprising~~ and ~~can might perhaps~~ be attributed to ~~the~~ physical damage induced by MPs ~~enabling further accumulation of reactive products. -leading to a slight accumulation of reactive products.~~ Furthermore, in the present study ~~the~~ increased GR activity in digestive glands ~~might reflect the can~~ ~~sustenance the rising~~ demand ~~for of~~ GSH as defense mechanism against oxidative stress ~~induced caused~~ by oil-based MPs (Jeong et al., 2016).

~~—THIS IS RESULTS! DELETE~~Results from this study described, for the first time, that exposure to bioplastic PHB MPs also showed impairment of antioxidant biomarkers. Activity of SOD decreased only in digestive glands, CAT in gills and Se-GPx in both tissues.

~~NO EVIDENCE FOR INFLAMMATION DELETE~~As for PE MPs, we may hypothesize a mechanic hurt, following inflammatory response. However, because of the biodegradability potential of PHB, chemical damages cannot be excluded. Indeed, it is feasible that potential toxic compounds have been generated during biopolymer degradation process. At our best knowledge, no studies have been carried out on the metabolism of PHB in marine mussels.~~METABOLISM NOT MEASURED~~ Though, it is known that the PHB degradation rate is in the order of few months in anaerobic environment, and years in seawater (Madison and Huisman, 1999; Verlinden et al., 2007). Within the organisms, PHB has a degradation rate, albeit low, due to its physical-chemical characteristics, such as the high crystallinity (Zinn et al., 2001; Anjum et al., 2016). Therefore, we may suppose that during the 96h trial, *Mytilus edulis* has start to degraded PHB MPs and a higher hydroxybutyrate tissue concentration occurred,

leading to a triggered ROS production in combination with physical damage. ~~YOU DID NOT MEASURE THIS DELETE~~

The present study also carried out a co-exposure and incubation scenario of both PE and PHB MPs with Flu and ~~reported the results have provided~~ information on oxidative stress related to the simultaneous exposure to compounds. In this scenario, a mussel experimental group exposed to Flu-only was used as positive control. ~~Previously Pan et al (2005) found that at high benzo(k)fluoranthene, a PAH, concentrations the activity of antioxidant enzymes WHICH STATE?? was diminished in the scallop *Chlamys farreri* in a~~ ~~The time--dependent manner--course tendency of antioxidant enzymes to restrain activities at high benzo(k)fluoranthene concentrations was already reported on scallop *Chlamys farreri* (Pan et al., 2005). Accordingly, Our findings that~~ the marked decrease of SOD ~~activity~~ levels and both GPx²s in digestive glands, ~~as well as of~~ SeGPx ~~also~~ in gills and ~~CAT catalase~~ activity in both tissues ~~indicate suggests~~ that this ~~Flu PAH~~ may exert detrimental effects when ~~singularly~~ administered ~~alone resulting in, causing~~ a severe oxidative stress ~~o~~in *Mytilus edulis*. The impairment of antioxidant biomarkers levels may be attributed to a chemical-~~mediated~~ damage. Indeed, the ~~enhanced triggered~~ GST activity measured in digestive glands ~~might~~ represent an important outcome, since ~~this --may~~ indicate ~~elevated a strengthening~~ of phase-II biotransformation metabolism. The significant induction of GST activity in mussels exposed to Flu-only ~~may~~ ~~can~~ be ~~attributed explained due to the~~ production of specific substrates for GST by phase I enzymes, representing an effective defense line against ~~Flu. fluoranthene. Babson et al (1986) showed that rat liver microsomes converted Flu to trans-2,3-dihydrodiol as the major metabolite. Metabolism study on rat liver microsomes showed that Flu is converted to trans 2,3-dihydrodiol as the major metabolite (Babson et al., 1986).~~ Although data ~~regarding about~~ Flu metabolites in blue mussels ~~are is~~ lacking, ~~we can it is~~ ~~suggested~~ that GST induction may be related to ~~the~~ production of this intermediate.

However, ~~b~~Biomarkers responses in digestive glands and gills of mussels exposed to both Flu and PE or PHB MPs (co-exposure and incubated treatment) generally followed a trend similar to ~~the~~ PAH ~~alone own traditional and or the~~ new plastic ~~alone~~ and no combined effects ~~of these agents was~~ apparent. The absence of combined effects in *Mytilus edulis* exposed to PE MPs and Flu is in agreement ~~consistent~~ with ~~the results of our previous study~~ our previous results (Magara et al., 2018). ~~This particular outcome may could be related to a MAKES NO SENSE retention mechanism?? of Flu exerted by PE MPs, due to MAKES NO SENSE specific physical properties, suggesting that MPs may act as a “sink” of environmental pollutants (Chua et al. 2014; Khan et al. 2015).~~ Chua et al. (2014), showed that the concentration of ~~P~~polybrominated ~~D~~diphenyl ~~E~~ethers (PBDE) was lower in *Allorchestes compressa* co-exposed to microplastics than specimens treated with ~~single~~ PBDE ~~alone~~. ~~The authors Chua et al (2014)~~ concluded that the presence of MPs may inhibit the uptake of PBDE, perhaps because this contaminant is strongly absorbed onto microplastics surface, making it less bioavailable. ~~(Chua et al., 2014).~~ It is already known that oil-based micropolymers have the propensity to aggregate in water ~~(Alimi et al., 2018) (NEED REF??).~~ ~~A p~~Previously ~~study~~ Khan et al (2015) found ~~o~~in zebrafish ~~showed~~ a decrease in silver (Ag) uptake by fish ~~attributed to , probably due to diminished a minimization of the~~ contacts between ~~the~~ Ag and tissues following oil-based MP aggregation. ~~(Khan et al., 2015).~~ Furthermore, Magara et al. (2018) reported that Flu tissues concentrations ~~were as~~ lower in *Mytilus edulis* exposed to both co-exposure ~~with and incubated~~ polyethylene and PAH compared to specimens treated with ~~Flu alone. fluoranthene-only~~. The absence of combined effects ~~of PE MPs and Flu in Mytilus edulis-exposed to PE MPs, either in co-exposure or incubated treatment,~~ and previous results on Flu uptake in ~~B~~blue mussel (Magara et al., 2018), suggests that the interaction of tissues with Flu ~~might could~~ be delayed by this aggregation mechanism exerted by microplastics. ~~THIS WAS NOT MEASURED, DELETE Concerning the biodegradable PHB polymer, biomarkers responses surprisingly allow to suppose characteristics similar to polyethylene, although knowledge on~~

~~PHB retention or aggregation properties are not available. However, due to its ability to degrade in tissues, further studies are needed to gather knowledge on the pro-oxidant role of bio-microplastic.~~

~~The results of the present research show for the first time a role of bioplastics in affecting antioxidant and THIS WAS NOT DONE detoxifying pathways.~~

Bioplastics are currently becoming the leading material for replacing oil-based polymers since it is more desirable than traditional plastic due to ~~a its~~ propensity to biodegrade in ~~the~~ environment (Anjum et al., 2016). ~~A r~~Recently Napper and Thompson (2019) study demonstrated that biodegradable polymers ~~may could~~ not undergo any substantial deterioration over a 3 year period in marine environment, but may be reduced in small fragments, ~~as is~~ similar ~~to for~~ oil-based plastics ~~(Napper and Thompson, 2019)~~. Our data suggest that PHB MPs ~~results in may lead to~~ altered levels of some oxidative stress biomarkers, ~~again which is~~ similar to oil-based MPs. The fact that bio-microplastics may exert stress effects and the potential low biodegradability rate ~~may~~ indicate a novel threat for the marine environment. Therefore, it is crucial to understand ~~the~~ degradation processes of bioplastics in the environment, ~~but whilst~~ also gathering knowledge regarding about their potential ecotoxicological impacts. ~~Whilst~~ Although the damage exerted by oil-based polymers is becoming increasingly understood, it is ~~still now~~ necessary to investigate the possible consequences impacts of bioplastics exposure, both alone and in combination with other environmental pollutants present in the aquatic ecosystem.

Conclusions

The results of the present study demonstrated suggest that PHB MPs ~~may~~ modify the baseline levels of biomarkers related to oxidative stress in *Mytilus edulis*. ~~These levels of~~ alterations were similar to those exerted by PE MPs treatments for ~~somefew~~ of the antioxidant detoxifying biomarkers. Furthermore, the responses measured in both co-treatments were similar to those noted with of MP-aloneonly exposures. Data indicated This suggests the absence of any no combined effects produced

caused by oil-based MPs or bioplastics MPs and Flu. fluoranthene. ~~Further studies are required to gather information on the potential consequences of new and innovative bioplastic polymer in the environment.~~

Acknowledgements

The visit of A. Inzirillo to Department of Science and Environment, Roskilde University, was funded by a LLP/Erasmus Placement program (2015/2016). The Authors thank Prodotti Gianni S.R.L. (Milano, Italy) company kindly providing the PHB beads.

References

- Al Kaddissi, S., O. Simon, A.C. Elia, P. Gonzalez, M. Floriani, I. Cavalie, V. Camilleri, S. Frelon, and A. Legeay. 2016. How toxic is the depleted uranium to crayfish *Procambarus clarkii* compared with cadmium? Environmental Toxicology 31:211–223.
- Al Kaddissi S., S. Frelon, A.C. Elia, A. Legeay, P. Gonzalez, F. Coppin, D. Orjollet, V. Camilleri, K. Beaugelin-Seiller, R. Gilbin, and O. Simon. 2012. Are antioxidant and transcriptional responses useful for discriminating between chemo- and radiotoxicity of uranium in the crayfish *Procambarus clarkii*? Ecotoxicology and Environmental Safety 80: 266-272.
- Alimba, C., and C. Faggio. 2019. Microplastics in the marine environment: current trends in environmental pollution and mechanisms of toxicological profile. Science of the Total Environment 68: 61-74.

- Alimi, O. S., J. Farner Budarz, J. L. M., Hernandez, L. M., and N. Tufenkji, N. 2018. Microplastics and nanoplastics in aquatic environments: aggregation, deposition, and enhanced contaminant transport. *Environmental Science and Technology* 52: 1704-1724.
- Al-Subiai, S. N., V. M. Arlt, P. E. Frickers, J. W. Readman, B. Stolpe, J. R. Lead, A. J. Moody, and A. N. Jha. 2012. Merging nano-genotoxicology with eco-genotoxicology: An integrated approach to determine interactive genotoxic and sub-lethal toxic effects of C 60 fullerenes and fluoranthene in marine mussels, *Mytilus* sp. *Mutation Researcher Genetic Toxicology and Environment Mutagenesis* 745: 92-103.
- Akerboom, T.P.M., and H. Sies. 1981. Assay of glutathione disulfide and glutathione mixed disulfide in biological samples. *Methods in Enzymology* 71: 373-382.
- Andrady, A.L. 2011. Microplastics in the marine environment. *Marine Pollution Bulletin* 62(8): 1596-1605.
- Anjum, A., M. Zuber, K.M. Zia, A. Noreen, M.N. Anjum, and S. Tabasum. 2016. International Journal of Biological Macromolecules Microbial production of polyhydroxyalkanoates (PHAs) and its copolymers: A review of recent advancements. *International Journal of Biological Macromolecules* 89: 161-174.
- Avio, C.G., S. Gorbi, M. Milan, M. Benedetti, D. Fattorini, G. d'Errico, M. Pauletto, L. Bargelloni, and F. Regoli. 2015. Pollutants bioavailability and toxicological risk from microplastics to marine mussels. *Environmental Pollution* 198: 211-222.
- Babson, J.R., S.E. Russo-Rodriguez, R.V. Wattley, P.L. Bergstein, W.H. Rastetter, H.L. Liber, B.M. Andon, W.G. Thilly, and G.N. Wogan. 1986. Microsomal activation of fluoranthene to mutagenic metabolites. *Toxicology and Applied Pharmacology* 85(3): 355-366.
- Barboza, L.G.A., L. R. Vieira, V. Branco, C. Carvalho, and L. Guilhermino. 2018. Microplastics increase mercury bioconcentration in gills and bioaccumulation in the liver, and cause oxidative

- [stress and damage in *Dicentrarchus labrax* juveniles. Scientific Reports 8: 15655](#)
- Browne, M.A.D., T.S. Galloway, D.M. Lowe, and R.C. Thompson. 2008. Ingested microscopic plastic translocates to the circulatory system of the mussel, *Mytilus edulis* (L). [_____Environmental Science and Technology 42\(13\):5026–5031.](#)
- Cheung, C.C.C., G.J. Zheng, A.M.Y. Li, B.J. Richardson, and P.K.S. Lam. 2001. Relationships between tissue concentrations of polycyclic aromatic hydrocarbons and antioxidative responses of marine mussels, *Perna viridis*. [_____Aquatic Toxicology 52\(3–4\):189–203.](#)
- Chua, E.M., J. Shimeta, D. Nugegoda, P.D. Morrison, and B.O. Clarke. 2014. Assimilation of polybrominated diphenyl ethers from microplastics by the marine amphipod, *Allorchestes compressa*. [_____Environmental Science and Technology 48\(14\):8127–8134.](#)
- Chung, P.M., R.E. Cappel, and H.F. Gilbert. 1991. Inhibition of glutathione disulfide reductase by glutathione. [_____Archives of Biochemistry and Biophysics 288:48–53.](#)
- Cole, M., P. Lindeque, C. Halsband, and T.S. Galloway. 2011. Microplastics as contaminants in the marine environment: A review. [_____Marine Pollution Bulletin 62\(12\):2588–2597.](#)
- Cole, M., P.K. Lindeque, E. Fileman, J. Clark, C. Lewis, C. Halsband, and T.S. Galloway. 2016. Microplastics alter the properties and sinking rates of zooplankton faecal pellets. [_____Environmental Science and Technology 50\(6\):3239–3246.](#)
- Cozzari, M., A.C. Elia, N. Pacini, B.D. Smith, D. Boyle, P.S. Rainbow, and F.R. Khan. 2015. Bioaccumulation and oxidative stress responses measured in the estuarine ragworm (*Nereis diversicolor*) exposed to dissolved, nano- and bulk-sized silver. [_____Environmental Pollution 198:32–40.](#)
- [Dacosta, C.F., J.A. Posada, and A. Ramirez. 2016. Techno-economic and carbon footprint assessment of methyl crotonate and methyl acrylate production from wastewater-based polyhydroxybutyrate \(PHB\). Journal of Cleaner Production 137: 942–952.](#)
- [Directive 2008/105/EC of the European Parliament and of the Council. Official Journal of the](#)

European Union.

Elia, A.C., G. Magara, C. Caruso, L. Masoero, M. Prearo, P. Arsieni, B. Caldaroni, M. Scoparo, A.J.M. Dörr, S. Salvati, P. Brizio, S. Squadrone, and M.C. Abete. 2018. A comparative study on subacute toxicity of arsenic trioxide and dimethylarsinic acid on antioxidants and antioxidant-related enzymes in ~~e~~Crandell ~~r~~Rees feline kidney (CRFK), human hepatocellular carcinoma (PCL/PRF/5) and epithelioma ~~p~~Papulosum cyprini (EPC) cell lines. Journal of Toxicology and Environmental Health A 81: ~~(10)~~: 333-348.

Elia, A. C., G. Magara, M. Righetti, A. J. M. Dörr, T. Scanzio, N. Pacini, M. C. Abete, and M. Prearo. 2017a. Oxidative stress and related biomarkers in cupric and cuprous chloride-treated rainbow trout. Environmental Science and Pollution Research 24:10205–19.

Elia, A.C., F. Giorda, N. Pacini, A.J.M. Dörr, T. Scanzio, and M. Prearo. 2017b. Subacute toxicity effects of deltamethrin on oxidative stress markers in rainbow trout. Journal of Aquatic Animal Health 29:165-172.

Greenwald, R.A. 1985. Handbook of mMethods for oOxygen rRadicals rResearch. CRC Press, Boca Raton, FL. Ed. by R. A. Greenwald, pp 447. NEED EDITOR NAME; NEED PAGES??

Guzzetti, E., A. Sureda, S.Tejada, and C. Faggio. 2018. Microplastic in marine organism: environmental and toxicological effects. Environmental Toxicology and Pharmacology 64: 164-171.

Habig, W.H., M.J. Pabst, and W.B. Jakoby. 1974. Glutathione S transferases. The first enzymatic step in mercapturic acid formation. Journal of Biological Chemistry 249: 7130–7139.

Jeong, C.B., H.M. Kang, M.C. Lee, D.H. Kim, J. Han, D.S. Hwang, S. Souissi, S.J. Lee, K.H. Shin, H.G. Park, and J.S. Lee. 2017. Adverse effects of microplastics and oxidative stress-induced MAPK/Nrf2 pathway-mediated defense mechanisms in the marine copepod Paracyclopsina nana”. Scientific Reports 7: 41323.

- Jeong, C., E. Won, H. Kang, M. Lee, D. Hwang, and U. Hwang. 2016. Microplastic size-dependent toxicity, oxidative stress induction, and p-JNK and p-p38 activation in the monogonont rotifer (*Brachionus koreanus*). *Environmental Science and Technology* 50(16): 8849-8857.
- Khan, F.R., K. Syberg, Y. Shashoua, and N.R. Bury. 2015. Influence of polyethylene microplastic beads on the uptake and localization of silver in zebra fish (*Danio rerio*). *Environmental Pollution* 206: 73–79.
- Koelmans, A. A., A. Bakir, G. A. Burton, and C.R. Janssen. 2016. Microplastic as a vector for chemicals in the aquatic environment: Critical review and model-supported reinterpretation of empirical studies. *Environmental Science and Technology* 50: 3315–3326.
- Lawrence, R.A., and R.F. Burk. 1976. Glutathione peroxidase activity in selenium-deficient rat liver. *Biochemical and Biophysical Research Communications* 71: 592–598.
- Lowry, O.H., N.J. Rosebrough, A.L. Farr, and R.J. Randall 1951. Protein measurement with the Folin phenol reagent. *Journal of Biological Chemistry* 193: 265–275.
- ~~YOU NEED TO CHECK THIS!! Lowry, O.H., Rosebrough,?? Farr and R.J. Randall, and A. Lewis. 1951. The folin by oliver. Readings, *Journal of Biological Chemistry* 193: (1), 265–275.~~
- Madison, L.L., and G.W. Huisman. 1999. Metabolic engineering of poly (3-hydroxyalkanoates): ~~f~~From ~~DNA~~ to plastic. *Microbiology and Molecular Biology Review* 63(1): 21–53.
- Magara, G., A.C. Elia, K. Syberg, and F.R. Khan. 2018. Single contaminant and combined exposures of polyethylene microplastics and fluoranthene: accumulation and oxidative stress response in the blue mussel, *Mytilus edulis*. *Journal of Toxicology and Environmental Health –Part A* 81(16): 761–773.

1
2 463 McCord, J.M., and I. Fridovich. 1969. Superoxide dismutase: an enzymatic function for erythrocuprein
3
4 454 (hemocuprein). *Journal of Biological Chemistry* 244:6049–6055.
5
6
7 465 Meister, A., and M.E. Anderson. 1983. Glutathione. *Annual Review of Biochemistry* 52:711–60.
8
9 466 Mohanty, A.K., M. Misra, and L.T. Drzal. 2002. Sustainable bio-composites from renewable resources:
10
11 457 Opportunities and challenges in the green materials world. _____ *Polish Journal of*
12
13 458 *Environmental Studies* 10(1-2): 19–26.
14
15
16 469 Napper, I.E., and R.C. Thompson. 2019. Environmental deterioration of biodegradable, oxo-
17
18 470 biodegradable, compostable, and conventional plastic carrier bags in the sea, soil, and open-air
19
20 471 over a 3-year period. _____ *Environmental Science and Technology* 53(9): 4775-4783.
21
22
23 472
24
25 473
26
27 474 Oliveira, M., A. Ribeiro, K. Hylland, and L. Guilhermino. 2013. Single and combined effects of
28
29 475 microplastics and pyrene on juveniles (0+ group) of the common goby *Pomatoschistus microps*
30
31 476 (Teleostei, Gobiidae). _____ *Ecological Indicators* 34: 641–647.
32
33
34 477 Pan, L., J. Ren, and J. Liu. 2005. Effects of benzo(k)fluoranthene exposure on the biomarkers of
35
36 478 scallop *Chlamys farreri*. _____ *Comparative Biochemistry and Physiology C* 141(3):248–256.
37
38
39 479 Paul-Pont, I., C. Lacroix, C. Gonzalez Fernandez, H. Helene, C, Lambert, N. Le Goic, L. Frère, A.L.
40
41 480 Cassone, R. Sussarellu, C. Fabioux, J. Guyomarch, M. Albentosa, A. Huvet, and P. Soudant.
42
43 481 2016. Exposure of marine mussels *Mytilus* spp. to polystyrene microplastics: Toxicity and
44
45 482 influence on fluoranthene bioaccumulation. _____ *Environmental Pollution* 216:724–737.
46
47
48 483 Sahlmann, A., R. Wolf, T. F. Holth, J. Titelman, and K. Hylland. 2017. Baseline and oxidative DNA
49
50 484 damage in marine invertebrates. *Journal of Toxicology and Environmental Health A* 80: 807-819.
51
52
53 485 Sahlmann, A. 2017. *Journal of Toxicology and Environmental Health A* 80:
54
55 486 807-819.
56
57 487 Syberg, K., F. R. Khan, H. Selck, A. Palmqvist, G.T. Banta, J. Daley, L. Sano, and M.B. Duhaime.
58
59
60

2015. Microplastics: Addressing ecological risk through lessons learned. *Environmental Toxicology and Chemistry SETAC* 34: 945–953.
- Verlinden, R.A.J., D.J. Hill, M.A. Kenward, C.D. Williams, and I. Radecka. 2007. Bacterial synthesis of biodegradable polyhydroxyalkanoates. *Journal of Applied Microbiology* 102: 1437–1449.
- Von Moos, N.B.P. 2012. Uptake and effects of microplastics on cells and tissue of the blue mussel *Mytilus edulis* L. after an experimental exposure. *Environmental Science and Technology* 46: (20): 11327–11335.
- Wright, S.L., R.C. Thompson, and T.S. Galloway. 2013. The physical impacts of microplastics on marine organisms: a review. *Environmental Pollution* 178: 483–492.
- Zinn, M., B. Witholt, and T. Egli. 2001. Occurrence, synthesis and medical application of bacterial polyhydroxyalkanoate. *Advanced Drug Delivery Reviews* 53(4): 5–21.

Legend of Figures

Figure 1. SOD (A and B) and CAT (C and D) activity in gills (A and C) and digestive gland (B and D) of *Mytilus edulis* exposed to different treatment groups A-H as follows: (A) Control (no added contaminants), (B) Flu only, (C) PE MPs only, (D) PHB MPs only, (E) PE MPs-Flu co-exposure, (F) PHB MPs-Flu co-exposure, (G) Flu-incubated PE MPs and (H) Flu-incubated PHB MPs. Bars show mean values $\pm$ S.D. Different letters (a, b, c) indicate statistical significant differences ($P < 0.05$, one-way ANOVA with post-hoc Bonferroni test). Note different scales on different x-axes.

Figure 2. GPx (A and B) and SeGPx (C and D) activity in gills (A and C) and digestive gland (B and D) of *Mytilus edulis* exposed to different treatment groups A-H as follows: (A) Control (no added contaminants), (B) Flu only, (C) PE MPs only, (D) PHB MPs only, (E) PE MPs-Flu co-exposure, (F)

1
2 513 PHB MPs-Flu co-exposure, (G) Flu-incubated PE MPs and (H) Flu-incubated PHB MPs. Bars show
3
4 514 mean values $\pm$ S.D. Different letters (a, b, c) indicate statistical significant differences ($P < 0.05$, one-
5
6 515 way ANOVA with post-hoc Bonferroni test). Note different scales on different x-axes.
7
8
9 516
10
11
12 517 **Figure 3.** GST (A and B) and GR (C and D) activity in gills (A and C) and digestive gland (B and D)
13
14 518 of *Mytilus edulis* exposed to different treatment groups A-H as follows: (A) Control (no added
15
16 519 contaminants), (B) Flu only, (C) PE MPs only, (D) PHB MPs only, (E) PE MPs-Flu co-exposure, (F)
17
18
19 520 PHB MPs-Flu co-exposure, (G) Flu-incubated PE MPs and (H) Flu-incubated PHB MPs.
20
21 521 Bars show mean values $\pm$ S.D. Different letters (a, b, c) indicate statistical significant differences ($P <$
22
23 522 0.05, one-way ANOVA with post-hoc Bonferroni test).
24
25
26 523 ~~**Figure 1.** SOD (A and B) and CAT (C and D) activity in gills (A and C) and digestive gland (B and D)~~
27
28 524 ~~of *Mytilus edulis*. Bars show mean values $\pm$ S.D. Different letters (a, b, c) indicate statistical significant~~
29
30 525 ~~differences ($P < 0.05$, one-way ANOVA with post-hoc Bonferroni test). Note different scales on~~
31
32 526 ~~different x-axes.~~
33
34
35 527
36
37 528 ~~**Figure 2.** GPx (A and B) and SeGPx (C and D) activity in gills (A and C) and digestive gland (B and~~
38
39 529 ~~D) of *Mytilus edulis*. Bars show mean values $\pm$ S.D. Different letters (a, b, c) indicate statistical~~
40
41 530 ~~significant differences ($P < 0.05$, one-way ANOVA with post-hoc Bonferroni test). Note different~~
42
43 531 ~~scales on different x-axes.~~
44
45
46 532
47
48
49 533 ~~**Figure 3.** GST (A and B) and GR (C and D) activity in gills (A and C) and digestive gland (B and D)~~
50
51 534 ~~of *Mytilus edulis*. Bars show mean values $\pm$ S.D. Different letters (a, b, c) indicate statistical significant~~
52
53 535 ~~differences ($P < 0.05$, one-way ANOVA with post-hoc Bonferroni test).~~
54
55
56
57
58
59
60

Figure 1

189x207mm (300 x 300 DPI)

Figure 2

199x217mm (300 x 300 DPI)

Figure 3

199x216mm (300 x 300 DPI)