

Nonionic surfactants modulate the transport activity of ATP-binding cassette (ABC) transporters and solute carriers (SLC)

Relevance to oral drug absorption

Al-Ali, A A A; Nielsen, R B; Steffansen, B; Holm, René; Nielsen, Carsten Uhd

Published in:
International Journal of Pharmaceutics

DOI:
[10.1016/j.ijpharm.2019.05.033](https://doi.org/10.1016/j.ijpharm.2019.05.033)

Publication date:
2019

Document Version
Peer reviewed version

Citation for published version (APA):
Al-Ali, A. A. A., Nielsen, R. B., Steffansen, B., Holm, R., & Nielsen, C. U. (2019). Nonionic surfactants modulate the transport activity of ATP-binding cassette (ABC) transporters and solute carriers (SLC): Relevance to oral drug absorption. *International Journal of Pharmaceutics*, 566, 410-433.
<https://doi.org/10.1016/j.ijpharm.2019.05.033>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Manuscript

Review

Nonionic surfactants modulate the transport activity of ATP-binding cassette (ABC) transporters and solute carriers (SLC): Relevance to oral drug absorption

Ahmed A. Abdulhussein Al-Ali, Rasmus Blaaholm Nielsen, Bente Steffansen, René Holm, Carsten Uhd Nielsen

PII: S0378-5173(19)30385-0
DOI: <https://doi.org/10.1016/j.ijpharm.2019.05.033>
Reference: IJP 18363

To appear in: *International Journal of Pharmaceutics*

Received Date: 23 March 2019
Revised Date: 10 May 2019
Accepted Date: 11 May 2019

Please cite this article as: A.A. Abdulhussein Al-Ali, R. Blaaholm Nielsen, B. Steffansen, R. Holm, C. Uhd Nielsen, Nonionic surfactants modulate the transport activity of ATP-binding cassette (ABC) transporters and solute carriers (SLC): Relevance to oral drug absorption, *International Journal of Pharmaceutics* (2019), doi: <https://doi.org/10.1016/j.ijpharm.2019.05.033>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**Nonionic surfactants modulate the transport activity of ATP-binding cassette (ABC) transporters
and solute carriers (SLC): Relevance to oral drug absorption**

Running title: Nonionic surfactants modulate membrane transport proteins

Ahmed A. Abdulhussein Al-Ali^a, Rasmus Blaaholm Nielsen^a, Bente Steffansen^{a, 1}, René Holm^{b, c} and
Carsten Uhd Nielsen^{a*}

^a: Department of Physics, Chemistry and Pharmacy, University of Southern Denmark, Campusvej 55,
DK-5230 Odense M, Denmark.

^b: Drug Product Development, Janssen R&D, Johnson & Johnson, Turnhoutseweg 30, 2340 Beerse,
Belgium

^c: Department of Science and Environment, Roskilde University, 4000 Roskilde, Denmark

*: Corresponding author at: Department of Physics, Chemistry and Pharmacy, University of Southern
Denmark, Campusvej 55, DK-5230 Odense M, Denmark. Phone: +45 6550 9427 e-mail: cun@sdu.dk
(Carsten Uhd Nielsen)

¹: Present address: Leo Pharma, Industriparken 55, 2750 Ballerup, Denmark

Abstract:

Recently, it has become evident that pharmaceutical excipients may interfere with the activity of ATP-binding cassette (ABC) transporters and solute carriers (SLC). The present review aims to provide an overview of surfactants shown to modulate substrate transport via SLCs and ABCs, and to discuss the relevance for oral drug absorption. *In vitro*, more than hundred surfactants have been suggested to decrease the efflux activity of P-glycoprotein (P-gp, ABCB1), and many of these surfactants also inhibit the breast cancer resistance protein (BCRP, ABCG2), while conflicting results have been reported for multidrug resistance-associated protein 2 (MRP2, ABCC2). In animals, surfactants such as pluronic® P85 and polysorbate 20 have been shown to enhance the oral absorption of P-gp and BCRP substrates. Many surfactants, including cremophor® EL and Solutol® HS 15 inhibiting ABC transporters, were also found to inhibit SLCs in cell cultures. These carriers were SLC16A1, SLC21A3, SLC21A9, SLC15A1-2, and SLC22A1-3. This overlap in specificity of surfactants that inhibit both transporters and carriers might influence the oral absorption of various drug substances, nutrients, and vitamins. Such biopharmaceutical elements may be relevant for future drug formulation design.

Key words: Nonionic surfactant, Co-surfactant, ATP-binding cassette transporters, Solute carriers, Oral absorption, Lipid-based formulations.

1. Introduction:

Today, many drug substances approved for oral and parenteral use are prepared in surfactant containing formulations such as lipid-based formulations, suspensions and solid dosage forms (Savla et al., 2017). In these formulations, nonionic surfactants are used as solubilizers, stabilizers, wetting agents etc. In the ~~last~~recent years, it has become evident that some surfactants may affect the function of ~~biological~~ membrane transport proteins by altering drug substance uptake (Engel et al., 2012; Rege et al., 2002) and/or cellular efflux (Batrakova et al., 2003a; Rege et al., 2002; Yamagata et al., 2007b). Membrane transport proteins relevant to drug transport are from two major families, i.e. the ATP-binding cassette (ABC) family of efflux transporters and the solute carrier (SLC) family of cellular influx and efflux carriers. In brief, members of the ABC family (hereafter termed “transporter”) depend directly on the use of cellular ATP to complete their transport cycle, which in mammals result in cellular efflux. Members of the SLC family do not directly depend on using cellular ATP, but are driven by substrate concentrations and in many cases the concentration gradient of other substrates such as ions. A SLC facilitates either cellular influx or efflux, and are hereafter termed “carrier”. Currently, the ABC and SLC families consist of 51 and 417 members (HUGO Gene Nomenclature Committee, 2019), respectively. The first indication that pharmaceutical excipients could alter the transport function of transporters and carriers came from observations that nonionic surfactants such as cremophor® EL (Woodcock et al., 1990), Solutol® HS 15 (Coon et al., 1991), and polysorbate 80 (Woodcock et al., 1992) reversed multidrug resistance in cancer cells. In cell cultures, nonionic surfactants inhibited different members of the ABC family such as P-glycoprotein (P-gp, MDR1, ABCB1) (Lo, 2003; Rege et al., 2002), breast cancer resistance protein (BCRP, ABCG2) (Yamagata et al., 2007a, b), and multidrug resistance-associated protein 2 (MRP2, ABCC2), although conflicting results have been

observed for MRP2 (Hanke et al., 2010; Li et al., 2013a, 2014). *In vivo*, in wild type animals, nonionic surfactants have been shown to enhance the intestinal absorption and bioavailability of P-gp substrates such as digoxin (Cornaire et al., 2004; Nielsen et al., 2016; Zhang et al., 2003), etoposide (Akhtar et al., 2017; Al-Ali et al., 2018a), and paclitaxel (Varma and Panchagnula, 2005), and the BCRP substrate topotecan (Yamagata et al., 2007b). Interestingly, corresponding control experiments in transporter deficient animals showed that co-administration of nonionic surfactants with digoxin (Nielsen et al., 2016) or etoposide (Al-Ali et al., 2018a) in *mdr1a* deficient rats, or topotecan (Yamagata et al., 2007b) in *abcg2* deficient mice did not alter the oral absorption and bioavailability of these substrates. This indeed indicates that surfactants increase intestinal absorption through P-gp or BCRP inhibition, and for the drug substances in question, not through unspecific effects related to solubilizing of the drug substances or through permeation enhancing effects.

Recently, *in vitro* studies have also shown that nonionic surfactants such as polysorbate 20 and cremophor® EL inhibit the transport via several SLCs expressed ~~ien~~ on the apical membrane of enterocytes such as the organic anion transporting polypeptide 1A2 (OATP1A2, SLC21A3) (Engel et al., 2012) and organic cation transporters (OCT1-3, SLC22A1-3) (Otter et al., 2017; Soodvilai et al., 2017). These observations indicate that biopharmaceutical considerations need to be an important part of new formulation development when formulations contain pharmaceutical excipients such as surfactants and co-surfactants because these excipients may have different impacts on transporters and carriers. Therefore, addition of surfactants to obtain an enabling formulation may potentially influence the oral absorption (positively or negatively) of a co-administered drug substance if this is a substrate for a carrier and/or transporter. On the other hand, enabling formulations provide the formulation scientist with the possibility to adjust drug absorption to become more consistent by selecting

appropriate excipients for drug substances that are ABC and/or SLC substrates. To do so, it becomes important to understand: 1) the different impacts of nonionic surfactants on carriers and/or transporters, 2) the mechanism behind surfactant-protein interactions, and 3) whether such impacts of surfactants on substrate-protein interactions observed in cell cultures may affect the pharmacokinetics parameters of the substrates *in vivo*. Currently, the translational aspects of how excipients affect carriers and transporters *in vivo* and how this may be exploited for formulation design are largely unexplored.

This review aims to provide an overview of surfactants shown to modulate substrate transport via transporters and carriers, and to discuss the relevance for oral drug absorption, and when possible the mechanism behind the interaction. In this paper, essential data is presented in tables, whereas more comprehensive overviews are provided in [a](#) supplementary tables in order to enhance the readability of the review.

2. Modulation of intestinal transporters and carriers

In the 1980s, modulation of membrane transport proteins was originally proposed as a strategy for chemo-sensitizing cancer cells and to increase the oral bioavailability of drug substances that were substrates for the efflux transporter P-gp in humans. Therefore, inhibitors of efflux transporters such as P-gp and BCRP were ~~then~~ identified and these inhibitors, for example verapamil (Tsuruo et al., 1981), dexverapamil (Gramatté and Oertel, 1999), valspodar (vanAsperen et al., 1997), and GF120918 (Hyafil et al., 1993), showed promising results in inhibiting P-gp (GF120918 also inhibited BCRP) (Maliepaard et al., 2001a), in pre-clinical studies. However, this strategy failed to produce safe and effective treatment in human clinical trials (Dalton et al., 1995; Greenberg et al., 2004; Lehnert et al.,

1998; Mross et al., 1999; Planting et al., 2005; Ries and Dicato, 1991; Sparreboom et al., 1999; Warner et al., 1998). Subsequently, other strategies based on natural products (Appendino et al., 2003; Yoshida et al., 2005) and pharmaceutical excipients (Lo, 2003; Rege et al., 2002; Regev et al., 1999; Zhang et al., 2003) were suggested. In the latter group of compounds, nonionic surfactants gained quite some attention since many surfactants were found to enhance the intracellular accumulation of anticancer drugs including daunorubicin, vinblastine, and etoposide in cancer cells (Buckingham et al., 1995; Woodcock et al., 1990), and to influence the translocation activity of several transporters (Batrakova et al., 2001; Lo, 2003; Rege et al., 2002) and carriers (Rege et al., 2002). In terms of intestinal absorption, it has been reported that nonionic surfactants such as cremophor® EL (Rege et al., 2002), polysorbate 80 (Lo, 2003), and d- α -tocopheryl polyethylene glycol 1000 succinate (TPGS 1000) (Bogman et al., 2005) increased the absorptive permeability and decreased the secretory permeability of the P-gp substrates drug substances rhodamine 123, epirubicin, and talinolol, respectively, *in vitro* using the Caco-2 cell monolayers model. Likewise, in intestinal segments of rats, surfactants such as TPGS 1000 (Varma and Panchagnula, 2005) and polysorbate 40 (Zhu et al., 2009) enhanced mucosal to serosal (M-S) permeability and decreased S-M permeability of P-gp substrates paclitaxel and rhodamine 123, respectively. Moreover, *in vivo*, it has been shown that Solutol® HS 15 (Bittner et al., 2002), polysorbate 80 (Zhang et al., 2003), and pluronic® P85 (Föger et al., 2006) enhanced the oral absorption and exposure of P-gp substrates colchicine, digoxin, and rhodamine 123, respectively. Consequently, it was evident that nonionic surfactants could be potential alternatives to conventional P-gp inhibitors.

Since the majority of research performed until now, has been focused on investigating the influence of nonionic surfactants on P-gp, BCRP, and MRP2 transporters, and on carriers including monocarboxylic

acid transporter (MCT), OATP1A2, OATP2B1, OCT1-3 and peptide transporters 1 and 2 (PEPT1 and 2), this review will therefore summarize and discuss the impact of nonionic surfactants on these transporters and carriers *in vitro* and on the intestinal absorption of substrate drug substances *in vivo*.

3. Expression of selected transporters and carriers along the human intestine

The intestinal expression of transporters and carriers have been studied for years using various mRNA-based techniques, such as northern blotting and RT-PCR, as well as protein quantification methods, e.g. western blotting. As an example, Broberg et al. studied the expression of the proton-coupled amino acid transporter PAT1 along the length of the rat intestine taking samples from each 5 cm and measured the *pat1* mRNA expression in each segment (Broberg et al., 2012). Recently, the emerging of high-resolution MS/MS-based techniques, membrane proteomics has received attention as a tool to describe the absolute transporter or carrier abundance in enterocytes. Knowing transporter and carrier abundancies in the intestine is important for understanding the absorption windows of drug substances (Oswald et al., 2006) and drug-drug interactions (Giacomini et al., 2010). Furthermore, absolute transporter and carrier abundance in different segments of the intestine is necessary for PBPK modelling (Harwood et al., 2013). The present section will briefly review the transporter and carrier abundance in the human intestine and Caco-2 cells of the selected transporter and carriers discussed in the present review.

Table 1 shows protein concentrations of selected transporters and carriers in Caco-2 cells and in different regions of the human intestine. Naturally, cell differentiation affects transport and carrier protein expression, which ~~is~~was also shown by Uchida et al. (Uchida et al., 2015). Therefore, we only

compare expression levels in cells with a similar degree of differentiation, and all three studies have ~~shown~~ investigated the expression in Caco-2 cells cultured for three weeks. Expression levels in cells obtained from different cell banks may vary to a high degree as illustrated for the glucose carrier SGLT1 by Steffansen and co-workers (Steffansen et al., 2017). For this reason, we have compared studies of Caco-2 cells from different cell banks (DSMZ, ECACC, and ATCC). The most extensive and systematic work in regional transporter and carrier expression in the intestine has been performed by Drozdzik and co-workers (Drozdzik et al., 2019; Drozdzik et al., 2014)

Regarding transporters, the P-gp expression ranged ds from low to high (Akazawa et al., 2018; Drozdzik et al., 2019; Drozdzik et al., 2014; Gröer et al., 2013; Harwood et al., 2015; Lloret-Linares et al., 2016), and the expression increased ds from the proximal small intestine towards the distal small intestine (Akazawa et al., 2018; Drozdzik et al., 2014; Gröer et al., 2013) and ~~drops~~ decreased in the colon to levels similar to those in the proximal small intestine (Table 1) (Drozdzik et al., 2019; Drozdzik et al., 2014). Caco-2 cells expressed ed slightly elevated levels of P-gp, compared to the small intestine (Brück et al., 2017; Uchida et al., 2015; Ölander et al., 2016). BCRP expression varied ds between the studies from low (Drozdzik et al., 2014) to very high (Akazawa et al., 2018) expression in the small intestine with an increasing expression from the proximal to the distal part of small intestine (Drozdzik et al., 2019; Drozdzik et al., 2014). In the colon, very low to intermediate expression of BCRP has been reported (Drozdzik et al., 2019; Drozdzik et al., 2014). Similarly, reported expression of BCRP in Caco-2 cells varied ds greatly. MRP2 and MRP3 generally seem were intermediately to highly expressed along the entire intestine (Akazawa et al., 2018; Drozdzik et al., 2019; Drozdzik et al., 2014; Gröer et al., 2013; Harwood et al., 2015) with a slight tendency of elevated expression in the colon (Drozdzik et al., 2019; Drozdzik et al., 2014). The expression of MRP2 in Caco-2 cells is ~~was~~ similar to the

expression in the small intestine (Brück et al., 2017; Uchida et al., 2015) with one exception, where lower expression was reported (Ölander et al., 2016). Ölander and co-workers quantified MRP3 in Caco-2 cells, and the expression of this protein was similar to the expression in the small intestine (Ölander et al., 2016). Brück et. al. and Uchida et. al. could not quantify MRP3 in Caco-2 cells (Brück et al., 2017; Uchida et al., 2015).

Regarding carriers, PEPT1 exhibit~~ed~~^{eds} high to very high expression in the small intestine (Akazawa et al., 2018; Drozdik et al., 2019; Drozdik et al., 2014; Gröer et al., 2013; Miyauchi et al., 2016), and increasing expression in the proximal to distal direction in the small intestine (Drozdik et al., 2019; Drozdik et al., 2014), however, the expression in the colon ~~is~~^{was} highly reduced (approx. 10- to 30-fold) (Drozdik et al., 2019; Drozdik et al., 2014); ~~(see~~^{see} Table 1). In Caco-2 cells, PEPT1 show~~ed~~^{eds} different levels of expression from colon-like levels to small intestine-like levels (Brück et al., 2017; Uchida et al., 2015; Ölander et al., 2016) (Table 1).

Absolute amounts of transporters and carriers in the human gastrointestinal tract highly depend on the site of sampling for the determination. Therefore, it is important to note that the site of sampling from anatomical structures varies between studies in the field (Drozdik et al., 2019; Drozdik et al., 2014; Lloret-Linares et al., 2016; Miyauchi et al., 2016). Additionally, tissue samples are occasionally only defined as ‘jejunal’, ‘ileal’, ‘distal jejunum’, or ‘distal ileum’ with no further definition (Akazawa et al., 2018; Gröer et al., 2013; Harwood et al., 2015). Likewise, great inter-individual variation of intestinal transporter and carrier protein expression is likely, and we have left out all statistical deviation parameters in (Table 1) to enhance the overview. Moreover, protein expression is affected by external factors, for example certain drug compounds (Lin and Yamazaki, 2003) and dietary elements (Erickson

et al., 1995) along with general health condition and diseases (Englund et al., 2007; Wojtal et al., 2009).

Finally, inter-laboratory variation is a well-documented factor, and we refer to the excellent cross-laboratory study by Wegler and co-workers (Wegler et al., 2017). Herein, the authors have shown large variabilities, depending on the methods applied for quantification, especially when they compare whole-lysate and membrane fractionation techniques in the sample preparation. Depending on the drug formulation, pharmaceutical excipients, such as nonionic surfactants, are likely to be present at different concentrations in different segments of the intestine. To firmly understand how the excipients will affect drug absorption influenced by transporters or carriers, it is crucial to obtain more PBPK modelling knowledge regarding transporter and carrier expression patterns.

4. Nonionic surfactants modulate the transport activity of several ABC transporters

4.1 Impact of nonionic surfactants on P-glycoprotein

The first discovered member of the ABC family, P-glycoprotein (Juliano and Ling, 1976), has until now been the most investigated transporter. ~~Since P-gp was found to mediate the cellular efflux of drug substances belonging to different drug classes, e.g. anticancer drugs and antibiotics, extensive research has been performed to modulate the transport activity of the transporter *in vitro* and *in vivo*. The next sections will focus on P-gp molecular characterization and substrate transport via P-gp, and cellular expression and tissue distribution of P-gp. In the subsequent sections, the effects of surfactants and co-surfactants on P-gp activity *in vitro* and on the intestinal absorption of P-gp substrates *in vivo* will be discussed.~~

~~P-glycoprotein molecular characterization and substrate transport~~

P-glycoprotein is a 170 kDa efflux transporter that requires energy from ATP hydrolysis to pump substrates out or across cellular membranes. It has been estimated that P-gp requires two ATP molecules to transport one substrate (Ambudkar et al., 1997; Sauna and Ambudkar, 2001). Due to the direct use of ATP, the transporter is able to transport its substrates against the concentration gradient, and therefore ~~P-gp can~~ limit the cellular accumulation and retention of certain drug substances. In human, P-gp is expressed in the cell membrane in different tissues such as the luminal membrane in enterocytes, the canalicular membrane in hepatocytes, in the luminal membrane in proximal tubular cell, in the luminal membrane of endothelial cells of the central nervous system and testes, bronchial cells of lungs, and placenta (Cordon-Cardo et al., 1990), hence affecting the ADMET properties of its substrates. P-gp substrates include a wide range of hydrophobic and amphipathic substrates such as drug substances and toxins with diverse molecular weight ranging from approximately 300-4000 Da (Fromm, 2004; Rao et al., 1999; Su et al., 2009). Hundreds of drug substances are P-gp substrates (Drugbank, 2019) including anti-cancer drugs, antibiotics, cardiac drugs, immuno-suppressants, lipid-lowering agents, HIV drugs, and hormones (Chan et al., 2004; Fromm, 2004; Seelig, 1998).

~~It has previously been suggested that P-gp substrates might diffuse through the membrane bilayer and reach the cytoplasmic leaflet, where the substrate gets access to the protein (Raviv et al., 1990). P-gp will then according to this mechanistic proposal act as a hydrophobic “vacuum cleaner” that pumps the substrate to the extracellular environment (Raviv et al., 1990). Another model referred as “flippase model” assumed that the P-gp substrate first partition into the lipid bilayer and reach the inner leaflet, where the substrate has access to P-gp. At this stage, the substrate will be pumped directly to the extracellular environment by P-gp or flipped by the protein to the outer leaflet of the lipid bilayer~~

(Higgins and Gottesman, 1992). In both cases, the movement of the substrate would be driven by either the equilibrium between the concentration of the substrate in the extracellular environment and in the outer leaflet, or between the concentration of the substrate in the inner leaflet and in the cytoplasm (Higgins and Gottesman, 1992). There is accumulating evidence supporting the flippase model for P-gp mediated substrate transport across the membrane bilayer (Abulrob and Gumbleton, 1999; Eckford and Sharom, 2005; Romsicki and Sharom, 2001; van Helvoort et al., 1996).

It was proposed that the transmembrane (TM) organization of P-gp consists of two TM domains (Chen et al., 1986), which each domain has six TM helices and one nucleotide-binding domain (NBD) located in the cytoplasm (Fig. 1a) (Chen et al., 1986) (Fig. 1a). Recently, the inward-facing conformation of mouse P-gp proposed that the TM helices are arranged to form an internal cavity of approximately 6000 Å³, which is integrated in the lipid bilayer (Aller et al., 2009). It has been suggested that P-gp might have portals open to the cytoplasmic region and to the inner leaflet of the lipid bilayer (Aller et al., 2009). The P-gp substrates may therefore via these portals get access to the binding sites in the internal cavity of the P-gp, where two P-gp substrates could simultaneously be accommodated (Aller et al., 2009). By using cryo-electron microscopy at 3.4 Å resolution, recent research has also shown the outward-facing conformation of human P-gp (Kim and Chen, 2018). In brief, a P-gp substrate may bind to the internal cavity of the inward-facing conformation of P-gp, as it has been suggested by Aller and co-workers (Aller et al., 2009), and this initiates ATP binding to the NBDs of the protein (Kim and Chen, 2018). In the process of reaching the outward-facing conformation, the NBDs may dimerize resulting in the NBD's becoming closer, while the TM helices re-arrange toward the extracellular space and compress, preventing the binding of the substrate to P-gp (Kim and Chen, 2018) (Fig. 1a). In its outward-facing conformation, the extracellular part of TM helices seems to be flexible for the substrate release and transport to the extracellular environment.

ATP hydrolysis will then reset the P-gp to the inward-facing conformation (Kim and Chen, 2018). ~~as it has been illustrated for the mouse P-gp (Aller et al., 2009). The inward and outward facing conformations of P-gp which were proposed to occur during the substrates translocation may support that the substrates transport across the membrane bilayer most likely occur by the flippase model.~~

4.1.1 Nonionic surfactants inhibit P-glycoprotein *in vitro* via different mechanisms

The most investigated nonionic surfactants that inhibit P-gp transport *in vitro* are cremophor[®] EL, Solutol[®] HS 15, TPGS 1000, polysorbate 20, polysorbate 80 and pluronic[®] P85, (see Table 2). Several *in vitro* assays have been utilized to investigate the P-gp inhibitory properties of these surfactants including: 1) measuring the impact of surfactants on substrate absorptive and/or secretory transport using cell-based systems or intestinal segments excised from animals, 2) measuring the ATPase activity ~~of P-gp~~ using membrane vesicles from cells overexpressing P-gp, and 3) fluorescence-based (e.g. calcein-AM) efflux assay using cells ~~highly overexpressing~~ P-gp. In bi-directional transport assays, nonionic surfactants were shown to enhance the absorptive permeability of ~~model~~ P-gp substrates such as digoxin (Al-Ali et al., 2018b; Batrakova et al., 2001; Collnot et al., 2010; Nielsen et al., 2016) and rhodamine 123 (Collnot et al., 2010; Guan et al., 2011; Kiss et al., 2014; Rege et al., 2002; Sachs-Barrable et al., 2007; Zhao et al., 2016), and ~~to~~ decrease the secretory permeability ~~of these substrates~~ across cell monolayers. Consequently, the data presented in Table 2 strongly support that these surfactants inhibit the efflux activity of P-gp, thus provide a promising approach to inhibit P-gp-mediated efflux of drug substances. In addition, Table S1 provides a comprehensive overview of many other nonionic surfactants and co-surfactants that have shown different abilities to inhibit P-gp-

mediated transport *in vitro*. ~~The results provided in this table may assist providing an overview of likely interactions.~~

Different mechanisms have been proposed to explain the mechanisms behind how nonionic surfactants increase drug absorption by inhibiting P-gp-mediated cellular efflux (see Fig. 2). Work from Seelig and co-worker has suggested that the inhibition may occur through partitioning of the hydrophobic tail of the surfactant into the cell membrane, while the hydrogen bond acceptor groups in the hydrophilic moiety of the surfactant form hydrogen bonds with the hydrogen bond donor groups in the TM domain of the protein (Li-Blatter et al., 2009; Seelig and Gerebtzoff, 2006), (Fig. 2). These hydrogen bonds between the surfactant and the TM domain in P-gp may explain the higher affinity of surfactants with large number of hydrogen bond acceptor groups such as n-octyl- β -D-maltopyranoside (C₈-malt) and 3-cyclohexyl-1-propyl- β -D-maltopyranoside (Cymal-3), than surfactants with fewer hydrogen bond acceptor groups e.g. n-heptyl- β -D-glucopyranoside (C₇-gluc). Thus C₈-malt and Cymal-3 with a similar number of hydrogen bond acceptor groups exhibit almost the same affinity to the membrane as C₇-gluc, but a higher affinity to the P-gp protein, due to the duplication of the sugar moiety in maltopyranoside based-surfactants compared to the single sugar moiety in glucopyranoside (Li-Blatter et al., 2012; Li-Blatter et al., 2009; Li-Blatter and Seelig, 2010; Xu et al., 2015). This observation may be supported by the results from a recent study (Al-Ali et al., 2018b), which reported that polysorbate 20 elicited higher affinity to P-gp in the calcein-AM efflux assay than the mono-saccharide based surfactants, e.g. lauroyl methyl glucamide, and the di-saccharides based surfactants, e.g. lauryl- β -D-maltoside, since the former surfactant had a higher number of hydrogen acceptor groups when retaining the laurate side chain in the surfactants (Al-Ali et al., 2018b). The latter study also suggested that extending the alkyl side chain to more than laurate e.g. stearate in polysorbate 60 and oleate in polysorbate 80, or attaching multiple

alkyl groups such as tri-stearate in polysorbate 65, while retaining the hydrophilic group in these surfactants, may decrease the affinity of the surfactant to P-gp (Al-Ali et al., 2018b). Therefore, the study concluded that *in vitro* both the hydrophobic and hydrophilic moieties in nonionic surfactant may contribute to the surfactant mediated P-gp inhibition.

The second proposed mechanism of P-gp inhibition by nonionic surfactants relates to the alteration of membrane bilayer fluidity induced by surfactants, an alteration that might indirectly inhibit the ATPase activity (Fig. 2). It was shown that polysorbate 20, Nonidet™ P-40 and Triton™ X-100, which all increase the fluidity of artificial membranes, inhibited P-gp ATPase activity in membrane vesicles prepared from Chinese hamster ovary AA8 cells (Regev et al., 1999), (Table 2). In addition, it was reported that polysorbate 80 and cremophor® EL, which increased membrane fluidity significantly, also inhibited P-gp. The inhibition resulted in a significant increase in the absorptive permeability across Caco-2 cells of the model P-gp substrate rhodamine 123 and a significant decrease in the secretory permeability (Rege et al., 2002). In a subsequent study, the surfactant N-octyl glucoside did not modulate membrane bilayer fluidity and did not change the absorptive and secretory permeability of rhodamine 123 (Rege et al., 2002).

Wei and co-workers proposed that the intracellular depletion of ATP was the main mechanism of P-gp inhibition by pluronic-based surfactants such as pluronic® P123 suggesting a third mechanism of P-gp inhibition (Wei et al., 2010; Wei et al., 2013), (see Fig. 2). This third suggested mechanism of P-gp inhibition is supported by previous studies which reported that pluronic® P85 and pluronic® L64 enhanced the intracellular accumulation of the P-gp substrate rhodamine 123 and decreased intracellular ATP *in vitro* (Batrakova et al., 2003a), (Table 2).

The fourth proposed mechanism of surfactant mediated P-gp inhibition suggested a combined effect of depleted intracellular ATP and alteration in cellular membrane fluidity (Batrakova et al., 2003a; Batrakova et al., 2003b), (see Fig. 2). In support of this, it was reported that pluronic® P85 and pluronic® L81, which enhanced membrane fluidity, were able to deplete the intracellular ATP, and significantly enhanced the intracellular accumulation of rhodamine 123 in bovine brain microvessel endothelial cells (Batrakova et al., 2003b; Batrakova et al., 2004), (Table S1). Consequently, the mechanisms of P-gp inhibition by surfactants seem complex. It could be that one or more mechanisms or more are involved in such inhibition. However, further research focused on further characterizing the underlying mechanism(s) of surfactant-mediated P-gp inhibition is needed, which might assist in choosing the appropriate surfactant(s) or developing new surfactant(s) that could be more potent than the ones available and perhaps transporter specific.

4.1.24.1.1 Surfactants used in preparing lipid-based formulations may inhibit P-glycoprotein *in vitro*

Surfactants can be used in pharmaceutical formulations such as lipid-based formulations (LBF) (Pouton, 2006). Currently, many drug substances available in the market are incorporated into LBFs such as tipranavir (Aptivus®), bexarotene (Targretin®), and sirolimus (Rapamune®), where the surfactants function as solubilizing agents and emulsifiers (Savla et al., 2017). For formulations of P-gp substrates, inclusion of surfactants that have P-gp inhibition properties might be advantageous with respect to enhancement of substrate transport across biological membranes. Therefore, studies have reported the use of the LBFs to enhance the oral absorption of different P-gp substrates (Akhtar et al., 2015; Zhao et al., 2013). The self-micro-emulsifying drug delivery systems (SMEDDS) containing

cremophor[®] RH 40, cremophor[®] EL, or polysorbate 80 (Zhao et al., 2013), and the self-nano-emulsifying drug delivery systems (SNEDDS) containing cremophor[®] RH40 and Transcutol[®] P (Akhtar et al., 2015) were shown to enhance etoposide permeability across intestinal tissues and cell monolayers partly due to the inhibition of P-gp by these surfactants. In human, several studies showed enhanced oral absorption of the P-gp substrate cyclosporine A when formulated in LBFs compared to the oral absorption of cyclosporine A from conventional oral dosage forms (Bekerman et al., 2004; Drewe et al., 1992; Postolache et al., 2002). Cyclosporine A is an immunosuppressant used in prophylaxis and treatment of graft rejection in organ transplantations, and in treatment of autoimmune diseases e.g. rheumatoid arthritis, aplastic anemia, and myasthenia gravis (Italia et al., 2006). Cyclosporine A has low aqueous solubility (0.04 mg mL⁻¹) (O'Leary et al., 1986), low permeability in cell cultures (Augustijns et al., 1993; Fricker et al., 1996), and high variations in oral bioavailability among patients (Czogalla, 2009; Lown, 1997). In the studies where cyclosporine A was prepared in LBFs (Bekerman et al., 2004; Drewe et al., 1992; Postolache et al., 2002), the possible P-gp inhibition effect of excipients, e.g. nonionic surfactants, was however not mentioned. Interestingly, the excipients used to prepare LBFs-containing cyclosporine A were polysorbate 80, cremophor RH 40 (Bekerman et al., 2004), sucrose monolaurate, hydrogenated castor oil, and polyethylene glycol (Drewe et al., 1992), which were later shown to possess P-gp inhibitory properties *in vitro* (Al-Ali et al., 2018a; Al-Saraf et al., 2016; Ashiru-Oredope et al., 2011; Chiu et al., 2003; Cornaire et al., 2004; Gurjar et al., 2018; Hanke et al., 2010; Hodaei et al., 2015; Hugger et al., 2002; Johnson et al., 2002; Kiss et al., 2014; Rege et al., 2002; Shono et al., 2004) (Table 2 and S1), and/or *in vivo* (Shimomura et al., 2016; Zhang et al., 2003; Zhao et al., 2013) (Table 3).

Furthermore, previous research have reported that several surfactants such as lauroyl methyl glucamide, lauryl- β -D-maltoside, and trehalose 6-laurate that inhibited the efflux of P-gp substrate calcein-AM in MDCKII MDR1 cells, (Table S1), might also possess paracellular and/or transcellular permeation enhancing effects in cell cultures (Al-Ali et al., 2018b; Eley and Triumalashetty, 2001; Petersen et al., 2012). Such effects might be advantageous when designing LBFs to enhance the oral absorption of P-gp substrate drug substances with limited oral bioavailability induced by intestinal P-gp.

Additionally, the mixed micelles formulations such as pluronic[®] 105/pluronic[®] F-127, pluronic[®] P123/pluronic[®] F127, and polysorbate 80/pluronic[®] F-127 were shown to inhibit the P-gp-mediated efflux of docetaxel (Chen et al., 2013), paclitaxel (Wei et al., 2010), and morin (Choi et al., 2015), respectively, in cells highly over-expressing P-gp (Table S1). Despite the fact, that pluronic[®] F-127 was used in the latter formulations and proposed to inhibit P-gp in another study (Guan et al., 2011), several studies reported that pluronic[®] F-127 did not inhibit P-gp mediated efflux of ~~several~~ P-gp substrates e.g. such as rhodamine 123 (Batrakova et al., 2003b; Wei et al., 2013), nelfinavir (Shaik et al., 2008), etoposide (Al-Ali et al., 2018a), and digoxin (Gurjar et al., 2018). In these types of formulations, one limitation could be that the drug substances might also be adsorbed to the core of surfactant micelles, which may decrease the free fraction of unbound substrate in the formulation, thus affecting the subsequent oral absorption and bioavailability of the substrate *in vivo*. Therefore, investigating the release of drug substances from micelles is important and should be performed *in vitro* in order to avoid or understand such impacts of surfactants *in vivo*. The LBFs containing nonionic surfactants that have P-gp inhibitory properties seem, however, as a promising approach to improve P-gp substrates permeability across cellular membranes.

4.1.34.1.2 Nonionic surfactants inhibited P-glycoprotein *in vitro* at below and above critical micelle concentrations

In several studies performed in cell cultures, it was observed that the inhibition of P-gp transport activity by nonionic surfactants decreased at concentrations at or above the surfactants critical micelles concentration (CMC) compared to concentration below CMC. It was reported that pluronic® P85 (Batrakova et al., 2003b; Batrakova et al., 2004), cremophor® EL (Shono et al., 2004), and polysorbate 40 (Zhu et al., 2009) were effective in inhibiting P-gp transport activity at concentrations lower than their CMCs *in vitro*; however, this inhibitory effect decreased at or above the surfactants CMC values (Batrakova et al., 2003a; Batrakova et al., 2003b; Batrakova et al., 2004; Shono et al., 2004; Zhu et al., 2009) (Table 2 and S1). A possible explanation to this observation could be that most P-gp substrates are lipophilic substances, which after incorporation into the hydrophobic core of micelles, lead to a decrease in the unbound fraction of the substrate available for the transcellular transport, and hence the observed reduction in substrate transport, termed the solubility/permeability interplay (Beig et al., 2017; Beig et al., 2015; Dahan et al., 2010; Miller et al., 2011).

In addition, using the parallel artificial membrane permeability assay (PAMPA), the passive permeability of the P-gp substrate paclitaxel across artificial membranes have been shown to decrease significantly when the TPGS 1000 concentration in the donor chamber was above the CMC (Varma and Panchagnula, 2005). Furthermore, using PAMPA, increasing the concentration of the surfactants such as sodium lauryl sulfate above the CMC values decreased the passive permeability of etoposide across artificial membranes (Beig et al., 2015). The PAMPA studies may support the hypothesis that the decreased permeability of the substrates was related to the incorporation of paclitaxel and etoposide

in the micelles, this may support the effect of surfactant on the thermodynamic activity of the substrate rather than the decreased inhibitory effect of surfactant above the CMC.

Moreover, many studies have reported that nonionic surfactants, used at concentrations above their CMCs, inhibited the efflux of P-gp substrates *in vitro* such as (Surfactant: P-gp substrate/s): (cremophor® RH 40: rhodamine 123) (Kiss et al., 2014), (Brij® 58: digoxin and rhodamine 123) (Gurjar et al., 2018; Zhao et al., 2016), (cremophor® EL: etoposide and digoxin) (Al-Ali et al., 2018a; Gurjar et al., 2018), (Labrasol®: rhodamine 123) (Lin et al., 2007), and (polysorbate 20: etoposide, doxorubicin, digoxin, and epirubicin) (Al-Ali et al., 2018b; Al-Saraf et al., 2016; Gurjar et al., 2018; Lo, 2003). Consequently, it seems that nonionic surfactants at concentration higher than their CMC values are able to inhibit P-gp *in vitro*; however, care should be taken in the interpretation due to the potential influence ~~for~~^{of} the solubility/permeability interplay.

4.1.44.1.3 Polyethylene glycol (PEG) derivatives inhibited P-glycoprotein *in vitro*

In addition to nonionic surfactants, co-surfactants such as polyethylene glycol (PEG) derivatives are used in a broad spectrum of drug delivery systems where these excipients are used as solubilizers, stabilizers, release-modifiers, and bioavailability enhancers (D'Souza and Shegokar, 2016). From Table 2 and S1, it can be noticed that several PEG derivatives with different molecular weight including PEG 300, PEG 400, PEG 2000, PEG 6000, and PEG 20000 were reported to decrease the efflux of several P-gp substrates in Caco-2 cells and in rat intestinal segments. *In vitro*, it has been shown that PEG 400 at concentration of 0.1-20% (w/v, or v/v) may decrease the efflux of several P-gp substrates e.g. digoxin (Johnson et al., 2002), ranitidine (Ashiru-Oredope et al., 2011), and rhodamine 123 (Hodaei et

al., 2015; Shen et al., 2006). Furthermore, recent studies have reported that PEG 400 may decrease the P-gp expression in Caco-2 cells (Hodaei et al., 2015) and increase the P-gp ATPase activity (Ashiru-Oredope et al., 2011), suggesting two different mechanism of P-gp inhibition. The latter effect might refer to the direct interaction of PEG 400 with P-gp, thus competitively inhibited the protein (Ashiru-Oredope et al., 2011). Moreover, PEG 300 was reported to inhibit P-gp through altering Caco-2 membrane fluidity (Hugger et al., 2002), suggesting a third mechanism of PEG derivatives mediated P-gp inhibition *in vitro*.

4.1.54.1.4 Nonionic surfactants increased the oral absorption of P-glycoprotein substrates *in vivo*

The impact of nonionic surfactants on the oral absorption of P-gp substrates have mainly been investigated in wild type rats (Table 3). Several surfactants such as polysorbate 20 and 80 have been shown to increase the oral absorption of different P-gp substrates ~~such as polysorbate 80, which has been shown to enhance the oral absorption of~~ such as digoxin (Zhang et al., 2003), etoposide (Zhao et al., 2013), and rifampicin (Shimomura et al., 2016). The concentrations of surfactants that increased the oral absorption of different P-gp substrates range from 1-25% (w/v) ~~(see~~ Table 3). Recently, wild type and *mdr1a* deficient rats have been used to investigate the role of intestinal P-gp for the oral absorption of digoxin (Nielsen et al., 2016) and etoposide (Al-Ali et al., 2018a). In these studies, there was approximately 2- and 8-fold increase in the AUC of digoxin and etoposide, respectively, in *mdr1a* deficient rats compared to wild type rats. When 5% and 10-25 % (v/v) polysorbate 20 was co-administered with etoposide and digoxin, respectively, in wild type rats, the oral bioavailability was enhanced significantly. However, in *mdr1a* deficient rats, the presence or absence of similar doses of the surfactant did not influence the bioavailability indicating that the enhanced oral absorption in the

wild type rats was most likely related to P-gp inhibition effects mediated by polysorbate 20 rather than enhancement of the substrate solubility by the surfactant.

In relation to scaling between *in vitro* and *in vivo* studies of the surfactants, it has been shown that a concentration of 20-500 μ M polysorbate 20 decreased the efflux ratio of digoxin and etoposide in cell cultures (Al-Ali et al., 2018a; Nielsen et al., 2016), whereas in pre-clinical studies, in wild type rats, the minimum doses of polysorbate 20 required to increase the oral bioavailability of digoxin (Nielsen et al., 2016) and etoposide (Al-Ali et al., 2018a) were 10% (v/v, 90 mM) and 5% (v/v, 45 mM), respectively. However, in *mdr1a* deficient rats, it was noticed that the oral absorption and bioavailability of etoposide decreased when co-administered with 25% (v/v) polysorbate 20 compared to the oral absorption with 0 or 5% (v/v) polysorbate 20, or without the surfactant (Al-Ali et al., 2018a). *In vitro* dialysis studies demonstrated that etoposide release from a 25% polysorbate 20 containing formulation was minimal, most likely due to the incorporation of etoposide into the micelles formed by the polysorbate (Al-Ali et al., 2018a). Furthermore, it has been reported that 5% labrasol[®] increased the oral bioavailability of etoposide more when co-administered with 5% labrasol[®] than with 10% labrasol[®] (Akhtar et al., 2017), indicating that etoposide release from the micelles was concentration dependent (Akhtar et al., 2017), (see also Table 3).

TPGS 1000 enhanced the oral bioavailability of paclitaxel in wild type rats with a factor of six relative to the bioavailability when administered without the surfactant (control) (Varma and Panchagnula, 2005), (Table 3). Verapamil was further demonstrated to enhance the oral bioavailability of similar doses of paclitaxel four times compared to control (Varma and Panchagnula, 2005). With respect to the use of surfactants *in vivo*, it should be noted that some of these surfactants undergo digestion in the intestinal tract (Christiansen et al., 2010; Cui   et al., 2008; Devraj et al., 2013; Mohsin, 2012), why

and might thus be less efficient ~~the use of these relative to~~ than small-molecular P-gp inhibitors like verapamil, ~~should be considered with respect to the hypothesis of the studies.~~

From Table 2, 3 and S1, it is evident that research is still needed to establish how inhibition observed *in vitro* translates into increased absorption *in vivo*, since many nonionic surfactants such as Brij[®] 78, pluronic[®] P123, and polysorbate 40, have not yet been investigated for their abilities to inhibit intestinal P-gp *in vivo*. Moreover, for some surfactants that were extensively investigated *in vitro* such as cremophor[®] EL, pluronic[®] 85 and Solutol[®] HS 15, (see also Table 2), only few *in vivo* studies have reported the effect ~~on PK~~ of these surfactants on PK in animals (Bittner et al., 2002; Föger et al., 2006; Zhao et al., 2013), (Table 3). Consequently, further *in vivo* studies are needed to advance the knowledge about the effect of ~~these~~ surfactants on the oral absorption of different P-gp substrates, which may assist in designing and performing clinical studies in humans.

4.2 Impact of nonionic surfactants on breast cancer resistance protein

The ABC transporter BCRP is as a monomeric protein of 72 kDa (Doyle and Ross, 2003; Mao, 2005) consisting of one TM domain of six TM helices, and one NBD located in the cytoplasm (Chen et al., 2015; Mao and Unadkat, 2015; Wang et al., 2008). Two BCRP monomers dimerize to form a functional BCRP transporter (Fig. 1b) (Rosenberg et al., 2010; Rosenberg et al., 2015). The helices are arranged to form a cavity, where BCRP substrates bind, while TM helices one and six are attached to amino and carboxyl termini in the cytoplasm, respectively (Wang et al., 2008). BCRP substrates belong to different therapeutic classes such as anticancer drugs, HIV drugs, antihistamines, and anti-hyperlipidemia drugs (Mao and Unadkat, 2015). BCRP shares many substrates with P-gp e.g.

topotecan (Jonker et al., 2000; Maliepaard et al., 1999), doxorubicin (Allen et al., 1999; Mechetner et al., 1998), irinotecan (Gupta et al., 1996; Maliepaard et al., 1999), and etoposide (Allen et al., 2003; Keller et al., 1992). In humans, BCRP is highly expressed in normal tissues such as the apical membrane of small intestinal and colonic enterocytes, canalicular membranes in the liver, endothelial cells of brain microvessels (Mao, 2005), veins and capillaries, and in cancer cells (Doyle and Ross, 2003; Maliepaard et al., 2001b). Since BCRP is expressed in different tissues, its modulation in humans may influence the ADMET properties of its substrates. In the past, BCRP inhibitors were developed to overcome the multidrug resistance phenomenon, as well as to enhance the oral absorption of the substrates (Gupta et al., 2006; Gupta et al., 2004; Houghton et al., 2004; Matsson et al., 2009). BCRP inhibitors may act as competitive inhibitors at the substrate binding sites, as allosteric inhibitors by binding to the protein at a site different from the substrate binding site ~~in the BCRP cavity~~, or by inhibiting ATPase activity (Mao and Unadkat, 2015). Of the BCRP inhibitors identified; some of these also inhibit P-gp e.g. GF120918, and the tyrosine kinase inhibitors imatinib, and the antifungal drug substance itraconazole (Mao and Unadkat, 2015; Matsson et al., 2009).

Until now, few studies were performed by Yamagata and co-workers to have investigated the effect of nonionic surfactants on BCRP transport activity *in vitro* and *in vivo* (Sawangrat et al., 2018a; Sawangrat et al., 2018b; Xiao et al., 2016; Xu et al., 2015; Yamagata et al., 2007a, b; Yamagata et al., 2009). In MDCKII BCRP cells, nonionic surfactants such as cremophor® EL, polysorbate 20, span 20, pluronic® P85, and Brij® 30 increased the uptake of the BCRP substrate mitoxantrone (Yamagata et al., 2007a) (Table 4). Yamagata and coworkers were also able to enhance the uptake of mitoxantrone in MDCKII MDR1 by the use of the same surfactants indicating the ability of these surfactants to modulate both BCRP and P-gp (Yamagata et al., 2007a), effects that wasere demonstrated both *in vitro*

(Al-Ali et al., 2018a; Al-Ali et al., 2018b; Al-Saraf et al., 2016; Gurjar et al., 2018; Li-Blatter and Seelig, 2010; Lo, 2003; Nielsen et al., 2016; Rege et al., 2002; Shaik et al., 2008), and *in vivo* (Al-Ali et al., 2018a; Föger et al., 2006; Nielsen et al., 2016; Zhao et al., 2013), for further details ~~see~~ (see Table 2, 3 and S1). With respect to the effect of nonionic surfactants on BCRP, ~~it another study was~~ reported that pluronic® P85 and polysorbate 20 enhanced the ~~mucosal-to-serosal~~ M-S transport of BCRP substrate topotecan across ileum everted sacs derived from wild type mice (~~see~~ Table 4) (Yamagata et al., 2007b). Interestingly, in everted intestinal sacs derived from *Abcg2* deficient mice, topotecan absorption rate was significantly enhanced in comparison to the absorption rate in everted intestinal sacs from wild type mice (Yamagata et al., 2007b). However, the presence of surfactants did not further improve the absorption rate of topotecan in *Abcg2* deficient everted intestinal sacs, demonstrating the surfactants' impacts in mediating ~~the~~ Bcrp inhibition in the wild type animals (Yamagata et al., 2007b). *In vivo*, pluronic® P85 and polysorbate 20 administered orally 15 min before oral administration of topotecan to wild type mice increased the AUC of topotecan significantly, when compared to the administration of similar doses of topotecan without the surfactant (Yamagata et al., 2007b). It was later noticed that the interaction of pluronic® P85 and polysorbate 20 with Bcrp was reversible and transient upon removal of these surfactants (Yamagata et al., 2009). It is worth noticing that these surfactants were also able to inhibit P-gp *in vitro* (Al-Ali et al., 2018a; Al-Ali et al., 2018b; Al-Saraf et al., 2016; Batrakova et al., 2003a; Batrakova et al., 2004; Gurjar et al., 2018; Nielsen et al., 2016; Shaik et al., 2008) (Table 2, S1) and *in vivo* (Al-Ali et al., 2018a; Föger et al., 2006; Nielsen et al., 2016) (Table 3).

Moreover, Recent research has reported that several surfactants, including 6-tetradecyl- β -D-maltopyranoside, (C6-malt) (Xu et al., 2015), cremophor® EL (Al-Ali et al., 2018a; Al-Saraf et al.,

2016; Gurjar et al., 2018; Rege et al., 2002; Shono et al., 2004). BL-9EX, Brij® 92, Brij® 97 (Zhao et al., 2016), and labrasol® (Akhtar et al., 2017; Cornaire et al., 2004; Lin et al., 2007; Ma et al., 2011), which have been shown to inhibit P-gp (Table 2), ~~have also been able to~~ inhibit BCRP in MDCKII BCRP cells (Xiao et al., 2016), in membrane vesicles containing human BCRP (Xu et al., 2015), and ~~Berp~~ in rat intestinal membrane and *in vivo* using the *in situ* closed intestinal loop method (Sawangrat et al., 2018a). The doses of the surfactants used to inhibit BCRP *in vitro* and *in vivo* (Sawangrat et al., 2018a; Xiao et al., 2016; Xu et al., 2015) were comparable to the doses used to inhibit P-gp (Akhtar et al., 2017; Lin et al., 2007; Ma et al., 2011; Zhao et al., 2016) (Table 2, 3, and S1). As similar doses of nonionic surfactants appeared to be able to inhibit P-gp and BCRP, and since many drug substances that are BCRP substrates share substrate specificity with P-gp, drug formulators should thus consider the surfactants used in their formulations, and avoid using the surfactants that may have overlap in inhibiting effect on both transporters in cases where this may have an influence on the biopharmaceutical properties of the ~~compound~~ substrate. Cremophor® EL was shown to enhance the absorptive permeability and decrease the secretory permeability of scutellarin in MDCKII BCRP (Xiao et al., 2016). Scutellarin is a flavonoid glucuronide approved in China to treat patients with cerebral infarction and paralysis caused by cerebrovascular diseases (Xiao et al 2016). In wild type rats, cremophor® EL enhanced scutellarin oral absorption, however, this study also reported that the surfactant affected other transporters such as MRP2 and MRP3 (See section 4.3). ~~Another recent~~ study by Sawangrat and co-workers showed that 0.05 % (w/v) cremophor® EL enhanced the absorptive permeability and decreased the secretory permeability of topotecan significantly in Caco-2 cells, and enhanced the intestinal absorption of topotecan in rats using the *in situ* closed-loop method (Sawangrat et al., 2018b). Similar concentration of cremophor® EL did, however, not influence the absorptive or secretory permeability of the BCRP substrate sulfasalazine across rat intestinal segments in diffusion

chambers (Sawangrat et al., 2018a). Furthermore, using the *in situ* closed-loop method, 0.05% (w/v) polysorbate 20 enhanced the intestinal absorption of topotecan significantly in rats (Sawangrat et al., 2018b). In contrast; however, higher concentration (0.1 and 0.5% w/v) of the surfactant did not enhance the absorption of sulfasalazine in another study (Sawangrat et al., 2018a). The effect of nonionic surfactants on the transport of BCRP substrates across the intestine therefore seems to differ as a function of substrate and/or method used. Further studies are needed to investigate the effects of nonionic surfactants on the transport activity of BCRP *in vitro* and *in vivo*. It is recommended that different BCRP substrates and different methods are used in the prospective investigations.

4.3 Impact of nonionic surfactants on the multidrug resistance-associated protein 2

In human tissues, the efflux transporter multidrug resistance-associated protein 2 (MRP2) is expressed in the hepatocyte canalicular membrane, gallbladder epithelial cells, the proximal tubule of the kidney, duodenum, jejunum, ileum, brain, bronchi, and placenta (Jedlitschky et al., 2006; Kool et al., 1997; Nies and Keppler, 2007). MRP2 is highly expressed in cancer cells, such as non-small cell lung cancer and adeno-lung carcinoma (Kool et al., 1997). MRP2 consists of two TM domains, each has six TM helices, linked intracellularly with two NBDs located in the cytoplasm (Fig. 1c) (Jedlitschky et al., 2006). To the NBDs, ATP molecules bind, which is required for hydrolysis initiating substrate transport (Jedlitschky et al., 2006). In addition, a third TM domain consisting of five helices is attached to the first TM domain via a linker (L0), which is located in the cytoplasm. The third TM domain is extracellularly attached to NH₂ terminus of the first TM domain (see-Fig. 1c) (Jedlitschky et al., 2006). MRP2 transports different endogenous compounds such as glutathione, leukotrienes, bilirubin

glucuronides and steroids, and drug substances of different classes, e.g. anticancer drugs, HIV drugs, antibiotics, and the metabolites of these substances (Dietrich et al., 2003; Jedlitschky et al., 2006).

The effect of nonionic surfactants on MRP2 has been investigated *in vitro* using different assays, e.g. bi-directional transport, uptake assay, ATP measurements, and phosphate release measurements. It was reported that pluronic® P85 enhanced the intracellular accumulation of the MRP2 substrates vincristine and doxorubicin in MDCKII MRP2 cells (Batrakova et al., 2003a). This was confirmed by decreased ATP levels in MDCKII MRP2 cells and decreased ATPase activity in the membrane vesicles isolated from these cells (Batrakova et al., 2004). Based upon these data, Batrakova and co-workers proposed that the mechanism of MRP2 inhibition could be related to the change in membrane fluidity or binding of the surfactant to the cell membrane, thereby competitively preventing the drug-protein interaction (Batrakova et al., 2004). Beside the effect of pluronic® P85 on MRP2, it was found that similar concentrations of the surfactant inhibited MRP1 and P-gp (Batrakova et al., 2004).

Recent studies have shown that surfactants including cremophor® EL, cremophor® RH 40, pluronic® F68, and pluronic® P-127, and co-surfactants PEG 400, and PEG 2000 decreased the efflux ratio of the MRP2 substrate scutellarin in Caco-2 cells (Li et al 2013, Li et al 2014). ~~Scutellarin is a flavonoid glucuronide approved in China to treat patients with cerebral infarction and paralysis caused by cerebrovascular diseases (Xiao et al 2016).~~ Scutellarin has a poor oral bioavailability, which is partly related to the efflux effect of membrane transporters such as MRP2 and BCRP (see Table 4 and 5). In agreement with Li and co-workers (Li et al 2013, Li et al 2014), Chen *et al.* have shown that pluronic® F68, pluronic® F-127, pluronic® P85, and pluronic® P105 increased A-B permeability and decreased B-A permeability of the MRP2 substrate, baicalein, in MDCK MRP2 cells (Chen et al., 2017), (Table 5).

602 Chen and co-workers have suggested that these observations were due to MRP2 inhibition (Chen et al.,
603 2017).

604 In contrast, vinblastine transport across MDCKII MRP2 cells was not affected significantly by 0.1 %
605 (w/v) pluronic® L61 (Evers et al., 2000). Likewise, using a 5-chloromethylfluorescein diacetate
606 (CMFDA) based accumulation assay in MDCK-MRP2 cells, Bogman *et al.* (2003) found that the
607 surfactants TPGS 1000, cremophor® EL, polysorbate 80, pluronic® F68, pluronic® L61, and pluronic®
608 L81 were unable to inhibit MRP2-mediated methylfluorescein-sulfogluthathione complex (MF-SG)
609 transport (Bogman et al., 2003).

610 From Table 4 and 5, it can be noticed that the dose of cremophor® EL needed to inhibit MRP2 or
611 BCRP in cell cultures and in rats are similar (Xiao et al 2016). Xiao and co-workers have also reported
612 that cremophor® EL was able to activate the efflux protein MRP3 (Xiao et al 2016), which was found
613 to be expressed on the basolateral membrane of enterocytes (Kool et al., 1997; Kool et al., 1999).
614 Therefore, effects of cremophor® EL on scutellarin seems to be related to the effect on multiple efflux
615 transporters. Thus, activating the efflux transporters, being located in the basolateral membrane in
616 enterocytes by surfactants could also be a strategy to improve the absorption of substrate drug
617 substances across intestinal membranes; however, further investigations are needed for this to be a
618 robust formulation strategy.

619 An interesting finding reported was that pluronic® F-127 decreased the efflux ratio of scutellarin and
620 baicalein in Caco-2 cells (Li et al 2013, Li et al 2014) and MDCKII MDRP2 (Chen et al., 2017),
621 respectively, but had no inhibitory effect on the P-gp substrates rhodamine 123 (Batrakova et al.,
622 2003b), nelfinavir (Shaik et al., 2008), etoposide (Al-Ali et al., 2018a), and digoxin (Gurjar et al.,
623 2018). Consequently, it may be that there is limited cross inhibitory effects for surfactants inhibiting

MRP2 towards other efflux membrane transporters and vice versa. The ability of nonionic surfactants to inhibit MRP2-mediated transport seems to be complex and dependent on the model system employed to investigate and understand the influence of the surfactant. Therefore, further studies and specific MRP2 model systems are needed to understand the consequences of MRP2 inhibition by nonionic surfactants.

5. Nonionic surfactants modulate solute carriers *in vitro*

In humans, solute carriers transport endogenous and exogenous (Yu Liang Siqi Li Ligong, 2015), charged, and uncharged substrates (Koepsell et al., 2007), in and/or out of cells in different tissues, e.g. intestine, kidney, liver and brain (Giacomini et al., 2010). The SLC family consists of 62 sub-families (HUGO Gene Nomenclature Committee, 2019). For a protein to be assigned to the SLC family they need to be responsible for membrane solute transport and to have an amino acid identity of > 20% to other members of the family (Hediger et al., 2013). Within the SLC family uniporters, symporters and antiporters are found. The symporters may depend on the driving force of ions such as K^+ , Na^+ , H^+ , or Cl^- , and at a cellular level, they are therefore known as secondary-active transporters (e.g. K^+ , Na^+ , or Cl^- -dependent carriers) or tertiary-active transporters (H^+ -dependent carriers), because the cellular homeostasis of ions eventually will involve transport by the active Na^+/K^+ -ATPase enzyme. The use of transporter in this context is a reminiscence of a notion and literature source present prior to the establishment of the SLC system, which was pioneered by Hediger (Hediger, 2004; Hediger et al., 2013).

SLC proteins are diverse in their structures, however, the most common predicted folds of this family proteins are the Major Facilitator Superfamily (MFS, LacY) and the Leucine transporter (LeuT) that has folding consisting of 12 and 10 TM helices, respectively. However, some carriers in the SLC family may possess a lower number of helices, e.g. the glucose uniporter that has a unique fold of seven TM helices (Colas et al., 2016). For the LacY like fold (Fig. 1d), the protein is oriented in a V-shape conformation opened to the extracellular side of plasma membrane where the substrate is assumed to bind. The substrate may then move to an intermediate state inside the protein, before it may release from the inverted V-shape conformation of the SLC to the cytoplasm (Fig. 1d) (Colas et al., 2016).

The oral absorption of a large variety of important nutrient such as amino acids, sugars, peptides, fatty acids, and vitamins are mediated by carriers, which are important for oral absorption of drug substances that are structurally similar to the nutrients (Steffansen et al., 2004). Despite the large number of carriers expressed in the intestine, the effect of nonionic surfactant on these transport systems is largely uninvestigated. In 2002, Rege and co-workers reported that polysorbate 80 decreased the absorptive permeability of the prototypic PEPT1 substrate glycyl-sarcosine and cremophor® EL decreased the transport of the monocarboxylic acid transporter (MCT) substrate benzoic acid in Caco-2 cells (Rege et al., 2002), (Table 6). Recently, these surfactants have been shown to inhibit other carriers in transfected cells models. Polysorbate 80 and cremophor® EL inhibited OCT1-3 and PEPT2 in MDCKII OCT1-3 cells and MDCKII PEPT2 cells, respectively. From Table 6, it can be noticed that polysorbate 80 appeared more potent than cremophor® EL with respect to inhibition of OCT1-3 and PEPT2 in cell cultures (Otter et al., 2017; Soodvilai et al., 2017). In addition, it was reported that cremophor® EL inhibited OATP1A2 and OATP2B1 in HEK OATP1A2 and HEK OATP2B1 cells, respectively (Engel

et al., 2012). Furthermore, poloxamer 188 and 407 (Otter et al., 2017), and polysorbate 20 and 60 (Otter et al., 2017; Soodvilai et al., 2017), have shown different abilities to inhibit the organic cation transporters in cell cultures. In MDCKII OCT1 cells, the estimated IC_{50} of poloxamer 407 (pluronic® F-127) was approximately 2600-fold and 900-fold higher than the estimated IC_{50} of polysorbate 80 and polysorbate 20, respectively (Otter et al., 2017).

Surfactants inhibiting carriers (Table 6) were also reported to inhibit transporters (Table 2-5, and S1) in cell cultures as exemplified by the observations that Solutol® HS 15 inhibited OATP1A2, OATP2B1 (Engel et al., 2012), OCT1-3, PEPT2 (Otter et al., 2017), and P-gp (Akhtar et al., 2017; Buckingham et al., 1995; Coon et al., 1991; Cornaire et al., 2004; Gurjar et al., 2018; Lamprecht and Benoit, 2006); polysorbate 80 inhibited OCT1-3, PEPT1-2 (Otter et al., 2017; Rege et al., 2002; Soodvilai et al., 2017), P-gp (Al-Ali et al., 2018b; Al-Saraf et al., 2016; Cornaire et al., 2004; Hanke et al., 2010; Kiss et al., 2014; Lo, 2003; Nerurkar et al., 1996; Nielsen et al., 2016; Shono et al., 2004; Woodcock et al., 1992; Yu et al., 2011), and MRP2 (Hanke et al., 2010); and cremophor® EL inhibited OATP1A2, OATP2B1 (Engel et al., 2012), OCT1-3, PEPT2 (Otter et al., 2017), MCT (Rege et al., 2002), P-gp (Al-Ali et al., 2018a; Al-Saraf et al., 2016; Buckingham et al., 1995; Chiu et al., 2003; Nerurkar et al., 1996; Rege et al., 2002; Shono et al., 2004; Woodcock et al., 1990; Woodcock et al., 1992), BCRP (Sawangrat et al., 2018a; Xiao et al., 2016; Yamagata et al., 2007a), and MRP2 (Hanke et al., 2010; Li et al., 2013a; Xiao et al., 2016). Cremophor® EL seems to be the surfactant with the widest range of inhibition of different carriers and transporters. Importantly, impact of surfactant on transporters and carriers simultaneously may lead to unpredictable drug-transporter interactions. Therefore, further knowledge about the ability of nonionic surfactants to inhibit carriers under relevant *in vivo* conditions

is needed for drug formulators to make enlightened choices on nonionic surfactants as pharmaceutical excipients.

As generally presented in this review, a broader class of surfactants that are often used as pharmaceutical excipients, may have effects on drug absorption through interactions with either transporters, or carriers, or both. While a lot of insights have been generated *in vitro*, less is available *in vivo* from non-clinical trials, and no information is publicly available from human trials systematically investigating the influence of pharmaceutical excipients on transporters or carriers. Drug prescribers and pharmacists should therefore be aware that in treatment of patients with polypharmacy prescriptions of drug compounds that are known substrates to transporters or carriers should be administered separately from drug products containing nonionic surfactants and co-surfactant polymers in order to avoid unexpected interactions of drug-excipient at transporters and/or carriers, as this might lead to unpredicted side effects.

6. Conclusion

From the present review, it is quite evident that pharmaceutical excipients are not just compounds required for processing drug formulations, but they also possess the ability to alter drug transport across biological barriers by interacting with transporters and carriers. Pharmaceutical excipients frequently used in enabling formulations, notably nonionic surfactants, alter the function of carriers and/or transporters, thereby affecting drug transport. The main body of evidence for this is based on *in vitro* experiments using cell culture models or excised tissue, whereas pre-clinical studies available in the literature are limited. Few studies have investigated if surfactants reduce transporter-mediated transport

and thereby increase the oral bioavailability; while to the best of our knowledge no *in vivo* study has investigated if excipients inhibit carriers *in vivo*, and hence could decrease oral bioavailability. Therefore, more pre-clinical studies are needed to investigate if surfactants at concentration likely to be reached in the intestinal lumen may alter the exposure of substrates of transporters and carriers. It seems likely that inhibition of transporters by surfactants could be incorporated into a formulation approach, while a potential inhibition of solute carriers should be avoided as this would decrease oral absorption. The key points missing are what the scaling between *in vitro* and *in vivo* effects is and if enabling formulations containing surfactants are safe i.e. without toxic effects. Such biopharmaceutical insight may assist in the active development of formulations where excipients are bioactive components included for inhibition of intestinal efflux transport. Interestingly, some surfactants, e.g. cremophor® EL, Solutol® HS 15 and polysorbate 20, have been shown to share inhibiting effects on several transporters and carriers *in vitro*. In addition, the concentration of surfactants that inhibit the efflux transporters P-gp, BCRP, and MRP2 were comparable in cell cultures. Hence, when drug substances such as doxorubicin and etoposide are substrates for multiple efflux transporters, co-administration of these with surfactants can generate a complex absorption mechanism.

As more and more discovered compounds have limited aqueous solubility, the need for enabling formulations, that may include surfactants, are increasing. Given that surfactants can have multiple physico-chemical as well as biopharmaceutical properties, drug formulators may need to bring this perspective into consideration when defining the formulations of the future.

Declaration of interest

The authors do not have any conflict of interest to report.

Author contribution

Writing - original draft: AAAA, CUN, and RBN. Writing - review & Editing: AAAA, RBN, BS, RH and CUN. Final approval of the version submitted: AAAA, RBN, BS, RH and CUN.

References:

- Abulrob, A.-n.G., Gumbleton, M., 1999. Transport of Phosphatidylcholine in MDR3-Negative Epithelial Cell Lines via Drug-Induced MDR1 P-Glycoprotein. *Biochem. Biophys. Res. Commun.* 262, 121-126.
- Akazawa, T., Uchida, Y., Miyauchi, E., Tachikawa, M., Ohtsuki, S., Terasaki, T., 2018. High Expression of UGT1A1/1A6 in Monkey Small Intestine: Comparison of Protein Expression Levels of Cytochromes P450, UDP-Glucuronosyltransferases, and Transporters in Small Intestine of Cynomolgus Monkey and Human. *Mol. Pharm.* 15, 127-140.
- Akhtar, N., Ahad, A., Khan, M.F., Allaham, A., Talegaonkar, S., 2017. The Ameliorated Pharmacokinetics of VP-16 in Wistar Rats: A Possible Role of P-Glycoprotein Inhibition by Pharmaceutical Excipients. *Eur. J. Drug Metab. Pharmacokinet.* 42, 191-199.

- 746 Akhtar, N., Talegaonkar, S., Ahad, A., Khar, R.K., Jaggi, M., 2015. Potential of a novel self
747 nanoemulsifying carrier system to overcome P-glycoprotein mediated efflux of etoposide: In vitro and
748 ex vivo investigations. *J Drug Deliv Sci Technol* 28, 18-27.
- 749 Al-Ali, A.A.A., Quach, J.R.C., Bundgaard, C., Steffansen, B., Holm, R., Nielsen, C.U., 2018a.
750 Polysorbate 20 alters the oral bioavailability of etoposide in wild type and *mdr1a* deficient Sprague-
751 Dawley rats. *Int. J. Pharm.* 543, 352-360.
- 752 Al-Ali, A.A.A., Steffansen, B., Holm, R., Nielsen, C.U., 2018b. Nonionic surfactants increase digoxin
753 absorption in Caco-2 and MDCKII MDR1 cells: Impact on P-glycoprotein inhibition, barrier function,
754 and repeated cellular exposure. *Int. J. Pharm.* 551, 270-280.
- 755 Al-Saraf, A., Holm, R., Nielsen, C.U., 2016. Tween 20 increases intestinal transport of doxorubicin in
756 vitro but not in vivo. *Int. J. Pharm.* 498, 66-69.
- 757 Allen, J.D., Brinkhuis, R.F., Wijnholds, J., Schinkel, A.H., 1999. The mouse *Bcrp1/Mxr/Abcp* gene:
758 amplification and overexpression in cell lines selected for resistance to topotecan, mitoxantrone, or
759 doxorubicin. *Cancer Res.* 59, 4237.
- 760 Allen, J.D., Van Dort, S.C., Buitelaar, M., van Tellingen, O., Schinkel, A.H., 2003. Mouse breast
761 cancer resistance protein (*Bcrp1/Abcg2*) mediates etoposide resistance and transport, but etoposide oral
762 availability is limited primarily by P-glycoprotein. *Cancer Res.* 63, 1339-1344.

- 763 Aller, S.G., Yu, J., Ward, A., Weng, Y., Chittaboina, S., Zhuo, R., Harrell, P.M., Trinh, Y.T., Zhang,
764 Q., Urbatsch, I.L., Chang, G., 2009. Structure of P-glycoprotein reveals a molecular basis for poly-
765 specific drug binding. *Science* 323, 1718-1722.
- 766 Ambudkar, S.V., Cardarelli, C.O., Pashinsky, I., Stein, W.D., 1997. Relation Between the Turnover
767 Number for Vinblastine Transport and for Vinblastine-stimulated ATP Hydrolysis by Human P-
768 glycoprotein. *J. Biol. Chem.* 272, 21160-21166.
- 769 Appendino, G., Della Porta, C., Conseil, G., Sterner, O., Mercalli, E., Dumontet, C., Di Pietro, A.,
770 Lund, U., Centre for, A., Synthesis, Lunds, u., Centrum för analys och, s., 2003. A new P-glycoprotein
771 inhibitor from the caper spurge (*Euphorbia lathyris*). *J. Nat. Prod.* 66, 140-142.
- 772 Ashiru-Oredope, D.A.I., Patel, N., Patel, R., Forbes, B., Basit, A.W., 2011. The effect of
773 polyoxyethylene polymers on the transport of ranitidine in Caco-2 cell monolayers. *Int. J. Pharm.* 409,
774 164-168.
- 775 Augustijns, P.F., Bradshaw, T.P., Gan, L.S.L., Hendren, R.W., Thakker, D.R., 1993. Evidence for a
776 Polarized Efflux System in Caco-2 Cells Capable of Modulating Cyclosporine A Transport. *Biochem.*
777 *Biophys. Res. Commun.* 197, 360-365.
- 778 Batrakova, E., Lee, S., Li, S., Venne, A., Alakhov, V., Kabanov, A., 1999. Fundamental Relationships
779 Between the Composition of Pluronic Block Copolymers and Their Hypersensitization Effect in MDR
780 Cancer Cells. *Pharm. Res.* 16, 1373-1379.

- 781 Batrakova, E.V., Li, S., Alakhov, V.Y., Elmquist, W.F., Miller, D.W., Kabanov, A.V., 2003a.
782 Sensitization of Cells Overexpressing Multidrug-Resistant Proteins by Pluronic P85. *Pharm. Res.* 20,
783 1581-1590.
- 784 Batrakova, E.V., Li, S., Alakhov, V.Y., Miller, D.W., Kabanov, A.V., 2003b. Optimal structure
785 requirements for pluronic block copolymers in modifying P-glycoprotein drug efflux transporter
786 activity in bovine brain microvessel endothelial cells. *J. Pharmacol. Exp. Ther.* 304, 845-854.
- 787 Batrakova, E.V., Li, S., Li, Y., Alakhov, V.Y., Kabanov, A.V., 2004. Effect of Pluronic P85 on
788 ATPase Activity of Drug Efflux Transporters. *Pharm. Res.* 21, 2226-2233.
- 789 Batrakova, E.V., Miller, D.W., Li, S., Alakhov, V.Y., Kabanov, A.V., Elmquist, W.F., 2001. Pluronic
790 P85 Enhances the Delivery of Digoxin to the Brain: In Vitro and in Vivo Studies. *J. Pharmacol. Exp.*
791 *Ther.* 296, 551.
- 792 Beig, A., Fine-Shamir, N., Porat, D., Lindley, D., Miller, J.M., Dahan, A., 2017. Concomitant
793 solubility-permeability increase: Vitamin E TPGS vs. amorphous solid dispersion as oral delivery
794 systems for etoposide. *Eur. J. Pharm. Biopharm.* 121, 97-103.
- 795 Beig, A., Miller, J.M., Lindley, D., Carr, R.A., Zocharski, P., Agbaria, R., Dahan, A., 2015. Head-To-
796 Head Comparison of Different Solubility-Enabling Formulations of Etoposide and Their Consequent
797 Solubility–Permeability Interplay. *J. Pharm. Sci.* 104, 2941-2947.

- 798 Bekerman, T., Golenser, J., Domb, A., 2004. Cyclosporin Nanoparticulate Lipospheres for Oral
799 Administration. *J. Pharm. Sci.* 93, 1264-1270.
- 800 Bittner, B., Guenzi, A., Fullhardt, P., Zuercher, G., González, R.C.B., Mountfield, R.J., 2002.
801 Improvement of the bioavailability of colchicine in rats by co-administration of D- α -tocopherol
802 polyethylene glycol 1000 succinate and a polyethoxylated derivative of 12-hydroxy-stearic acid.
803 *Arzneimittelforschung* 52, 684-688.
- 804 Bogman, K., Erne-Brand, F., Alsenz, J., Drewe, J., 2003. The role of surfactants in the reversal of
805 active transport mediated by multidrug resistance proteins. *J. Pharm. Sci.* 92, 1250-1261.
- 806 Bogman, K., Zysset, Y., Degen, L., Hopfgartner, G., Gutmann, H., Alsenz, J., Drewe, J., 2005. P-
807 Glycoprotein and Surfactants: Effect on Intestinal Talinolol Absorption. *Clin. Pharmacol. Ther.* 77, 24-
808 32.
- 809 Broberg, M.L., Holm, R., Tønsberg, H., Frølund, S., Ewon, K.B., Nielsen, A.L., Brodin, B., Jensen, A.,
810 Kall, M.A., Christensen, K.V., Nielsen, C.U., 2012. Function and expression of the proton-coupled
811 amino acid transporter PAT1 along the rat gastrointestinal tract: implications for intestinal absorption
812 of gaboxadol. *Br. J. Pharmacol.* 167, 654-665.
- 813 Brück, S., Strohmeier, J., Busch, D., Drozdzik, M., Oswald, S., 2017. Caco-2 cells - expression,
814 regulation and function of drug transporters compared with human jejunal tissue: Transporter
815 Expression, Regulation and Function in Caco-2 Cells. *Biopharm. Drug Disposition* 38, 115-126.

- 816 Buckingham, L.E., Buckingham, L.E., Balasubramanian, M., Emanuele, R.M., Emanuele, R.M.,
817 Clodfelter, K.E., Clodfelter, K.E., Coon, J.S., Coon, J.S., 1995. Comparison of Solutol HS 15,
818 Cremophor EL and novel ethoxylated fatty acid surfactants as multidrug resistance modification agents.
819 Int. J. Cancer 62, 436-442.
- 820 Chan, L.M.S., Lowes, S., Hirst, B.H., 2004. The ABCs of drug transport in intestine and liver: efflux
821 proteins limiting drug absorption and bioavailability. Elsevier B.V, Netherlands, pp. 25-51.
- 822 Chen, C.-j., Chin, J.E., Ueda, K., Clark, D.P., Pastan, I., Gottesman, M.M., Roninson, I.B., 1986.
823 Internal duplication and homology with bacterial transport proteins in the *mdr1* (P-glycoprotein) gene
824 from multidrug-resistant human cells. Cell 47, 381-389.
- 825 Chen, T.K., Li, Y., Li, C.W., Yi, X., Wang, R.B., Lee, S.M.Y., Zheng, Y., 2017. Pluronic P85/F68
826 Micelles of Baicalein Could Interfere with Mitochondria to Overcome MRP2-Mediated Efflux and
827 Offer Improved Anti-Parkinsonian Activity. Mol. Pharm. 14, 3331-3342.
- 828 Chen, Y., Sha, X., Zhang, W., Zhong, W., Fan, Z., Ren, Q., Chen, L., Fang, X., 2013. Pluronic mixed
829 micelles overcoming methotrexate multidrug resistance: in vitro and in vivo evaluation. Int J
830 Nanomedicine 8, 1463-1476.
- 831 Chen, Z., Shi, T., Zhang, L., Zhu, P., Deng, M., Huang, C., Hu, T., Jiang, L., Li, J., 2015. Mammalian
832 drug efflux transporters of the ATP binding cassette (ABC) family in multidrug resistance: A review of
833 the past decade. Cancer Lett. 370, 153-164.

- 834 Chiu, Y.-Y., Higaki, K., Neudeck, B.L., Barnett, J.L., Welage, L.S., Amidon, G.L., 2003. Human
835 Jejunal Permeability of Cyclosporin A: Influence of Surfactants on P-Glycoprotein Efflux in Caco-2
836 Cells. *Pharm. Res.* 20, 749-756.
- 837 Choi, Y.A., Yoon, Y.H., Choi, K., Kwon, M., Goo, S.H., Cha, J.-S., Choi, M.-K., Lee, H.S., Song, I.-
838 S., 2015. Enhanced Oral Bioavailability of Morin Administered in Mixed Micelle Formulation with
839 PluronicF127 and Tween80 in Rats. *Biol. Pharm. Bull.* 38, 208-217.
- 840 Christiansen, A., Backensfeld, T., Weitschies, W., 2010. Effects of non-ionic surfactants on in vitro
841 triglyceride digestion and their susceptibility to digestion by pancreatic enzymes. *Eur. J. Pharm. Sci.*
842 41, 376-382.
- 843 Colas, C., Ung, P.M.-U., Schlessinger, A., 2016. SLC Transporters: Structure, Function, and Drug
844 Discovery. *MedChemComm* 7, 1069-1081.
- 845 Collnot, E.-M., Baldes, C., Schaefer, U.F., Edgar, K.J., Wempe, M.F., Lehr, C.-M., 2010. Vitamin E
846 TPGS P-glycoprotein inhibition mechanism: influence on conformational flexibility, intracellular ATP
847 levels, and role of time and site of access. *Mol. Pharm.* 7, 642-651.
- 848 Collnot, E.-M., Baldes, C., Wempe, M.F., Kappl, R., Hüttermann, J., Hyatt, J.A., Edgar, K.J., Schaefer,
849 U.F., Lehr, C.-M., 2007. Mechanism of inhibition of P-glycoprotein mediated efflux by vitamin E
850 TPGS: influence on ATPase activity and membrane fluidity. *Mol. Pharm.* 4, 465-474.

- 851 Coon, J.S., Knudson, W., Clodfelter, K., Lu, B., Weinstein, R.S., 1991. Solutol HS 15, nontoxic
852 polyoxyethylene esters of 12-hydroxystearic acid, reverses multidrug resistance. *Cancer Res.* 51, 897-
853 902.
- 854 Cornaire, G., Woodley, J., Hermann, P., Cloarec, A., Arellano, C., Houin, G., 2004. Impact of
855 excipients on the absorption of P-glycoprotein substrates in vitro and in vivo. *Int. J. Pharm.* 278, 119-
856 131.
- 857 Cuiné, J.F., McEvoy, C.L., Charman, W.N., Pouton, C.W., Edwards, G.A., Benameur, H., Porter,
858 C.J.H., 2008. Evaluation of the Impact of Surfactant Digestion on the Bioavailability of Danazol after
859 Oral Administration of Lipidic Self-Emulsifying Formulations to Dogs. *J. Pharm. Sci.* 97, 995-1012.
- 860 Czogalla, A., 2009. Oral cyclosporine A - the current picture of its liposomal and other delivery
861 systems. *Cell. Mol. Biol. Lett.* 14, 139-152.
- 862 D'Souza, A.A., Shegokar, R., 2016. Polyethylene glycol (PEG): a versatile polymer for pharmaceutical
863 applications, England, pp. 1257-1275.
- 864 Dahan, A., Miller, J.M., Hoffman, A., Amidon, G.L., Amidon, G.E., 2010. The Solubility–Permeability
865 Interplay in Using Cyclodextrins as Pharmaceutical Solubilizers: Mechanistic Modeling and
866 Application to Progesterone. *J. Pharm. Sci.* 99, 2739-2749.

- 867 Dallas, S., Miller, D.S., Bendayan, R., 2006. Multidrug Resistance-Associated Proteins: Expression
868 and Function in the Central Nervous System. *Pharmacol. Rev.* 58, 140-161.
- 869 Dalton, W.S., Dalton, W.S., Crowley, J.J., Crowley, J.J., Salmon, S.S., Salmon, S.S., Grogan, T.M.,
870 Grogan, T.M., Laufman, L.R., Laufman, L.R., Weiss, G.R., Weiss, G.R., Bonnet, J.D., Bonnet, J.D.,
871 1995. A phase III randomized study of oral verapamil as a chemosensitizer to reverse drug resistance in
872 patients with refractory myeloma. A southwest oncology group study. *Cancer* 75, 815-820.
- 873 Devraj, R., Williams, H.D., Warren, D.B., Mohsin, K., Porter, C.J.H., Pouton, C.W., 2013. In vitro
874 assessment of drug-free and fenofibrate-containing lipid formulations using dispersion and digestion
875 testing gives detailed insights into the likely fate of formulations in the intestine. *Eur. J. Pharm. Sci.* 49,
876 748-760.
- 877 Dietrich, C.G., Geier, A., Oude Elferink, R.P.J., 2003. ABC of oral bioavailability: transporters as
878 gatekeepers in the gut. *Gut* 52, 1788-1795.
- 879 Dong, X., Mattingly, C.A., Tseng, M.T., Cho, M.J., Liu, Y., Adams, V.R., Mumper, R.J., 2009.
880 Doxorubicin and paclitaxel-loaded lipid-based nanoparticles overcome multidrug resistance by
881 inhibiting P-glycoprotein and depleting ATP. *Cancer Res.* 69, 3918-3926.
- 882 Doyle, L.A., Ross, D.D., 2003. Multidrug resistance mediated by the breast cancer resistance protein
883 BCRP (ABCG2). *Oncogene* 22, 7340-7358.

- 884 Drewe, J., Meier, R., Vonderscher, J., Kiss, D., Posanski, U., Kissel, T., Gyr, K., 1992. Enhancement of
885 the oral absorption of cyclosporin in man. *Br. J. Clin. Pharmacol.* 34, 60-64.
- 886 Drozdik, M., Busch, D., Lapczuk, J., Müller, J., Ostrowski, M., Kurzawski, M., Oswald, S., 2019.
887 Protein Abundance of Clinically Relevant Drug Transporters in the Human Liver and Intestine: A
888 Comparative Analysis in Paired Tissue Specimens. *Clin. Pharmacol. Ther.*
- 889 Drozdik, M., Groer, C., Penski, J., Lapczuk, J., Ostrowski, M., Lai, Y., Prasad, B., Unadkat, J.D.,
890 Siegmund, W., Oswald, S., 2014. Protein abundance of clinically relevant multidrug transporters along
891 the entire length of the human intestine. *Mol. Pharm.* 11, 3547-3555.
- 892 Drugbank, 2019. P-glycoprotein substrates.
893 [https://www.drugbank.ca/unearth/q?utf8=%E2%9C%93&searcher=drugs&query=P-](https://www.drugbank.ca/unearth/q?utf8=%E2%9C%93&searcher=drugs&query=P-glycoprotein+substrate)
894 [glycoprotein+substrate](https://www.drugbank.ca/unearth/q?utf8=%E2%9C%93&searcher=drugs&query=P-glycoprotein+substrate). (accessed 01.003.2019).
- 895 Eckford, P.D.W., Sharom, F.J., 2005. The reconstituted P-glycoprotein multidrug transporter is a
896 flippase for glucosylceramide and other simple glycosphingolipids. *Biochem. J.* 389, 517-526.
- 897 Eley, J.G., Triumalashetty, P., 2001. In vitro assessment of alkylglycosides as permeability enhancers.
898 *AAPS PharmSciTech* 2, 81-87.
- 899 Engel, A., Oswald, S., Siegmund, W., Keiser, M., 2012. Pharmaceutical excipients influence the
900 function of human uptake transporting proteins. *Mol. Pharm.* 9, 2577-2581.

- Englund, G., Jacobson, A., Rorsman, F., Artursson, P., Kindmark, A., Rönnblom, A., Medicinska, f.,
Medicinska och farmaceutiska, v., Uppsala, u., Institutionen för medicinska, v., Institutionen för, f.,
Farmaceutiska, f., 2007. Efflux transporters in ulcerative colitis: decreased expression of BCRP
(ABCG2) and Pgp (ABCB1). *Inflamm. Bowel Dis.* 13, 291-297.
- Erickson, R.H., Gum, J.R., Lindstrom, M.M., McKean, D., Kim, Y.S., 1995. Regional Expression and
Dietary Regulation of Rat Small Intestinal Peptide and Amino Acid Transporter mRNAs. *Biochem.*
Biophys. Res. Commun. 216, 249-257.
- Evers, R., Kool, M., Smith, A.J., van Deemter, L., de Haas, M., Borst, P., 2000. Inhibitory effect of the
reversal agents V-104, GF120918 and Pluronic L61 on MDR1 Pgp-, MRP1- and MRP2-mediated
transport. *Br. J. Cancer* 83, 366-374.
- Fricker, G., Drewe, J., Huwyler, J., Gutmann, H., Beglinger, C., 1996. Relevance of p-glycoprotein for
the enteral absorption of cyclosporin A: in vitro-in vivo correlation. *Br. J. Pharmacol.* 118, 1841-1847.
- Fromm, M.F., 2004. Importance of P-glycoprotein at blood–tissue barriers. *Trends Pharmacol. Sci.* 25,
423-429.
- Föger, F., Hoyer, H., Kafedjiiski, K., Thaurer, M., Bernkop-Schnürch, A., 2006. In vivo comparison of
various polymeric and low molecular mass inhibitors of intestinal P-glycoprotein. *Biomaterials* 27,
5855-5860.

- 918 Giacomini, K.M., Huang, S.-M., Tweedie, D.J., Benet, L.Z., Brouwer, K.L.R., Chu, X., Dahlin, A.,
919 Evers, R., Fischer, V., Hillgren, K.M., Hoffmaster, K.A., Ishikawa, T., Keppler, D., Kim, R.B., Lee,
920 C.A., Niemi, M., Polli, J.W., Sugiyama, Y., Swaan, P.W., Ware, J.A., Wright, S.H., Yee, S.W., Zamek-
921 Gliszczynski, M.J., Zhang, L., International Transporter, C., 2010. Membrane transporters in drug
922 development. *Nat. Rev. Drug Discov.* 9, 215-236.
- 923 Gramatté, T., Oertel, R., 1999. Intestinal secretion of intravenous talinolol is inhibited by luminal
924 R-verapamil. *Clin. Pharmacol. Ther.* 66, 239-245.
- 925 Greenberg, P.L., Lee, S.J., Advani, R., Tallman, M.S., Sikic, B.I., Letendre, L., Dugan, K., Lum, B.,
926 Chin, D.L., Dewald, G., Paietta, E., Bennett, J.M., Rowe, J.M., 2004. Mitoxantrone, Etoposide, and
927 Cytarabine With or Without Valspodar in Patients With Relapsed or Refractory Acute Myeloid
928 Leukemia and High-Risk Myelodysplastic Syndrome: A Phase III Trial (E2995). *J. Clin. Oncol.* 22,
929 1078-1086.
- 930 Gröer, C., Brück, S., Lai, Y., Paulick, A., Busemann, A., Heidecke, C.D., Siegmund, W., Oswald, S.,
931 2013. LC-MS/MS-based quantification of clinically relevant intestinal uptake and efflux transporter
932 proteins. *J. Pharm. Biomed. Anal.* 85, 253-261.
- 933 Guan, Y., Huang, J., Zuo, L., Xu, J., Si, L., Qiu, J., Li, G., 2011. Effect of pluronic P123 and F127
934 block copolymer on P-glycoprotein transport and CYP3A metabolism. *Arch. Pharm. Res.* 34, 1719-
935 1728.

- 936 Gupta, A., Dai, Y., Vethanayagam, R.R., Hebert, M.F., Thummel, K.E., Unadkat, J.D., Ross, D.D.,
937 Mao, Q., 2006. Cyclosporin A, tacrolimus and sirolimus are potent inhibitors of the human breast
938 cancer resistance protein (ABCG2) and reverse resistance to mitoxantrone and topotecan. *Cancer*
939 *Chemother. Pharmacol.* 58, 374-383.
- 940 Gupta, A., Zhang, Y., Unadkat, J.D., Mao, Q., 2004. HIV Protease Inhibitors Are Inhibitors but Not
941 Substrates of the Human Breast Cancer Resistance Protein (BCRP/ABCG2). *J. Pharmacol. Exp. Ther.*
942 310, 334-341.
- 943 Gupta, E., Safa, A.R., Wang, X., Ratain, M.J., 1996. Pharmacokinetic modulation of irinotecan and
944 metabolites by cyclosporin A. *Cancer Res.* 56, 1309-1314.
- 945 Gurjar, R., Chan, C., Curley, P., Sharp, J., Chiong, J., Rannard, S., Siccardi, M., Owen, A., 2018.
946 Inhibitory effects of commonly used excipients on P-glycoprotein in vitro. *Mol. Pharm.* 15, 4835-4842.
- 947 Hanke, U., May, K., Rozehnal, V., Nagel, S., Siegmund, W., Weitschies, W., 2010. Commonly used
948 nonionic surfactants interact differently with the human efflux transporters ABCB1 (p-glycoprotein)
949 and ABCC2 (MRP2). *Eur. J. Pharm. Biopharm.* 76, 260-268.
- 950 Harwood, M.D., Achour, B., Russell, M.R., Carlson, G.L., Warhurst, G., Rostami-Hodjegan, A., 2015.
951 Application of an LC–MS/MS method for the simultaneous quantification of human intestinal
952 transporter proteins absolute abundance using a QconCAT technique. *J. Pharm. Biomed. Anal.* 110, 27-
953 33.

- 954 Harwood, M.D., Neuhoﬀ, S., Carlson, G.L., Warhurst, G., Rostami-Hodjegan, A., 2013. Absolute
955 abundance and function of intestinal drug transporters: a prerequisite for fully mechanistic in vitro–in
956 vivo extrapolation of oral drug absorption. *Biopharm. Drug Disposition* 34, 2-28.
- 957 Hediger, M.A., 2004. The ABCs of solute carriers: physiological, pathological and therapeutic
958 implications of human membrane transport proteins. *Pflügers Arch.* 447, 465-468.
- 959 Hediger, M.A., Cléménçon, B., Burrier, R.E., Bruford, E.A., 2013. The ABCs of membrane
960 transporters in health and disease (SLC series): Introduction. *Mol. Aspects Med.* 34, 95-107.
- 961 Higgins, C.F., Gottesman, M.M., 1992. Is the multidrug transporter a ﬂippase? *Trends Biochem. Sci.*
962 17, 18-21.
- 963 Hodaei, D., Baradaran, B., Valizadeh, H., Zakeri-Milani, P., 2015. Effects of polyethylene glycols on
964 intestinal eﬄux pump expression and activity in Caco-2 cells. *BJPS* 51, 745-753.
- 965 Houghton, P.J., Germain, G.S., Harwood, F.C., Schuetz, J.D., Stewart, C.F., Buchdunger, E., Traxler,
966 P., 2004. Imatinib mesylate is a potent inhibitor of the ABCG2 (BCRP) transporter and reverses
967 resistance to topotecan and SN-38 in vitro. *Cancer Res.* 64, 2333-2337.
- 968 Hugger, E.D., Novak, B.L., Burton, P.S., Audus, K.L., Borchardt, R.T., 2002. A comparison of
969 commonly used polyethoxylated pharmaceutical excipients on their ability to inhibit P-glycoprotein
970 activity in vitro. *J. Pharm. Sci.* 91, 1991-2002.

- 971 HUGO Gene Nomenclature Committee, 2019. ATP binding cassette transporters (ABC).
 972 <https://www.genenames.org/cgi-bin/genefamilies/set/417>. Solute carriers (SLC).
 973 <https://www.genenames.org/cgi-bin/genefamilies/set/752>., (accessed 12.01.2019).
- 974 Hyafil, F., Vergely, C., Vignaud, P.D., Grand-Perret, T., 1993. In vitro and in vivo reversal of
 975 multidrug resistance by GF120918, an acridonecarboxamide derivative. *Cancer Res.* 53, 4595-4602.
- 976 Italia, J.L., Bhardwaj, V., Ravi Kumar, M.N.V., 2006. Disease, destination, dose and delivery aspects
 977 of ciclosporin: the state of the art. *Drug Discov Today* 11, 846-854.
- 978 Jedlitschky, G., Hoffmann, U., Kroemer, H.K., 2006. Structure and function of the MRP2 (ABCC2)
 979 protein and its role in drug disposition. *Expert Opin. Drug Metab. Toxicol.* 2, 351-366.
- 980 Johnson, B.M., Charman, W.N., Porter, C.J.H., 2002. An in vitro examination of the impact of
 981 polyethylene glycol 400, pluronic P85, and vitamin E d- α -tocopheryl polyethylene glycol 1000
 982 succinate on P-glycoprotein efflux and enterocyte-based metabolism in excised rat intestine. *AAPS*
 983 *PharmSci* 4, 193-205.
- 984 Johnson, Z.L., Chen, J., 2017. Structural Basis of Substrate Recognition by the Multidrug Resistance
 985 Protein MRP1. *Cell* 168, 1075-1085.e1079.
- 986 Jonker, J.W., Smit, J.W., Brinkhuis, R.F., Maliepaard, M., 2000. Role of breast cancer resistance
 987 protein in the bioavailability and fetal penetration of topotecan. *J. Natl. Cancer Inst.* 92, 1651.

- 988 Juliano, R.L., Ling, V., 1976. A surface glycoprotein modulating drug permeability in Chinese hamster
989 ovary cell mutants. *Biochim. Biophys. Acta* 455, 152-162.
- 990 Keller, R.P., Altermatt, H.J., Nooter, K., Poschmann, G., Laissue, J.A., Bollinger, P., Hiestand, P.C.,
991 1992. SDZ PSC 833, a non-immunosuppressive cyclosporine: its potency in overcoming P-
992 glycoprotein-mediated multidrug resistance of murine leukemia. *Int. J. Cancer* 50, 593-597.
- 993 Kim, Y., Chen, J., 2018. Molecular structure of human P-glycoprotein in the ATP-bound, outward-
994 facing conformation. *Science* 359, 915-919.
- 995 Kiss, L., Hellinger, É., Pilbat, A.M., Kittel, Á., Török, Z., Füredi, A., Szakács, G., Veszeka, S., Sipos,
996 P., Ózsvári, B., Puskás, L.G., Vastag, M., Szabó-Révész, P., Deli, M.A., 2014. Sucrose Esters Increase
997 Drug Penetration, But Do Not Inhibit P-Glycoprotein in Caco-2 Intestinal Epithelial Cells. *J. Pharm.*
998 *Sci.* 103, 3107-3119.
- 999 Koepsell, H., Lips, K., Volk, C., 2007. Polyspecific organic cation transporters: structure, function,
1000 physiological roles, and biopharmaceutical implications. *Pharm. Res.* 24, 1227-1251.
- 1001 Kool, M., de Haas, M., Scheffer, G.L., Scheper, R.J., van Eijk, M.J., Juijn, J.A., Baas, F., Borst, P.,
1002 1997. Analysis of expression of cMOAT (MRP2), MRP3, MRP4, and MRP5, homologues of the
1003 multidrug resistance-associated protein gene (MRP1), in human cancer cell lines. *Cancer Res.* 57,
1004 3537-3547.

- 1005 Kool, M., Marcel van der, L., Haas, M.d., Scheffer, G.L., Vree, J.M.L.D., Smith, A.J., Jansen, G.,
1006 Peters, G.J., Ponne, N., Scheper, R.J., Ronald, P.J.O.E., Baas, F., Borst, P., 1999. MRP3, an Organic
1007 Anion Transporter Able to Transport Anti-Cancer Drugs. *Proc. Natl. Acad. Sci. U. S. A.* 96, 6914-
1008 6919.
- 1009 Kumar, H., Finer-Moore, J.S., Jiang, X., Smirnova, I., Kasho, V., Pardon, E., Steyaert, J., Kaback,
1010 H.R., Stroud, R.M., 2018. Crystal Structure of a ligand-bound LacY–Nanobody Complex. *Proc. Natl.*
1011 *Acad. Sci. U. S. A.* 115, 8769.
- 1012 Kumar, H., Kasho, V., Smirnova, I., Finer-Moore, J.S., Kaback, H.R., Stroud, R.M., 2014. Structure of
1013 sugar-bound LacY. *Proc. Natl. Acad. Sci.* 111, 1784.
- 1014 Lamprecht, A., Benoit, J.-P., 2006. Etoposide nanocarriers suppress glioma cell growth by intracellular
1015 drug delivery and simultaneous P-glycoprotein inhibition. *J. Control. Release* 112, 208-213.
- 1016 Lehnert, M., Mross, K., Schueller, J., Thuerlimann, B., Kroeger, N., Kupper, H., 1998. Phase II trial of
1017 dexverapamil and epirubicin in patients with non-responsive metastatic breast cancer. *Br. J. Cancer* 77,
1018 1155-1163.
- 1019 Li-Blatter, X., Beck, A., Seelig, A., 2012. P-glycoprotein-ATPase modulation: the molecular
1020 mechanisms. *Biophys. J.* 102, 1383-1393.

- 1021 Li-Blatter, X., Nervi, P., Seelig, A., 2009. Detergents as intrinsic P-glycoprotein substrates and
1022 inhibitors. *BBA - Biomembranes* 1788, 2335-2344.
- 1023 Li-Blatter, X., Seelig, A., 2010. Exploring the P-glycoprotein binding cavity with polyoxyethylene
1024 alkyl ethers. *Biophys. J.* 99, 3589-3598.
- 1025 Li, L., Yi, T., Lam, C.W.-k., 2013a. Interactions between human multidrug resistance related protein
1026 (MRP2; ABCC2) and excipients commonly used in self-emulsifying drug delivery systems (SEDDES).
1027 *Int. J. Pharm.* 447, 192-198.
- 1028 Li, L., Yi, T., Lam, C.W.-k., 2013b. Interactions between human multidrug resistance related protein
1029 (MRP2; ABCC2) and excipients commonly used in self-emulsifying drug delivery systems (SEDDES).
1030 *Int. J. Pharm.* 447, 192-198.
- 1031 Li, L., Yi, T., Lam, C.W.-K., 2014. Inhibition of human efflux transporter ABCC2 (MRP2) by self-
1032 emulsifying drug delivery system: influences of concentration and combination of excipients. *J. Pharm.*
1033 *Pharm. Sci.* 17, 447.
- 1034 Lin, J.H., Yamazaki, M., 2003. Role of P-glycoprotein in pharmacokinetics. *Clin. Pharmacokinet.* 42,
1035 59-98.

- 1036 Lin, Y., Shen, Q., Katsumi, H., Okada, N., Fujita, T., Jiang, X., Yamamoto, A., 2007. Effects of
1037 Labrasol and Other Pharmaceutical Excipients on the Intestinal Transport and Absorption of
1038 Rhodamine123, a P-Glycoprotein Substrate, in Rats. *Biol. Pharm. Bull.* 30, 1301-1307.
- 1039 Lloret-Linares, C., Miyauchi, E., Luo, H., Labat, L., Bouilliot, J.-L., Poitou, C., Oppert, J.-M.,
1040 Laplanche, J.-L., Mouly, S., Scherrmann, J.-M., Uchida, Y., Tachikawa, M., Terasaki, T., Bergmann,
1041 J.-F., Declèves, X., 2016. Oral Morphine Pharmacokinetic in Obesity: The Role of P-Glycoprotein,
1042 MRP2, MRP3, UGT2B7, and CYP3A4 Jejunal Contents and Obesity-Associated Biomarkers. *Mol.*
1043 *Pharm.* 13, 766-773.
- 1044 Lo, Y.-l., 2003. Relationships between the hydrophilic–lipophilic balance values of pharmaceutical
1045 excipients and their multidrug resistance modulating effect in Caco-2 cells and rat intestines. *J.*
1046 *Controlled Release* 90, 37-48.
- 1047 Lown, K.S., 1997. Role of intestinal P-glycoprotein (mdr1) in interpatient variation in the oral
1048 bioavailability of cyclosporine. *Clin. Pharmacol. Ther.* 62, 248-260.
- 1049 Ma, L., Wei, Y., Zhou, Y., Ma, X., Wu, X.a., 2011. Effects of Pluronic F68 and Labrasol on the
1050 intestinal absorption and pharmacokinetics of rifampicin in rats. *Arch. Pharm. Res.* 34, 1939-1943.
- 1051 Maliepaard, M., Margôt, A.v.G., Tohgo, A., Hausheer, F.H., Robert, C.A.M.v.W., Laurina, A.d.J.,
1052 Pluim, D., Beijnen, J.H., Jan, H.M.S., 2001a. Circumvention of Breast Cancer Resistance Protein

- 1053 (BCRP)-mediated Resistance to Camptothecins in Vitro Using Non-Substrate Drugs or the BCRP
1054 Inhibitor GF120918. Clin. Cancer. Res. 7, 935-941.
- 1055 Maliepaard, M., Scheffer, G.L., Faneyte, I.F., van Gastelen, M.A., Pijnenborg, A.C., Schinkel, A.H.,
1056 van de Vijver, M.J., Scheper, R.J., Schellens, J.H., 2001b. Subcellular localization and distribution of
1057 the breast cancer resistance protein transporter in normal human tissues. Cancer Res. 61, 3458-3464.
- 1058 Maliepaard, M., van Gastelen, M.A., de Jong, L.A., Pluim, D., van Waardenburg, R., Ruevekamp-
1059 Helmers, M.C., Floom, B.G.J., Schellens, J.H.M., 1999. Overexpression of the BCRP/MXR/ABCP
1060 Gene in a Topotecan-selected Ovarian Tumor Cell Line. Cancer Res. 59, 4559-4563.
- 1061 Mao, Q., 2005. Role of the breast cancer resistance protein (ABCG2) in drug transport. The AAPS
1062 Journal 7, E118-E133.
- 1063 Mao, Q., Unadkat, J.D., 2015. Role of the Breast Cancer Resistance Protein (BCRP/ABCG2) in Drug
1064 Transport—an Update. The AAPS Journal 17, 65-82.
- 1065 Matsson, P., Pedersen, J.M., Norinder, U., Bergström, C.A.S., Artursson, P., Medicinska och
1066 farmaceutiska, v., Uppsala, u., Institutionen för, f., Farmaceutiska, f., 2009. Identification of novel
1067 specific and general inhibitors of the three major human ATP-binding cassette transporters P-gp, BCRP
1068 and MRP2 among registered drugs. Pharm. Res. 26, 1816-1831.

- 1069 Mechetner, E., Kyshtoobayeva, A., Zonis, S., Kim, H., Stroup, R., Garcia, R., Parker, R.J., Fruehauf,
1070 J.P., 1998. Levels of multidrug resistance (MDR1) P-glycoprotein expression by human breast cancer
1071 correlate with in vitro resistance to taxol and doxorubicin. Clin. Cancer. Res. 4, 389.
- 1072 Miller, J.M., Beig, A., Krieg, B.J., Carr, R.A., Borchardt, T.B., Amidon, G.E., Amidon, G.L., Dahan,
1073 A., 2011. The solubility-permeability interplay: mechanistic modeling and predictive application of the
1074 impact of micellar solubilization on intestinal permeation. Mol. Pharm. 8, 1848-1856.
- 1075 Miyauchi, E., Tachikawa, M., Declèves, X., Uchida, Y., Bouillot, J.-L., Poitou, C., Oppert, J.-M.,
1076 Mouly, S., Bergmann, J.-F., Terasaki, T., Scherrmann, J.-M., Lloret-Linares, C., 2016. Quantitative
1077 Atlas of Cytochrome P450, UDP-Glucuronosyltransferase, and Transporter Proteins in Jejunum of
1078 Morbidly Obese Subjects. Mol. Pharm. 13, 2631-2640.
- 1079 Mohsin, K., 2012. Design of Lipid-Based Formulations for Oral Administration of Poorly Water-
1080 Soluble Drug Fenofibrate: Effects of Digestion. AAPS PharmSciTech 13, 637-646.
- 1081 Mross, K., Kröger, N., Herbst, K., Gastl, G., Hossfeld, D., 1999. Alteration in Epirubicin
1082 Pharmacokinetics and Metabolism by Dexverapamil: Results from a Phase II Study in Patients with
1083 Metastatic Breast Cancer. Oncol Res Treat 22, 35-40.
- 1084 Nerurkar, M.M., Burton, P.S., Borchardt, R.T., 1996. The use of surfactants to enhance the
1085 permeability of peptides through Caco-2 cells by inhibition of an apically polarized efflux system.
1086 Pharm. Res. 13, 528-534.

- 1087 Nielsen, C.U., Abdulhussein, A.A., Colak, D., Holm, R., 2016. Polysorbate 20 increases oral
1088 absorption of digoxin in wild-type Sprague Dawley rats, but not in *mdr1a*(-/-) Sprague Dawley rats. *Int.*
1089 *J. Pharm.* 513, 78-87.
- 1090 Nies, A.T., Keppler, D., 2007. The apical conjugate efflux pump ABCC2 (MRP2). *Pflugers Arch.* 453,
1091 643-659.
- 1092 O'Leary, T.J., Ross, P.D., Lieber, M.R., Levin, I.W., 1986. Effects of cyclosporine A on
1093 biomembranes. Vibrational spectroscopic, calorimetric and hemolysis studies. *Biophys. J.* 49, 795-801.
- 1094 Oswald, S., Haenisch, S., Fricke, C., Sudhop, T., Remmler, C., Giessmann, T., Jedlitschky, G., Adam,
1095 U., Dazert, E., Warzok, R., Wacke, W., Cascorbi, I., Kroemer, H.K., Weitschies, W., Bergmann, K.,
1096 Siegmund, W., 2006. Intestinal expression of P-glycoprotein (ABCB1), multidrug resistance associated
1097 protein 2 (ABCC2), and uridine diphosphate-glucuronosyltransferase 1A1 predicts the disposition and
1098 modulates the effects of the cholesterol absorption inhibitor ezetimibe in humans. *Clin. Pharmacol.*
1099 *Ther.* 79, 206-217.
- 1100 Otter, M., Oswald, S., Siegmund, W., Keiser, M., 2017. Effects of frequently used pharmaceutical
1101 excipients on the organic cation transporters 1–3 and peptide transporters 1/2 stably expressed in
1102 MDCKII cells. *Eur. J. Pharm. Biopharm.* 112, 187-195.

- 1103 Patil, S., Choudhary, B., Rathore, A., Roy, K., Mahadik, K., 2015. Enhanced oral bioavailability and
1104 anticancer activity of novel curcumin loaded mixed micelles in human lung cancer cells.
1105 *Phytomedicine* 22, 1103-1111.
- 1106 Petersen, S.B., Nolan, G., Maher, S., Rahbek, U.L., Guldbrandt, M., Brayden, D.J., 2012. Evaluation of
1107 alkylmaltosides as intestinal permeation enhancers: comparison between rat intestinal mucosal sheets
1108 and Caco-2 monolayers. *Eur. J. Pharm. Sci.* 47, 701-712.
- 1109 Planting, A.S.T., Sonneveld, P., van der Gaast, A., Sparreboom, A., van der Burg, M.E.L., Luyten,
1110 G.P.M., de Leeuw, K., de Boer-Dennert, M., Wissel, P.S., Jewell, R.C., Paul, E.M., Purvis Jr, N.B.,
1111 Verweij, J., 2005. A phase I and pharmacologic study of the MDR converter GF120918 in combination
1112 with doxorubicin in patients with advanced solid tumors. *Cancer Chemother. Pharmacol.* 55, 91-99.
- 1113 Postolache, P., Petrescu, O., Dorneanu, V., Zanini, A., 2002. Cyclosporine bioavailability of two
1114 physically different oral formulations. *Eur. Rev. Med. Pharmacol. Sci.* 6, 127-131.
- 1115 Pouton, C.W., 2006. Formulation of poorly water-soluble drugs for oral administration:
1116 Physicochemical and physiological issues and the lipid formulation classification system. *Eur. J.*
1117 *Pharm. Sci.* 29, 278-287.
- 1118 Radestock, S., Forrest, L.R., 2011. The Alternating-Access Mechanism of MFS Transporters Arises
1119 from Inverted-Topology Repeats. *J. Mol. Biol.* 407, 698-715.

- 1120 Rao, V.V., Dahlheimer, J.L., Bardgett, M.E., Snyder, A.Z., Finch, R.A., Sartorelli, A.C., Piwnica-
1121 Worms, D., 1999. Choroid Plexus Epithelial Expression of MDR1 P Glycoprotein and Multidrug
1122 Resistance-Associated Protein Contribute to the Blood-Cerebrospinal-Fluid Drug-Permeability Barrier.
1123 Proc. Natl. Acad. Sci. U. S. A. 96, 3900-3905.
- 1124 Raviv, Y., Pollard, H.B., Bruggemann, E.P., Pastan, I., Gottesman, M.M., 1990. Photosensitized
1125 labeling of a functional multidrug transporter in living drug-resistant tumor cells. J. Biol. Chem. 265,
1126 3975-3980.
- 1127 Rege, B.D., Kao, J.P., Polli, J.E., 2002. Effects of nonionic surfactants on membrane transporters in
1128 Caco-2 cell monolayers. Eur. J. Pharm. Sci. 16, 237-246.
- 1129 Regev, R., Assaraf, Y.G., Eytan, G.D., 1999. Membrane fluidization by ether, other anesthetics, and
1130 certain agents abolishes P-glycoprotein ATPase activity and modulates efflux from multidrug-resistant
1131 cells. Eur. J. Biochem. 259, 18-24.
- 1132 Ries, F., Dicato, M., 1991. Treatment of advanced and refractory breast cancer with doxorubicin,
1133 vincristine and continuous infusion of verapamil. a phase I-II clinical trial. Med. Oncol. Tumor
1134 Pharmacother. 8, 39-43.
- 1135 Romsicki, Y., Sharom, F.J., 2001. Phospholipid flippase activity of the reconstituted P-glycoprotein
1136 multidrug transporter. Biochemistry 40, 6937-6947.

- 1137 Rosenberg, M.F., Bikadi, Z., Chan, J., Liu, X., Ni, Z., Cai, X., Ford, R.C., Mao, Q., 2010. The Human
1138 Breast Cancer Resistance Protein (BCRP/ABCG2) Shows Conformational Changes with Mitoxantrone.
1139 Structure 18, 482-493.
- 1140 Rosenberg, M.F., Bikadi, Z., Hazai, E., Starborg, T., Kelley, L., Chayen, N.E., Ford, R.C., Mao, Q.,
1141 2015. Three-dimensional structure of the human breast cancer resistance protein (BCRP/ABCG2) in an
1142 inward-facing conformation. Acta Crystallographica Section D 71, 1725-1735.
- 1143 Sachs-Barrable, K., Thamboo, A., Lee, S.D., Wasan, K.M., 2007. Lipid excipients Peceol and Gelucire
1144 44/14 decrease P-glycoprotein mediated efflux of rhodamine 123 partially due to modifying P-
1145 glycoprotein protein expression within Caco-2 cells. J. Pharm. Pharm. Sci. 10, 319-331.
- 1146 Sauna, Z.E., Ambudkar, S.V., 2001. Characterization of the catalytic cycle of ATP hydrolysis by
1147 human P-glycoprotein. The two ATP hydrolysis events in a single catalytic cycle are kinetically similar
1148 but affect different functional outcomes. J. Biol. Chem. 276, 11653-11661.
- 1149 Savla, R., Browne, J., Plassat, V., Wasan, K.M., Wasan, E.K., 2017. Review and analysis of FDA
1150 approved drugs using lipid-based formulations. Drug Dev. Ind. Pharm. 43, 1743-1758.
- 1151 Sawangrat, K., Morishita, M., Kusamori, K., Katsumi, H., Sakane, T., Yamamoto, A., 2018a. Effects of
1152 Various Pharmaceutical Excipients on the Intestinal Transport and Absorption of Sulfasalazine, a
1153 Typical Substrate of Breast Cancer Resistance Protein Transporter. J. Pharm. Sci. 107, 2946-2956.

- 1154 Sawangrat, K., Yamashita, S., Tanaka, A., Morishita, M., Kusamori, K., Katsumi, H., Sakane, T.,
1155 Yamamoto, A., 2018b. Modulation of intestinal transport and absorption of topotecan, a BCRP
1156 substrate by various pharmaceutical excipients and their inhibitory mechanisms of BCRP transporter. *J.*
1157 *Pharm. Sci.*
- 1158 Seelig, A., 1998. A general pattern for substrate recognition by P-glycoprotein. *Eur. J. Biochem.* 251,
1159 252-261.
- 1160 Seelig, A., Gerebtzoff, G., 2006. Enhancement of drug absorption by noncharged detergents through
1161 membrane and P-glycoprotein binding. *Expert Opin. Drug Metab. Toxicol.* 2, 733-752.
- 1162 Shaik, N., Pan, G., Elmquist, W.F., 2008. Interactions of pluronic block copolymers on P-gp efflux
1163 activity: Experience with HIV-1 protease inhibitors. *J. Pharm. Sci.* 97, 5421-5433.
- 1164 Shen, Q., Li, W., Lin, Y., Katsumi, H., Okada, N., Sakane, T., Fujita, T., Yamamoto, A., 2008.
1165 Modulating effect of polyethylene glycol on the intestinal transport and absorption of prednisolone,
1166 methylprednisolone and quinidine in rats by in-vitro and in-situ absorption studies. *J. Pharm.*
1167 *Pharmacol.* 60, 1633-1641.
- 1168 Shen, Q., Lin, Y., Handa, T., Doi, M., Sugie, M., Wakayama, K., Okada, N., Fujita, T., Yamamoto, A.,
1169 2006. Modulation of intestinal P-glycoprotein function by polyethylene glycols and their derivatives by
1170 in vitro transport and in situ absorption studies. *Int. J. Pharm.* 313, 49-56.

- 1171 Shimomura, H., Nogami, R., Shigeno, A., Shimada, S., Aoyama, T., 2016. Influence of Food on
1172 Rifampicin Pharmacokinetics in Rats. *Biol. Pharm. Bull.* 39, 49-53.
- 1173 Shono, Y., Nishihara, H., Matsuda, Y., Furukawa, S., Okada, N., Fujita, T., Yamamoto, A., 2004.
1174 Modulation of Intestinal P-Glycoprotein Function by Cremophor EL and Other Surfactants by an In
1175 Vitro Diffusion Chamber Method Using the Isolated Rat Intestinal Membranes. *J. Pharm. Sci.* 93, 877-
1176 885.
- 1177 Soodvilai, S., Soodvilai, S., Chatsudthipong, V., Ngawhirunpat, T., Rojanarata, T., Opanasopit, P.,
1178 2017. Interaction of pharmaceutical excipients with organic cation transporters. *Int. J. Pharm.* 520, 14-
1179 20.
- 1180 Sparreboom, A., Planting, A.S., Jewell, R.C., Loos, W., Nooter, K., Chandler, L., Paul, E., Wissel, P.,
1181 Verweij, J., 1999. Clinical pharmacokinetics of doxorubicin in combination with GF120918, a potent
1182 inhibitor of MDR1 P-glycoprotein. *Anti-Cancer Drugs* 10, 719-728.
- 1183 Steffansen, B., Nielsen, C.U., Brodin, B., Eriksson, A.H., Andersen, R., Frokjaer, S., 2004. Intestinal
1184 solute carriers: an overview of trends and strategies for improving oral drug absorption. Elsevier B.V,
1185 Netherlands, pp. 3-16.
- 1186 Steffansen, B., Pedersen, M.D.L., Laghmoch, A.M., Nielsen, C.U., 2017. SGLT1-Mediated Transport
1187 in Caco-2 Cells Is Highly Dependent on Cell Bank Origin. *J. Pharm. Sci.* 106, 2664-2670.

- 1188 Su, L., Cheng, C.Y., Mruk, D.D., 2009. Drug transporter, P-glycoprotein (MDR1), is an integrated
1189 component of the mammalian blood–testis barrier. *Int. J. Biochem. Cell Biol.* 41, 2578-2587.
- 1190 Tsuruo, T., Tsuruo, T., Iida, H., Iida, H., Tsukagoshi, S., Tsukagoshi, S., Sakurai, Y., Sakurai, Y.,
1191 1981. Overcoming of vincristine resistance in P388 leukemia in vivo and in vitro through enhanced
1192 cytotoxicity of vincristine and vinblastine by verapamil. *Cancer Res.* 41, 1967-1972.
- 1193 Uchida, Y., Ohtsuki, S., Kamiie, J., Ohmine, K., Iwase, R., Terasaki, T., 2015. Quantitative targeted
1194 absolute proteomics for 28 human transporters in plasma membrane of Caco-2 cell monolayer cultured
1195 for 2, 3, and 4 weeks. *Drug Metab. Pharmacokinet.* 30, 205-208.
- 1196 van Helvoort, A., Smith, A.J., Sprong, H., Fritzsche, I., Schinkel, A.H., Borst, P., van Meer, G., 1996.
1197 MDR1 P-Glycoprotein Is a Lipid Translocase of Broad Specificity, While MDR3 P-Glycoprotein
1198 Specifically Translocates Phosphatidylcholine. *Cell* 87, 507-517.
- 1199 van Asperen, J., van Tellingen, O., Sparreboom, A., Schinkel, A.H., Borst, P., Nooijen, W.J., Beijnen,
1200 J.H., 1997. Enhanced oral bioavailability of paclitaxel in mice treated with the P-glycoprotein blocker
1201 SDZ PSC 833. *Br. J. Cancer* 76, 1181-1183.
- 1202 Varma, M.V.S., Panchagnula, R., 2005. Enhanced oral paclitaxel absorption with vitamin E-TPGS:
1203 Effect on solubility and permeability in vitro, in situ and in vivo. *Eur. J. Pharm. Sci.* 25, 445-453.

- 1204 Wang, H., Lee, E.-W., Cai, X., Ni, Z., Zhou, L., Mao, Q., 2008. Membrane topology of the human
1205 breast cancer resistance protein (BCRP/ABCG2) determined by epitope insertion and
1206 immunofluorescence. *Biochemistry* 47, 13778-13787.
- 1207 Warner, E., Hedley, D., Andrulis, I., Myers, R., Trudeau, M., Warr, D., Pritchard, K.I., Blackstein, M.,
1208 Goss, P.E., Franssen, E., Roche, K., Knight, S., Webster, S., Fraser, R.A., Oldfield, S., Hill, W., Kates,
1209 R., 1998. Phase II study of dexverapamil plus anthracycline in patients with metastatic breast cancer
1210 who have progressed on the same anthracycline regimen. *Clin. Cancer. Res.* 4, 1451-1457.
- 1211 Wegler, C., Gaugaz, F.Z., Andersson, T.B., Wiśniewski, J.R., Busch, D., Gröer, C., Oswald, S., Norén,
1212 A., Weiss, F., Hammer, H.S., Joos, T.O., Poetz, O., Achour, B., Rostami-Hodjegan, A., van de Steeg,
1213 E., Wortelboer, H.M., Artursson, P., Medicinska, f., Medicinska och farmaceutiska, v.,
1214 Gastrointestinalkirurgi, Farmaceutiska, f., Uppsala, u., Institutionen för kirurgiska, v., Institutionen för,
1215 f., 2017. Variability in Mass Spectrometry-based Quantification of Clinically Relevant Drug
1216 Transporters and Drug Metabolizing Enzymes. *Mol. Pharm.* 14, 3142-3151.
- 1217 Wei, Z., Shi, Y., Chen, Y., Yu, S., Hao, J., Luo, J., Sha, X., Fang, X., 2010. Enhanced antitumor
1218 efficacy by Paclitaxel-loaded Pluronic P123/F127 mixed micelles against non-small cell lung cancer
1219 based on passive tumor targeting and modulation of drug resistance. *Eur. J. Pharm. Biopharm.* 75, 341-
1220 353.

- 1221 Wei, Z., Yuan, S., Hao, J., Fang, X., 2013. Mechanism of inhibition of P-glycoprotein mediated efflux
1222 by Pluronic P123/F127 block copolymers: Relationship between copolymer concentration and
1223 inhibitory activity. *Eur. J. Pharm. Biopharm.* 83, 266-274.
- 1224 Wojtal, K.A., Eloranta, J.J., Hruz, P., Gutmann, H., Drewe, J.r., Staumann, A., Beglinger, C., Fried, M.,
1225 Kullak-Ublick, G.A., Vavricka, S.R., 2009. Changes in mRNA Expression Levels of Solute Carrier
1226 Transporters in Inflammatory Bowel Disease Patients. *Drug Metab. Dispos.* 37, 1871-1877.
- 1227 Woodcock, D.M., Jefferson, S., Linsenmeyer, M.E., Crowther, P.J., Chojnowski, G.M., Williams, B.,
1228 Bertoncello, I., 1990. Reversal of the multidrug resistance phenotype with cremophor EL, a common
1229 vehicle for water-insoluble vitamins and drugs. *Cancer Res.* 50, 4199.
- 1230 Woodcock, D.M., Linsenmeyer, M.E., Chojnowski, G., Kriegler, A.B., Nink, V., Webster, L.K.,
1231 Sawyer, W.H., 1992. Reversal of multidrug resistance by surfactants. *Br. J. Cancer* 66, 62-68.
- 1232 Xiao, L., Yi, T., Chen, M., Lam, C.W.K., Zhou, H., 2016. A new mechanism for increasing the oral
1233 bioavailability of scutellarin with Cremophor EL: Activation of MRP3 with concurrent inhibition of
1234 MRP2 and BCRP. *Eur. J. Pharm. Sci.* 93, 456-467.
- 1235 Xu, Y., Egido, E., Li-Blatter, X., Müller, R., Merino, G., Bernèche, S., Seelig, A., 2015. Allocrite
1236 Sensing and Binding by the Breast Cancer Resistance Protein (ABCG2) and P-Glycoprotein (ABCB1).
1237 *Biochemistry* 54, 6195.

- 1238 Yamagata, T., Kusuhara, H., Morishita, M., Takayama, K., Benameur, H., Sugiyama, Y., 2007a. Effect
1239 of excipients on breast cancer resistance protein substrate uptake activity. *J. Control. Release* 124, 1-5.
- 1240 Yamagata, T., Kusuhara, H., Morishita, M., Takayama, K., Benameur, H., Sugiyama, Y., 2007b.
1241 Improvement of the oral drug absorption of topotecan through the inhibition of intestinal xenobiotic
1242 efflux transporter, breast cancer resistance protein, by excipients. *Drug Metab. Dispos.* 35, 1142-1148.
- 1243 Yamagata, T., Morishita, M., Kusuhara, H., Takayama, K., Benameur, H., Sugiyama, Y., 2009.
1244 Characterization of the inhibition of breast cancer resistance protein-mediated efflux of mitoxantrone
1245 by pharmaceutical excipients. *Int. J. Pharm.* 370, 216-219.
- 1246 Yoshida, N., Takagi, A., Kitazawa, H., Kawakami, J., Adachi, I., 2005. Inhibition of P-glycoprotein-
1247 mediated transport by extracts of and monoterpenoids contained in *Zanthoxyli Fructus*. *Toxicol. Appl.*
1248 *Pharmacol.* 209, 167-173.
- 1249 Yu, H., Hu, Y.Q., Ip, F.C.F., Zuo, Z., Han, Y.F., Ip, N.Y., 2011. Intestinal transport of
1250 bis(12)-hupyrindone in Caco-2 cells and its improved permeability by the surfactant Brij-35. *Biopharm.*
1251 *Drug Disposition* 32, 140-150.
- 1252 Yu Liang Siqi Li Ligong, C., 2015. The physiological role of drug transporters. *Protein Cell* 6, 334-
1253 350.

- 1254 Zhang, H., Yao, M., Morrison, R.A., Chong, S., 2003. Commonly used surfactant, Tween 80, improves
1255 absorption of P-glycoprotein substrate, digoxin, in rats. *Arch. Pharm. Res.* 26, 768-772.
- 1256 Zhao, G., Huang, J., Xue, K., Si, L., Li, G., 2013. Enhanced intestinal absorption of etoposide by self-
1257 microemulsifying drug delivery systems: Roles of P-glycoprotein and cytochrome P450 3A inhibition.
1258 *Eur. J. Pharm. Sci.* 50, 429-439.
- 1259 Zhao, W., Uehera, S., Tanaka, K., Tadokoro, S., Kusamori, K., Katsumi, H., Sakane, T., Yamamoto,
1260 A., 2016. Effects of Polyoxyethylene Alkyl Ethers on the Intestinal Transport and Absorption of
1261 Rhodamine 123: A P-glycoprotein Substrate by In Vitro and In Vivo Studies. *J. Pharm. Sci.* 105, 1526-
1262 1534.
- 1263 Zhu, S., Huang, R., Hong, M., Jiang, Y., Hu, Z., Liu, C., Pei, Y., 2009. Effects of polyoxyethylene (40)
1264 stearate on the activity of P-glycoprotein and cytochrome P450. *Eur. J. Pharm. Sci.* 37, 573-580.
- 1265 Ölander, M., Wiśniewski, J.R., Matsson, P., Lundquist, P., Artursson, P., Science for Life Laboratory,
1266 S., Medicinska och farmaceutiska, v., Uppsala, u., Institutionen för, f., Farmaceutiska, f., 2016. The
1267 Proteome of Filter-Grown Caco-2 Cells With a Focus on Proteins Involved in Drug Disposition. *J.*
1268 *Pharm. Sci.* 105, 817-827.

1270 **Figures Legend:**

1271 **Figure 1:** Structures of membrane transport proteins in the absence and presence of a substrate.

1272 Cartoons illustrate: **a)** Inward-facing P-glycoprotein (P-gp) (mouse Abcb1, left) and the outward-facing
 1273 P-gp (human ABCB1, right, substrate release), transmembrane domain 1 (TMD1) (1-6 transmembrane
 1274 helices (TMH)), TMD2 (7-12 TMH), extracted with modifications from (Aller et al., 2009; Kim and
 1275 Chen, 2018); **b)** Two monomers of breast cancer resistance protein (BCRP), BCRP monomer-1 (1-6
 1276 TMH, white) and -2 (1-6 TMH, dark gray), substrate-free state (left) using MsbA from *Escherichia coli*
 1277 as a template, substrate-bound state (right) using mouse Abcb1 as the template, extracted with
 1278 modifications from (Rosenberg et al., 2010; Rosenberg et al., 2015); **c)** Multidrug resistance-
 1279 associated protein 2 (MRP2), TM0 (1-5 TMH), TMD1 (6-11 TMH), TMD3 (12-17 TMH), Lasso motif
 1280 (L_0), left is when substrate-free state, and right when substrate-bound state, the molecular structure is
 1281 determined using bovine Mrp1 as a template, extracted with modifications from (Dallas et al., 2006;
 1282 Johnson and Chen, 2017); **d)** Lactose permease of *Escherichia coli* (LacY) representing a solute carrier
 1283 (SLC) member with 12 TMHs, LacY consists of two segments, each containing two repeat units of
 1284 three TMHs (1-3, 4-6, 7-9 and 10-12) as dark gray, black, white and light gray rods, respectively ,
 1285 outward-open conformation (V-shape, substrate-free state, left) and inward-open conformation
 1286 (inverted V-shape, substrate-released state, right) facing the extracellular and cytoplasmic side of the
 1287 cellular membrane, respectively, extracted with modifications from (Colas et al., 2016; Kumar et al.,
 1288 2018; Kumar et al., 2014; Radestock and Forrest, 2011). Nucleotide binding domain (NBD), adenosine
 1289 tri-phosphate (ATP), TMHs are depicted as rods, straight-dashed arrow represents the direction of
 1290 substrate movement, curved arrow represents the direction of helices movement during conformational
 1291 changes, post-translational modifications are not shown in the sub-figures, and black circle is a
 1292 substrate.

1293

Figure 2: Proposed mechanisms of P-glycoprotein inhibition by nonionic surfactants.

Cartoon shows: I) Surfactant-P-gp interaction via hydrogen bonding, the hydrophobic moiety of the surfactant partitions into the cell membrane, while the hydrogen bond acceptor groups in the hydrophilic moiety of the surfactant form hydrogen bonds with the hydrogen bond donor groups in P-gp, II) Alteration of membrane fluidity and/or (III) depleted intracellular ATP. Transmembrane helices are depicted as rods. Nucleotide binding domain (NBD), adenosine tri-phosphate (ATP), adenosine diphosphate (ADP), extracellular (Ex.), intracellular (In.), black triangle (P-gp substrate), red circle attached to a tail (nonionic surfactant), and black dashed lines (hydrogen bonds).

Table 1: Expression of selected transporters and carriers along the human intestine and in Caco-2 cells.

Logarithmic 10-step color scale and annotation of expression levels (very low-very high) have arbitrarily been defined for overview in the range 0-15 pmol/mg total protein and 0-450 fmol/mg total tissue^d.

To be inserted as a footnote under Table 1: LC-MS/MS-determined protein concentrations (pmol/mg total protein) of selected transporters and carriers in Caco-2 cells and segments of the human gastrointestinal tract. Protein concentrations from Caco-2 cells were obtained three weeks after seeding. Caco-2 cells were from three different sources: American Type Culture Collection (ATCC)^a, The European Collection of Authenticated Cell Cultures (ECACC)^b, and Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ)^c. The average values are depicted without statistical deviation parameters. For (Akazawa et al., 2018), the average were obtained from two reported values from two humans, and if one of the two values was below the lower limit of quantification (LLOQ), the other value is depicted. Intestinal segments were adapted from (Drozdik et al., 2014): Duodenum (D),

jejunum (J1-2), ileum (I1-2), and colon (C1-4). BLQ = below the LLOQ. ~~Logarithmic 10-step color scale and annotation of expression level (very low-very high) have arbitrarily been defined for overview in the range 0-15 pmol/mg total protein and 0-450 fmol/mg total tissue^d.~~

Table 2: *In vitro* ~~h~~ impact of selected nonionic surfactants and polyethylene glycol (PEG) derivatives on P-glycoprotein *in vitro*.

~~To be inserted as a footnote under Table 2:~~ Accumulation (accum.), Approximately (approx.), Respectively (resp.), Surfactant (surf.), Concentration (conc.), Dependent (dep.), Not specified in the study (ns), Apical to basolateral (A-B), Mucosal to serosal (M-S), Permeability (P_{app}), Any increase or decrease described in the table means significant $P < 0.05$, Resistance Modification Index (RMI), Mouse embryo fibroblasts transfected with MDR1 (NIH-MDR1-G185), Mouse embryo fibroblasts transfected with MDR1 (NIH-MDR1-G185), P-gp overexpressing human melanoma cell line (MDA-MB-435/LCC6MDR1), P-gp overexpressing human ovarian carcinoma cell line (NCI/ADR-RES), Adriamycin-resistant of murine leukaemia P388 cells (P388/ADR), P-gp variant of human epithelial cells KB 3-1 (KB 8-5-11 cells), In Vitro Diffusion Chamber Method (In vitro DCM), ATPlite 1step Assay kit was from PerkinElmer, P-gp containing membranes of Chinese hamster lung fibroblasts (DC-3F/ADX cells), MDR cell subline of Chinese hamster ovary cells Aux-B1(CH⁺C5), Bovine brain microvessel endothelial cells (BBMEC), Vinblastine-resistant derivative of Human Caucasian acute lymphoblastic leukaemia CCRF-CEM cells (R100 cells), Human lung adenocarcinoma cell line A549 treated with paclitaxel (A549/Taxol), Porcine kidney epithelial cell line (LLC-PK1-MDR1), LLC-PK1 stably expressing MDR1 (LLC-MDR1), P-gp overexpressing human oral epidermal carcinoma (KBv), For cremophor[®] EL and Solutol[®] HS 15, RMI was measured at 10 $\mu\text{g/mL}$, P-gp overexpressing human melanoma cell line (MDA-MB-435/LCC6MDR1), P-gp overexpressing human ovarian carcinoma cell

line NCI/ADR-RES, MDR cell subline of human breast carcinoma MCF-7 cells (MCF7/ADR), Resistance reversion index ($\text{Log}(\text{IC}_{50.0}/\text{IC}_{50})$) was determined as a ratio of IC_{50} of Doxorubicin in the assay buffer and surfactant solution, Vincristine-resistant derivative of K562 (7962 cells), Human lung cancer cells (A549), Human P-gp overexpressing membranes obtained from baculovirus-infected insect cells (High Five, BTI-TN5B1-4), Clonal isolate derived from the *Spodoptera frugiperda* cell line IPLB-Sf-21-AE (Sf9), The disappearance of the drug in perfusate (P_{lumen}) as well as the appearance of the drug in mesenteric vein blood (apparent permeability coefficient, P_{blood}), P-gp overexpressing of Chinese hamster ovary AA8 cells (Emt^{R1}). For an overview of the effects of more surfactants on P-gp, (see Table S1).

Table 43: Impact of nonionic surfactants on intestinal P-glycoprotein in rats.

In vivo pre-clinical studies were performed in male ^a: Sprague-Dawley rats, ^b: Wistar albino rats.

Synonyms of surfactants are available in Table 2.

Table 34: In vitro and in vivo ~~I~~mpact of nonionic surfactants on breast cancer resistance protein, BCRP, in vitro.

To be inserted as a footnote under Table 4: Concentration (conc.), Approximately (approx.), Respectively (resp.), Plasma membrane vesicle of cells containing human ABCG2 (Membrane vesicles BCRP), Clonal isolate derived from the *Spodoptera frugiperda* cell line IPLB-Sf-21-AE (Sf9 insect cells), *In Vitro* Diffusion Chamber Method (*In vitro* DCM), Serosal to mucosal (S-M), *In situ* closed-

loop method (*In situ* CLM), Wild type (WT), Sprague-Dawley (SD), Synonyms of surfactants are available in Table 2.

~~Table 4: Impact of nonionic surfactants on intestinal P-glycoprotein in rats.~~

~~*In vivo* pre-clinical studies were performed in male ^a: Sprague-Dawley rats, ^b: Wistar albino rats.~~

~~Synonyms of surfactants are available in Table 2.~~

Table 5: *In vitro* and *in vivo* impact of nonionic surfactants and co-surfactants on multidrug resistance-associated protein 2, MRP2, *in vitro*.

To be inserted as a footnote under Table 5: Concentration (Conc.), Respectively (resp.), Membrane vesicles prepared from *Spodoptera frugiperda* (Sf9) insect cells over-expressing human MRP2 (Membrane vesicles of Sf9 MRP2), ATP measurements were performed using ATP luciferin/Luciferase assay, Wild type (WT), Synonyms of surfactants are available in Table 2 and 3.

Table 6: Nonionic surfactants inhibited solute carriers (SLCs) *in vitro*.

To be inserted as a footnote under Table 6: IC₅₀ were estimated from uptake transport assay. For Regev et al. 2002, impact of surfactant on bi-directional transport assay was shown. 1-methyl-4-phenylpyridinium acetate (MPP⁺), Monocarboxylic acid transporter (MCT, SLC16A1), Organic cation transporter 1 (OCT1, SLC22A1), (OCT2, SLC22A2), (OCT3, SLC22A3), Peptide transporter 1 (PEPT1, SLC15A1), (PEPT2, SLC15A2), Organic anion transporting polypeptide 1A2 (OATP1A2, SLC21A3), (OATP2B1, SLC21A9). Human embryonic kidney cells stably transfected with OATP1A2

(HEK OATP1A2), or with OATP2B1 (HEK OATP2B1), Chinese hamster ovary cells stably transfected with rbOCT1(CHO-K1 rbOCT1), Madin-Darby canine kidney cells stably transfected with OCT1-3 (MDCKII OCT1-3), or with PEP2 (MDCKII PEPT2). Synonyms of surfactants available in Table 2 and 5.

Table S1: *In vitro* impact of nonionic surfactants and polyethylene glycol (PEG) derivatives on P-glycoprotein *in vitro*.

Accumulation (accum.), Approximately (approx.), Respectively (resp.), Surfactant (surf.), Concentration (conc.), Dependent (dep.), Not specified in the study (ns), Apical to basolateral (A-B), Mucosal to serosal (M-S), Permeability (P_{app}), Any increase or decrease described in the table means significant $P < 0.05$, Resistance Modification Index (RMI), Mouse embryo fibroblasts transfected with MDR1 (NIH-MDR1-G185), Mouse embryo fibroblasts transfected with MDR1 (NIH-MDR1-G185), P-gp overexpressing human melanoma cell line (MDA-MB-435/LCC6MDR1), P-gp overexpressing human ovarian carcinoma cell line (NCI/ADR-RES), Adriamycin-resistant of murine leukemia P388 cells (P388/ADR), P-gp variant of human epithelial cells KB 3-1 (KB 8-5-11 cells), In Vitro Diffusion Chamber Method (In vitro DCM), ATPlite 1step Assay kit was from PerkinElmer, P-gp containing membranes of Chinese hamster lung fibroblasts (DC-3F/ADX cells), Concentration of half-maximum activation (K_1), Concentration of half-maximum inhibition (K_2), MDR cell subline of Chinese hamster ovary cells Aux-B1(CH⁺C5), Bovine brain microvessel endothelial cells (BBMEC), Vinblastine-resistant derivative of Human Caucasian acute lymphoblastic leukemia CCRF-CEM cells (R100 cells), Human lung adenocarcinoma cell line A549 treated with paclitaxel (A549/Taxol), Porcine kidney epithelial cell line (LLC-PK1-MDR1), LLC-PK1 stably expressing MDR1 (LLC-MDR1), P-gp overexpressing human oral epidermal carcinoma (KBv), For cremophor[®] EL and Solutol[®] HS 15, RMI

was measured at 10 $\mu\text{g/mL}$, P-gp overexpressing human melanoma cell line (MDA-MB-435/LCC6MDR1), P-gp overexpressing human ovarian carcinoma cell line NCI/ADR-RES, MDR cell subline of human breast carcinoma MCF-7 cells (MCF7/ADR), Resistance reversion index ($\text{Log}(\text{IC}_{50.0}/\text{IC}_{50})$) was determined as a ratio of IC_{50} of Doxorubicin in the assay buffer and surfactant solution, Vincristine-resistant derivative of K562 (7962 cells), Human lung cancer cells (A549), Human P-gp overexpressing membranes obtained from baculovirus-infected insect cells (High Five, BTI-TN5B1-4), Clonal isolate derived from the *Spodoptera frugiperda* cell line IPLB-Sf-21-AE (Sf9), The disappearance of the drug in perfusate (P_{lumen}) as well as the appearance of the drug in mesenteric vein blood (apparent permeability coefficient, P_{blood}), P-gp overexpressing of Chinese hamster ovary AA8 cells (Emt^{R1}).

Table 1: Expression of selected transporters and carriers along the human intestine and in Caco-2 cells.

~~LC-MS/MS-determined protein concentrations (pmol/mg total protein) of selected transporters and carriers in Caco-2 cells and segments of the human gastrointestinal tract. Protein concentrations from Caco-2 cells were obtained three weeks after seeding. Caco-2 cells were from three different sources: American Type Culture Collection (ATCC)^a, The European Collection of Authenticated Cell Cultures (ECACC)^b, and Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ)^c. The average values are depicted without statistical deviation parameters. For (Akazawa et al., 2018), the average were obtained from two reported values from two humans, and if one of the two values was below the lower limit of quantification (LLOQ), the other value is depicted. Intestinal segments were adapted from (Drozdik et al., 2014): Duodenum (D), jejunum (J1-2), ileum (I1-2), and colon (C1-4). BLQ = below the LLOQ. Logarithmic 10-step color scale and annotation of expression level (very low-very~~

high) have *arbitrarily* been defined for overview in the range 0-15 pmol/mg total protein and 0-450

fmol/mg total tissue^d:

	<i>Very low expression</i>		<i>Low expression</i>		<i>Intermediate expression</i>	
pmol protein/mg total protein	0-0.0099	0.0100-0.0248	0.0249-0.0621	0.0622-0.154	0.155-0.386	0.387-0.9
fmol protein/mg total tissue ^d	0-0.299	0.300-0.747	0.748-1.86	1.87-4.65	4.66-11.5	11.6-28

Transp orter or carrier	Caco- 2	Reference	Intestinal segment								Reference	
			D	J1	J2	I1	I2	C1	C2	C3		C4
P-gp	2.06 ^a	(Ölander et al., 2016)	7.67 ^d	33.02 ^d	47.41 ^d		70.78 ^d		9.98 ^d			(Drozdzik et al., 2019)
	4.1 ^b	(Uchida et al., 2015)	0.290	0.408	0.475	0.711	1.06	0.145	0.304	0.228	0.368	(Drozdzik et al., 2014)
	1.0 ^c	(Brück et al., 2017)		1.22								(Lloret-Linares et al., 2016)
				0.614			0.656					(Gröer et al., 2013)
					1.89		0.20					(Harwood et al., 2015)
				2.43		4.93						(Akazawa et al., 2018)
BCRP	0.0117 ^a	(Ölander et al., 2016)	5.51 ^d	19.58 ^d	26.97 ^d		30.47 ^d		5.13 ^d			(Drozdzik et al., 2019)
	1.79 ^b	(Uchida et al., 2015)	0.190	0.277	0.356	0.405	0.359	0.150	0.0438	0.153	0.160	(Drozdzik et al., 2014)
	0.5 ^c	(Brück et al., 2017)			1.25							(Miyauchi et al., 2016)
				0.574			0.241					(Gröer et al., 2013)
					2.56		1.60					(Harwood et al., 2015)

				4.65	8.07					2015) (Akazawa et al., 2018)
MRP2	0.134 ^a	(Ölander et al., 2016)	11.8 8 ^d	22. 37 ^d	22.5 2 ^d		19.8 4 ^d		16.6 9 ^d	(Drozdzik et al., 2019)
	0.649 ^b	(Uchida et al., 2015)	0.75 8	1.0 3	0.94 6	0.7 64	0.80 8	1.4 1	1.77 3	(Drozdzik et al., 2014)
	0.8 ^c	(Brück et al., 2017)		0.1 16						(Lloret-Linares et al., 2016)
				1.07		0.350				(Gröer et al., 2013)
					0.59		BL Q			(Harwood et al., 2015)
MRP3				0.835		1.16				(Akazawa et al., 2018)
	0.423 ^a	(Ölander et al., 2016)	17.2 8 ^d	30. 47 ^d	31.2 5 ^d		22.5 8 ^d		28.7 9 ^d	(Drozdzik et al., 2019)
	BLQ ^b	(Uchida et al., 2015)	0.85 0	0.6 39	0.50 6	0.5 52	0.69 6	1.5 3	2.10 1	(Drozdzik et al., 2014)
	BLQ ^c	(Brück et al., 2017)		1.9 1						(Lloret-Linares et al., 2016)
				0.309		0.686				(Gröer et al., 2013)
PEPT1				0.501		0.303				(Akazawa et al., 2018)
	0.342 ^a	(Ölander et al., 2016)	25.6 1 ^d	84. 17 ^d	109. 6 ^d		107. 3 ^d		3.27 d	(Drozdzik et al., 2019)
	1.48 ^b	(Uchida et al., 2015)	2.63	3.4 9	4.23	4.6 2	4.89	0.2 98	0.21 0	(Drozdzik et al., 2014)
	5.2 ^c	(Brück et al., 2017)			1.60					(Miyauchi et al., 2016)
				2.45		4.73				(Gröer et al., 2013)
OATP2 B1				8.34		10.7				(Akazawa et al., 2018)
	2.66 ^a	(Ölander et al., 2016)	5.30 d	7.2 1 ^d	8.02 d		8.06 d		8.00 d	(Drozdzik et al., 2019)
	0.771 ^b	(Uchida et al., 2015)	0.42 8	0.5 56	0.48 6	0.4 64	0.48 2	0.4 78	0.73 1	(Drozdzik et al., 2014)
	3.3 ^c	(Brück et al., 2017)			0.54 0					(Miyauchi et al., 2016)
				0.299		0.267				(Gröer et al., 2013)
				BLQ		BLQ				(Akazawa et al., 2018)

											2018)	
OATP1 A2	BLQ ^b	(Uchida et al., 2015)	BL Q	BL Q	BL Q		BL Q		BL Q		(Drozdzik et al., 2019)	
	BLQ ^c	(Brück et al., 2017)	BL Q	BL Q	BL Q	BL Q	BL Q	BL Q	BL Q	BL Q	(Drozdzik et al., 2014)	
					BL Q						(Miyauchi et al., 2016)	
				BLQ		BLQ					(Gröer et al., 2013)	
				0.336		0.189					(Akazawa et al., 2018)	
OCT1	BLQ ^a	(Ölander et al., 2016)	1.61 _d	4.2 _{2^d}	6.02 _d		5.12 _d		2.79 _d		(Drozdzik et al., 2019)	
	BLQ ^b	(Uchida et al., 2015)	0.66 5	0.6 47	0.56 6	0.8 02	0.84 2	0.4 69	0.69 5	0.7 25	0.6 32	(Drozdzik et al., 2014)
	BLQ ^c	(Brück et al., 2017)			BL Q							(Miyauchi et al., 2016)
				BLQ		0.480						(Gröer et al., 2013)
OCT3	BLQ ^b	(Uchida et al., 2015)	BL Q	BL Q	BL Q		BL Q					(Drozdzik et al., 2019)
	BLQ ^c	(Brück et al., 2017)	0.06 70	0.0 564	0.06 29	0.0 531	0.0 687	0.1 26	0.10 7	0.1 16	0.1 35	(Drozdzik et al., 2014)
					BL Q							(Miyauchi et al., 2016)
				BLQ		0.077						(Gröer et al., 2013)
				0.551		BLQ					(Akazawa et al., 2018)	
MCT	1.72 ^a	(Ölander et al., 2016)	61.1 3 ^d	78. 81 ^d	75.1 4 ^d		43.6 5 ^d		112. 6 ^d			(Drozdzik et al., 2019)
	0.871 _b	(Uchida et al., 2015)			1.85							(Miyauchi et al., 2016)
				1.54		2.41						(Akazawa et al., 2018)

1438

1439

1440

1441

1442

1443

Table 2: *In vitro* impact of selected nonionic surfactants and polyethylene glycol (PEG) derivatives on P-glycoprotein *in vitro*.

Accumulation (accum.), Approximately (approx.), Respectively (resp.), Surfactant (surf.), Concentration (conc.), Dependent (dep.), Not specified in the study (ns), Apical to basolateral (A-B), Mucosal to serosal (M-S), Permeability (P_{app}), Any increase or decrease described in the table means significant $P < 0.05$, Resistance Modification Index (RMI), Mouse embryo fibroblasts transfected with MDR1 (NIH-MDR1-G185), Mouse embryo fibroblasts transfected with MDR1 (NIH-MDR1-G185), P-gp overexpressing human melanoma cell line (MDA-MB-435/LCC6MDR1), P-gp overexpressing human ovarian carcinoma cell line (NCI/ADR-RES), Adriamycin-resistant of murine leukaemia P388 cells (P388/ADR), P-gp variant of human epithelial cells KB-3-1 (KB-8-5-11 cells), In Vitro Diffusion Chamber Method (In vitro DCM), ATPlite 1step Assay kit was from PerkinElmer, P-gp containing membranes of Chinese hamster lung fibroblasts (DC-3F/ADX cells), MDR cell subline of Chinese hamster ovary cells Aux-B1(CH⁺C5), Bovine brain microvessel endothelial cells (BBMEC), Vinblastine-resistant derivative of Human Caucasian acute lymphoblastic leukaemia CCRF-CEM cells (R100 cells), Human lung adenocarcinoma cell line A549 treated with paclitaxel (A549/Taxol), Porcine kidney epithelial cell line (LLC-PK1-MDR1), LLC-PK1 stably expressing MDR1 (LLC-MDR1), P-gp overexpressing human oral epidermal carcinoma (KBv), For cremophor[®]-EL and Solutol[®]-HS 15, RMI was measured at 10 $\mu\text{g/mL}$, P-gp overexpressing human melanoma cell line (MDA-MB-435/LCC6MDR1), P-gp overexpressing human ovarian carcinoma cell line NCI/ADR-RES, MDR cell subline of human breast carcinoma MCF-7 cells (MCF7/ADR), Resistance reversion index ($\text{Log}(\text{IC}_{50-0}/\text{IC}_{50})$) was determined as a ratio of IC_{50} of Doxorubicin in the assay buffer and surfactant solution, Vincristine-resistant derivative of K562 (7962 cells), Human lung cancer cells (A549), Human P-gp overexpressing membranes obtained from baculovirus-infected insect cells (High Five, BTI-TN5B1-4), Clonal isolate derived from the *Spodoptera frugiperda* cell line IPLB-Sf-21-AE (Sf9), The disappearance of the drug in perfusate (P_{lumen}) as well as the appearance of the drug in mesenteric

1472 vein blood (apparent permeability coefficient, P_{blood}), P-gp overexpressing of Chinese hamster ovary
1473 AA8 cells (Emt^{R1}). For an overview of the effects of more surfactants on P-gp, (see Table S1).

	Conc.	Substrate	Cell line/Tissue	Assay	Impact	Refer
EL, astor astor 5	(1:1000)	Daunorubicin	R100 cells	Uptake transport	Increased intracellular accum.	(Wood
	(1:1000)	Daunorubicin	7962 cells	Uptake transport	Increased intracellular accum.	(Wood
	(1:1000)	Daunorubicin	R100 cells	Uptake transport	Increased intracellular accum.	(Wood
	(1:1000)	Daunorubicin	P388/ADR	Uptake transport	Increased intracellular accum.	(Wood
	0.0001-0.1% (w/v)	Acf(N-Mef) ₂ NH ₂	Caco-2 Cell monolayers	Bi-directional transport	Increased A-B and decreased B-A P _{app} .	(Nerur
	100 µg/mL	Rh 123	KB 8-5-11 cells	Uptake transport	Enhanced the fluorescence of Rh 123 by 3-fold.	(Buck
	3-20 µg/mL	Doxorubicin	KB 8-5-11 cells	MTT	Decreased IC ₅₀ in a conc. dep. manner. RMI = 1.5 ± 0.0	(Buck
	3-20 µg/mL	Vinblastine	KB 8-5-11 cells	MTT	Decreased IC ₅₀ in a conc. dep. manner. RMI = 1.1 ± 0.1	(Buck
	3-20 µg/mL	Colchicine	KB 8-5-11 cells	MTT	Decreased IC ₅₀ in a conc. dep. manner. RMI = 1.3 ± 0.1	(Buck
	3-20 µg/mL	Etoposide	KB 8-5-11 cells	MTT	Decreased IC ₅₀ in a conc. dep. manner. RMI = 1.2 ± 0.3	(Buck
	3-20 µg/mL,	Actinomycin D	KB 8-5-11 cells	MTT	Decreased IC ₅₀ in a conc. dep. manner. RMI = 1 ± 0.2	(Buck
	0.01-1 mM	Rh 123	Caco-2 cell monolayers	Bi-directional transport	Increased A-B and decreased B-A P _{app} in a conc. dependent manner.	(Rege
	0.02-2% (w/v)	Cyclosporine A	Caco-2 cell monolayers	Bi-directional transport	Decreased B-A P _{app} .	(Chiu
	0.005-0.5% (w/v)	Rh 123	Rat intestinal membrane	Bi-directional transport (In vitro DCM)	Increased S-M and decreased M-S P _{app} .	(Shon
glycol	400 µM	Doxorubicin	Caco-2 cell monolayers	Bi-directional transport	Increased A-B P _{app} .	(Al-Sa
	300 µM	Etoposide	Caco-2 cell monolayers	Bi-directional transport	Increased A-B and decreased B-A P _{app} .	(Al-A
	1% SMEDDS containing 50% (w/w) of surf.	Etoposide (SMEDDS)	Intestinal segments from rats' ileum	In situ single- pass perfusion experiments	Increased intestinal P _{app} . Increased P _{Blood} and P _{Lumen} .	(Zhao
	0.3-1000 µM	Digoxin	MDCKII MDR1	Uptake transport	Increased intracellular accum. in a conc. dependent manner. IC ₅₀ = 12 µM	(Gurja
	2.5-20% (w/v)	Paclitaxel	Caco-2 cell monolayers	Bi-directional transport	Increased A-B and decreased B-A P _{app} in a conc. dependent manner.	(Hugg

20% (w/v)	Paclitaxel	Caco-2 cell monolayers	Bi-directional transport	Increased A-B and decreased B-A P_{app} in a conc. dependent manner.	(Hugg
20% (v/v)	Ranitidine	Caco-2 cell monolayers	Bi-directional transport	Decreased ranitidine ER.	(Ashir
300 μ M			PREDEASY ATPase Kit	Increased P-gp ATPase activity.	al., 20
1, 5 and 20% (w/v)	Digoxin	Rat jejunal membrane	Bi-directional transport (In vitro DCM)	Decreased S-M flux by 47, 57 and 64%, resp., compared to control.	(Ashir al., 20
0.1-20% (v/v)	Rh 123	Rat intestinal membrane	Bi-directional transport (In vitro DCM)	Decreased S-M P_{app} in a conc. dep. manner.	(Johns 2002)
20% (v/v)	Ranitidine	Caco-2 cell monolayers	Bi-directional transport	Decreased ER.	(Shen
0.5 and 1% (v/v)	Ranitidine	Caco-2 cell monolayers	Bi-directional transport	Enhanced A-B and decreased B-A P_{app} .	(Ashir al., 20
300 μ M			PREDEASY ATPase Kit	Increased P-gp ATPase activity.	(Ashir al., 20
1 and 2% (w/v)	Rh 123	Caco-2 cells	Uptake transport	Enhanced Rh 123 intracellular accum.	(Hoda
1 and 2% (w/v)		Caco-2 cells	Western blotting	Decreased P-gp expression.	(Hoda
	Doxorubicin	KBv		$\text{Log (IC}_{50.0}/\text{IC}_{50}) = 0.7$	(Batra 1999)
	Doxorubicin	MCF7/ADR		$\text{Log (IC}_{50.0}/\text{IC}_{50}) = 0.8$	(Batra 1999)
	Doxorubicin	CH ^r C5		$\text{Log (IC}_{50.0}/\text{IC}_{50}) = 2$	(Batra 1999)
Log M = -5	Rh 123	KBv	Uptake transport	Enhanced Rh 123 accum. by approx. 6.5-fold.	(Batra 1999)
0.001-1%	Rh 123	LLC-PK1-MDR1	Uptake transport	Increased Rh 123 accum.	(Batra 2001)
0.001-1%	Digoxin	LLC-PK1-MDR1	Uptake transport	Increased digoxin accum.	(Batra 2001)
0.01-1%	Digoxin	BBMEC	A-B transport	Increased A-B transport.	(Batra 2001)
0.01%	Digoxin	BBMEC	A-B transport	Increased A-B and decreased transport.	(Batra 2001)
0.01 and 0.1% (w/v)	Digoxin	Rat jejunal membrane	Bi-directional transport (In vitro DCM)	Decreased S-M flux.	(Johns 2002)
0.01% (w/v)	Rh 123	BBMEC	Uptake transport	Enhanced Rh 123 accum. by approx. 2-fold. Depleted intracellular ATP content. Decreased the P-gp ATPase	(Batra 2003)

0.01% (w/v)	Rh 123	BBMEC	ATP luciferin/ luciferase	activity. Depleted intracellular ATP content.	(Batra 2003)
0.01% (w/v)	Rh 123	KBv	Pgp ATPase activity	Decreased the P-gp ATPase activity.	(Batra 2003)
0.1% w/v		P-gp membranes from Gentest Co.	P-gp ATPase Assay	Decreased V_{max} and increased K_m significantly.	(Batra 2004)
0.01 and 0.1% w/v	Vincristine	P-gp membranes from Gentest Co.	P-gp ATPase Assay	Decreased V_{max} and increased K_m significantly.	(Batra 2004)
0.5 % (w/v)	Rh 123	Rats' jejunal segments	M-S transport. (Ussing chamber)	Increased M-S P_{app} by 1.9-fold.	(Föger 2003)
0.1% w/w		P-gp membranes (High Five, BTI- TN5B1-4)	P-gp ATPase Assay	Abolished P-gp ATPase activity completely.	(Shaik 2004)
0.01% w/w	Verapamil	P-gp membranes (High Five, BTI- TN5B1-4)	P-gp ATPase Assay	Inhibited verapamil-stimulated P-gp ATPase activity.	(Shaik 2004)
0.01% w/w	Nelfinavir	P-gp membranes (High Five, BTI- TN5B1-4)	P-gp ATPase Assay	Abolished the nelfinavir stimulated P-gp ATPase activity.	(Shaik 2004)
0.01% w/w	Nelfinavir	MDCKII MDR1	Uptake transport	Enhanced nelfinavir accum.	(Shaik 2004)
0.01% w/w	Saquinavir	MDCKII MDR1	Uptake transport	Increased saquinavir accum. by 2- fold.	(Shaik 2004)
0.01% w/w	Saquinavir	LLC-PK1-MDR1	Uptake transport	Increased saquinavir accum. by 5- fold.	(Shaik 2004)
0-600 ng/mL		Membrane vesicles of Emt ^{R1} cells	Phosphate release measurements	Reduced P-gp ATPase activity in a conc. dep. manner.	(Rege 2004)
0-300 ng/mL	Doxirubicin	Large unilamellar vesicles (LUV)	Trans-bilayer movement	Decreased Flip-Flop Life-Time of doxorubicin in a conc. dep. manner.	(Rege 2004)
30-100 ng/mL	Clacein-AM	Emt ^{R1} cells	Calcein-AM efflux	Enhanced calcein-AM uptake in a conc. dep. manner.	(Rege 2004)
0.5% (w/v)	Digoxin	Rat everted gut sac model	Uptake transport	Enhanced digoxin accum.	(Corna 2004)
200 μ M	Epirubicin	Caco-2 cell monolayers	Bi-directional transport	Increased A-B and decreased B-A P_{app} .	(Lo, 2 004)
20-200 μ M	Epirubicin	Caco-2 cells	Uptake transport	Enhanced fluorescent epirubicin accum. in a conc. dep. manner.	(Lo, 2 004)
200 μ M	Epirubicin	Everted sacs of rat's jejunum or ileum	M-S transport	Increased M-S P_{app} .	(Lo, 2 004)
200 μ M	Doxorubicin	Caco-2 cell monolayers	Bi-directional transport	Increased A-B P_{app} .	(Al-Sa 2016)

	200 μ M	Digoxin	Caco-2 cell monolayers	Bi-directional transport	Increased A-B and decreased B-A P_{app} .	(Niels)
	0.2-500 μ M	Digoxin	Caco-2 cell monolayers	Bi-directional transport	Increased A-B and decreased B-A P_{app} in a conc. dep. manner.	(Niels)
	0.2-500 μ M	Digoxin	MDCKII MDR1 cell monolayers	Bi-directional transport	Increased A-B and decreased B-A P_{app} in a conc. dep. manner.	(Niels)
	200-500 μ M	Etoposide	Caco-2 cell monolayers	Bi-directional transport	Increased A-B P_{app} .	(Al-A)
	0.2-500 μ M	Etoposide	Caco-2 cell monolayers	Bi-directional transport	Decreased B-A P_{app} .	(Al-A)
	20-500 μ M	Etoposide	MDCKII MDR1 cell monolayers	Bi-directional transport	Increased A-B and decreased B-A P_{app} .	(Al-A)
		Calcein-AM	MDCKII MDR1	Calcein-AM efflux	Increased calcein fluorescence in a conc. dep. manner. $IC_{50} = 11 \mu$ M.	(Al-A) 2018b
	200 μ M	Digoxin	MDCKII MDR1	Bi-directional transport	Increased A-B and decreased B-A P_{app} . Increased intracellular accum. of digoxin from the apical side.	(Al-A) 2018b
	0.3-1000 μ M	Digoxin	MDCKII MDR1	Uptake transport	Increased intracellular accum. in a conc. dependent manner. $IC_{50} = 74 \mu$ M.	(Gurja)
0,	200 μ M	Epirubicin	Everted sacs of jejunum or ileum of rats	M-S transport	Increased M-S P_{app}	(Lo, 2)
ne	20-200 μ M	Epirubicin	Caco-2 cells	Uptake transport	Enhanced intracellular accum. of fluorescent epirubicin in a conc. dep. manner.	(Lo, 2)
oleate	200 μ M	Epirubicin	Caco-2 cell monolayers	Bi-directional transport	Increased A-B and decreased B-A P_{app} .	(Lo, 2)
	0.5 w/v	Digoxin	Rat everted gut sac model	Uptake transport	Enhanced digoxin uptake.	(Corma) 2004)
	0.01-1 mM	Rh 123	Caco-2 cell monolayers	Bi-directional transport	Increased A-B and decreased B-A P_{app} in a conc. dep. manner.	(Rege)
		Rh 123	Caco-2	Uptake transport	Increased Rh 123 accum.	(Kiss)
		Rh 123	Caco-2 cell monolayers	Bi-directional transport	Increased A-B and decreased B-A P_{app} .	(Kiss)
		Calcein-AM	Caco-2 cells	Uptake transport	Increased calcein accum.	(Kiss)
	(1:10000)	Daunorubicin	R100 cells	Uptake transport	Increased intracellular daunorubicin accum.	(Wood) 1992)
	0.0001-1 % (w/v)	Acf(N-Mef) ₂ NH ₂	Caco-2 Cell monolayers	Bi-directional transport	Increased A-B and decreased B-A P_{app} .	(Nerun) 1996)
	0.1% (w/v)	Rh 123	Rat intestinal membrane	Bi-directional transport	Reduced S-M/M-S ratio.	(Shon)

S, S 15, glycol ate, ted aric ate	0.06-0.66 μ M	Verapamil	Membrane vesicles of NIH-MDR1-G185	(In vitro DCM) Phosphate release measurements	Inhibition of verapamil-induced P-gp ATPase activity.	(Li-BL 2009)
	0.001-0.05 w/v	Clacein-AM	MDCKII MDR1	Calcein-AM efflux	Enhanced calcein fluorescence by approx. 2-fold.	(Hank
	150 μ M	Bis(12)-hupyridone	Caco-2 cell monolayers	Bi-directional transport	Increased A-B P_{app} and decreased B-A P_{app} .	(Yu et
	10 μ M	Bis(12)-hupyridone	Caco-2 cell monolayers	Bi-directional transport	Decreased B-A P_{app} .	(Yu et
	200 μ M	Doxorubicin	Caco-2 cell monolayers	Bi-directional transport	Increased A-B P_{app} .	(Al-Sa 2016)
	200 μ M	Digoxin	Caco-2 cell monolayers	Bi-directional transport	Increased A-B and decreased B-A flux.	(Niels
	300 μ M	Etoposide	Caco-2 cell monolayers	Bi-directional transport	Increased A-B and decreased B-A P_{app} .	(Al-A
	1% SMEDDS containing 50% (w/w) of surf.	Etoposide (SMEDDS)	Intestinal segments from rats' ileum	In situ single-pass perfusion experiments	Increased intestinal P_{app} . Increased P_{Blood} and P_{Lumen} .	(Zhao
		Calcein-AM	MDCKII MDR1	Calcein-AM efflux	Increased calcein fluorescence in a conc. dep. manner. IC_{50} = 69 μ M	(Al-A 2018b
	200 μ M	Digoxin	MDCKII MDR1	Bi-directional transport	Decreased B-A P_{app} . Increased intracellular accum. of digoxin from the apical side.	(Al-A 2018b
	0.3-100 μ M	Digoxin	MDCKII MDR1	Uptake transport	Increased intracellular accum. in a conc. dep. manner. IC_{50} = 45 μ M	(Gurja
	0.05-0.5 % (w/v)	Digoxin	Rat everted gut sac model	Uptake transport	Enhanced digoxin accum.	(Corna 2004)
	(1:10000)	Daunorubicin	R100 cells	Uptake transport	Increased intracellular daunorubicin.	(Wood 1992)
		Etoposide	C6 glioma cells	MTT	Decreased IC_{50} by 10-fold.	(Lamp Benoi
		Etoposide	F98 glioma cells	MTT	Decreased IC_{50} by 3-fold.	(Lamp Benoi
		Etoposide	9L glioma cells	MTT	Decreased IC_{50} by 8-fold.	(Lamp Benoi
	35-39% Lipid nanoparticles		P-gp exhibiting membrane vesicles	ATPase kit (SPIbio®, Massy, France)	Decreased ATPase activity.	(Lamp Benoi
	5-100 μ g/mL	Rh 123	KB 8-5-11 cells	Uptake transport	Enhanced fluorescence of Rh 123 in a conc. dep. manner.	(Buck 1995)
	3-20 μ g/mL	Doxorubicin	KB 8-5-11 cells	MTT	Decreased IC_{50} in a conc. dep. manner. $RMI = 6 \pm 3.2$	(Buck 1995)

I glycol e, GS, glycol	3-20 µg/mL	Vinblastine	KB 8-5-11 cells	MTT	Decreased IC ₅₀ in a conc. dep. manner. RMI = 2 ± 1	(Buck 1995)
	3-20 µg/mL	Colchicine	KB 8-5-11 cells	MTT	Decreased IC ₅₀ in a conc. dep. manner. RMI = 4.2 ± 0.7	(Buck 1995)
	3-20 µg/mL	Etoposide	KB 8-5-11 cells	MTT	Decreased IC ₅₀ in a conc. dep. manner. RMI = 2.7 ± 0.7	(Buck 1995)
	3-20 µg/mL	Actinomycin D	KB 8-5-11 cells	MTT	Decreased IC ₅₀ in a conc. dep. manner. RMI = 2.3 ± 0.9	(Buck 1995)
	3-20 µg/mL	Paclitaxel	KB 8-5-11 cells	MTT	Decreased IC ₅₀ in a conc. dep. manner. RMI = 10 ± 1.2	(Buck 1995)
	0.1 – 100 µM	Colchicine	KB 8-5-11 cells	Colorimetric (Crystal violet)	Decreased IC ₅₀ in a conc. dep. manner. RMI = 34.5 ± 2.5	(Coon 1995)
	0.1 – 100 µM	Vinblastine	KB 8-5-11 cells	Colorimetric (Crystal violet)	Decreased IC ₅₀ in a conc. dep. manner. RMI = 27.7 ± 2.3	(Coon 1995)
	0.1 – 100 µM	Doxorubicin	KB 8-5-11 cells	Colorimetric (Crystal violet)	Decreased IC ₅₀ in a conc. dep. manner. RMI = 41.7 ± 3	(Coon 1995)
	70 µM	Rh 123	KB 8-5-11 cells	Uptake transport	Increased accum. by 50-fold.	(Coon 1995)
	0.1-1 % (w/v)	Etoposide	Everted sacs of ileum of rats	M-S and S-M transport	Increased A-B P _{app}	(Akht 1995)
	0.3-1000 µM	Digoxin	MDCKII MDR1	Uptake transport	Increased intracellular accum. in a conc. dep. manner. IC ₅₀ = 180 µM	(Gurja 1995)
	0.05-0.5% w/v	Digoxin	Rat everted gut sac model	Uptake transport	Enhanced digoxin accum.	(Corna 2004)
	0.05 and 0.5 w/v	Celiprolol	Rat everted gut sac model	Uptake transport	Enhanced celiprolol accum.	(Corna 2004)
	0.002-1 mg/mL	Paclitaxel	Intestinal segments from rats' ileum	Bi-directional transport (Ussing chamber)	Decreased B-A P _{app} in a conc. dep. manner. Increased A-B P _{app} .	(Varm 1995)
	0.1 and 1 mg/mL	Paclitaxel	Intestinal segments from rats' ileum	In situ single-pass perfusion experiments	Increased intestinal P _{app} .	(Varm 1995)
		Verapamil	P-gp membranes from Sf9	ATPase	Inhibited substrate induced ATPase activity. IC ₅₀ (µM) = 3.18 ± 1.97	(Colln 1995)
		Quinidine	P-gp membranes from Sf9	ATPase	Inhibited substrate induced ATPase activity. IC ₅₀ (µM) = 0.82 ± 0.47	(Colln 1995)
		Progesterone	P-gp membranes from Sf9	ATPase	Inhibited substrate induced ATPase activity. IC ₅₀ (µM) = 3.25 ± 1.29	(Colln 1995)
		Nicardipine	P-gp membranes from Sf9	ATPase	Inhibited substrate induced ATPase activity. IC ₅₀ (µM) = 0.40 ± 0.17	(Colln 1995)
	33.0 µM	Rh 123	Caco-2 monolayers	Bi-directional transport	Increase A-B and decrease B-A P _{app} .	(Colln 1995)
	33.0 µM	Digoxin	Caco-2 monolayers	Bi-directional transport	Increase A-B and decrease B-A P _{app} .	(Colln 1995)

	0.005%	Calcein-AM	NCI/ADR-RES	Calcein-AM efflux	Dose-dependent increase in calcein fluorescence.	(Dong et al., 2005)
		Calcein-AM	MDA-MB-435/LCC6MDR1	Calcein-AM efflux	Dose-dependent increase in calcein fluorescence.	(Dong et al., 2005)
		Talinolol	Caco-2 Cell	Bi-directional	Increased A-B P_{app} .	(Bognár et al., 2005)
Dose of surfactant		Substrate (Dose)		Impact		Reference
EL	1% SMEDDS containing 50% (w/w) surfactant	Etoposide ^a (12 mg/kg)		Increased AUC, C_{max} , and F by 1.7-, 1.3-, and 1.7-fold, respectively.		(Zhao et al., 2014)
RH 40	1% SMEDDS containing 43% (w/w) surfactant	Etoposide ^a (12 mg/kg)		Increased AUC, C_{max} , and F by 1.4-, 1.3-, and 1.4-fold, respectively.		(Zhao et al., 2014)
or oil						
royl	240 mg/kg	Rifampicin ^b		Increased AUC by 1.5-fold, prolonged $t_{1/2}$ by 25%, and decreased CL to 60%		(Ma et al., 2014)
lycerides,	1% (w/v)	Etoposide ^b (4.5 mg/kg)		Increased AUC, C_{max} , and F by 1.8-, 4.7-, and 1.8-fold, respectively.		(Akhtar et al., 2014)
ric	5% (w/v)	Etoposide ^b (4.5 mg/kg)		Increased AUC, C_{max} , and F by 3-, 7-, and 3-fold, respectively.		(Akhtar et al., 2014)
	10% (w/v)	Etoposide ^b (4.5 mg/kg)		Increased AUC, C_{max} , and F by 1.6-, 6-, and 1.6-fold, respectively.		(Akhtar et al., 2014)
lene (40)	8.5 mg/tablet	Rh 123 ^a (1.5 mg/tablet)		Increased AUC by 3.4-fold.		(Föger et al., 2014)

1476

1477

1478 **Table 43:** Impact of nonionic surfactants on intestinal P-glycoprotein in rats.

1479 *In vivo* pre-clinical studies were performed in male ^a: Sprague-Dawley rats, ^b: Wistar albino rats.

1480 Synonyms of surfactants are available in Table 2.

1481	240 mg/kg	Rifampicin ^b	Increased AUC by 1.5-fold, prolonged $t_{1/2}$ by 38%, and decreased CL to 60%.	(Ma et al.
1482	8.5 mg/tablet	Rh123 ^a	Increased AUC by 1.6-fold.	(Föger et
1483	10-25% (v/v)	Digoxin ^a (0.2 mg/kg)	Increased AUC by 1.4-fold, increased C_{max} by 1.4-1.8-fold, increased k_e by 1.4-1.6-fold, and increased F by approx. 1.5-fold.	(Nielsen et
1484	5 and 25% (v/v)	Etoposide ^a (20 mg/kg)	Increased AUC by 1.8-fold, increased C_{max} by 1.5-2.1-fold, CL decreased by half, and increased F by 1.7-fold.	(Al-Ali et
1485	1 and 10% (v/v)	Digoxin ^a (0.2 mg/kg)	Increased AUC by 1.3-1.6-fold and increased C_{max} by 2.5-fold	(Zhang et
1486	10%	Rifampicin ^b (30 mg/kg)	Increased AUC by 1.7-fold and decreased $t_{1/2}$ to 36%.	(Shimomura et al., 2016)
1487	1% SMEDDS containing 50% (w/w) surfactant	Etoposide ^a (12 mg/kg)	Increased C_{max} by 3.5-fold, increased F and AUC by 2.5-fold.	(Zhao et al.)
1488	1 mg/kg	Digoxin ^a (0.25 mg/kg)	Increased AUC by 1.4-fold and decreased t_{max} by 4.5-fold.	(Cornaier et al.)
1489	10%	Colchicine (5mg/kg)	Increased AUC by 4-fold.	(Bittner et al.)
1490	10%	Colchicine (5mg/kg)	Increased AUC by 2-fold.	(Bittner et al.)
1491	50 mg/kg	Paclitaxel ^a (25 mg/kg)	Increased AUC, C_{max} , and F by 6.3-, 3.1-, and 6.4-fold, respectively.	(Varma and Panchagnanam)

Table 34: *In vitro* and *in vivo* impact of nonionic surfactants on breast cancer resistance protein, BCRP, *in vitro* and *in vivo*.

Concentration (conc.), Approximately (approx.), Respectively (resp.), Plasma membrane vesicle of cells containing human ABCG2 (Membrane vesicles BCRP), Clonal isolate derived from the *Spodoptera frugiperda* cell line IPLB-Sf21-AE (Sf9 insect cells), *In Vitro* Diffusion Chamber Method

Surfactants	Conc.	Substrate	Cells/Tissue/ <u>Animal</u>	Assay	Impact of surfactant	Ref.
β-D-glucoside, (C₆-malt)			Membrane vesicles BCRP	Phosphate release measurements	Reduced Pgp ATPase activity. $K_2 = 4.6 \cdot 10^3 \mu\text{M}$	(Xu)
glycol, ene (9) lauryl 99	0.05% and 0.075%	Sulfasalazine	Rat intestinal membrane	In vitro DCM	Decreased <u>S-M B-A</u> transport.	(Sav 2018)
	0.05 %	Sulfasalazine	WT male Wistar rat	<i>In situ</i> CLM	Increased AUC and C _{max} by 1.45 and 1.4- folds, resp.	(Sav 2018)
	0.1 %	Sulfasalazine	WT male Wistar rat	<i>In situ</i> CLM	Increased AUC and C _{max} by 2.2 and 2.1-folds, resp.	(Sav 2018)
ene (4) lauryl 4, Brij® L4	50 and 100 μM	Mitoxantrone	MDCKII BCRP	Uptake transport	Increased the uptake by approx. 1.7-fold.	(Yar 2007)

~~(*In vitro* DCM), *In situ* closed-loop method (*In situ* CLM), Wild type (WT), Sprague-Dawley (SD),~~

~~Synonyms of surfactants are available in Table 2.~~

polyoxyethylene er EL	0.01% and 0.05%	Sulfasalazine	Rat intestinal membrane	<i>In vitro</i> DCM	Decreased S-M B-A transport.	(Sav 2018
	0.1 %	Sulfasalazine	WT male Wistar rat	<i>In situ</i> CLM	Increased AUC and C _{max} by 1.8 and 2.3-fold, resp.	(Sav 2018
	50 µM	Mitoxantrone	MDCKII BCRP	Uptake transport	Increased the uptake by approx. 1.4-fold.	(Yar 2007
	6.25-100 nM	Scutellarin	Membrane vesicles of Sf9 BCRP	Uptake transport	Increased the uptake in a conc. dependent manner.	(Xia 2016
	1 and 5 µg/mL	Scutellarin	MDCKII BCRP	Bi-directional transport	Increased A-B and decreased B-A P _{app} .	(Xia 2016
	5 µg/mL	Scutellarin	WT Male SD rats		Increased AUC and C _{max2} by 1.6 and 1.9-folds, resp.	(Xia 2016
	0.025 and 0.05% (w/v)	Topotecan	Caco-2 cell monolayers	Bi-directional transport	Increased A-B and/or decreased B-A P _{app} .	(Sav 2018
methyl-β-D- side, (Cymal-1)	0.05 % (w/v)	Topotecan	WT male Wistar rat	<i>In situ</i> CLM	Increased AUC 2.3-folds.	(Sav 2018
			Membrane vesicles BCRP	Phosphate release measurements	Reduced P-gp ATPase activity. K ₂ = 1.51 10 ⁴ µM	(Xu 2018
	0.075%	Sulfasalazine	Rat intestinal membrane	<i>In vitro</i> DCM	Decreased S-M B-A transport.	(Sav 2018
	0.025 and 0.05% (w/v)	Topotecan	Caco-2 cell monolayers	Bi-directional transport	Increased A-B P _{app} .	(Sav 2018
	0.025 and 0.05% (w/v)	Topotecan	Caco-2 cell monolayers	Bi-directional transport	Decreased B-A P _{app} .	(Sav 2018
	20 µM	Mitoxantrone	MDCKII BCRP	Uptake transport	Increased the uptake by approx. 1.8-fold.	(Yar 2007
	250 mg kg ⁻¹ (Oral)	Topotecan (Oral)	WT mice		Increased the AUC by 2- folds.	(Yar 2007
ene (8) lauryl yloctaglyol,	20 µM	Topotecan	Everted sacs from WT mice ileum	Transport	Increased the intestinal absorption rate of topotecan.	(Yar 2007
	20 µM	Mitoxantrone	MDCKII BCRP	Uptake transport	Increased the uptake.	(Yar 2009
			Membrane vesicles BCRP	Phosphate release measurements	Reduced Pgp ATPase activity. K ₂ = 6.93 µM	(Xu 2018
	100 and 250 µM	Mitoxantrone	MDCKII BCRP	Uptake transport	Increased the uptake by approx. 1.6-fold.	(Yar 2007
	100 mg kg ⁻¹	Topotecan	WT mice		Increased the AUC by 2-	(Yar 2007

	(Oral) 250 μ M	(Oral) Topotecan	Everted	Transport	fold. Increased the intestinal	2007 (Yan)
	Conc.	Substrate	Cells or animal	Assay	Impact of surfactant	Reference
EL	0.005-0.05% (v/v)	Calcein-AM	MDCKII MRP2	Bi-directional transport	Decreased B-A P_{app} .	(Hank)
	100 μ g/mL	Scutellarin	Membrane vesicles of Sf9 MRP2	Uptake transport	Increased the uptake.	(Li et al)
	0.1%		membrane		transport.	2013
	0.025 and 0.05% (w/v)	Topotecan	Caco-2 cell monolayers	Bi-directional transport	Increased A-B and decreased B-A P_{app} .	(Sav)
	0.05 % (w/v)	Topotecan	WT male Wistar rat	<i>In situ</i> CLM	Increased AUC 2.5-folds.	(Sav)
	100 μ M	Mitoxantrone	MDCKII BCRP	Uptake transport	Increased the uptake by approx. 1.4-fold.	(Yan)
molaurate						2007

Table 5: *In vitro* and *in vivo* impact of nonionic surfactants and co-surfactants on multidrug resistance-associated protein 2 MRP2.

Concentration (Conc.), Respectively (resp.), Membrane vesicles prepared from *Spodoptera frugiperda* (Sf9) insect cells over-expressing human MRP2 (Membrane vesicles of Sf9 MRP2), ATP measurements were performed using ATP luciferin/Luciferase assay, Wild type (WT), Synonyms of surfactants are available in Table 2 and 3.

	100 µg/mL	Scutellarin	Caco-2 cell monolayers	Bi-directional transport	Decrease ER.	(Li et al.)
	0.1-100 µg/mL	Scutellarin	Caco-2 cell monolayers	Bi-directional transport	Decreased ER in a conc. dependent manner.	(Li et al.)
	0.1-100 µg/mL	Scutellarin	Membrane vesicles of Sf9 MRP2	Uptake transport	Increased the uptake in a conc. dependent manner.	(Li et al.)
	6.25-100 nM	Scutellarin	Membrane vesicles of Sf9 MRP2	Uptake transport	Increased the uptake in a conc. dependent manner.	(Xiao et al.)
	1 and 5 µg/mL	Scutellarin	MDCKII MRP2	Bi-directional transport	Decreased B-A P_{app} .	(Xiao et al.)
	5 µg/mL	Scutellarin	WT Male Sprague-Dawley rats		Increased AUC and C_{max} by 1.6 and 1.9-folds, resp.	(Xiao et al.)
EL + 27	100 µg/ml + 100 µg/ml	Scutellarin	Membrane vesicles of Sf9 MRP2	Uptake transport	Increased the uptake.	(Li et al.)
EL +	100 µg/ml + 100 µg/ml	Scutellarin	Membrane vesicles of Sf9 MRP2	Uptake transport	Increased the uptake.	(Li et al.)
RH 40	0.02-0.04% (v/v)	Calcein-AM	MDCKII MRP2	Bidirectional transport	Decreased B-A P_{app} .	(Hank et al.)
	100 µg/mL	Scutellarin	Membrane vesicles of Sf9 MRP2	Uptake transport	Increased the uptake.	(Li et al.)
	100 µg/mL	Scutellarin	Caco-2 cell monolayers	Bi-directional transport	Decreased ER.	(Li et al.)
	0.1-100 µg/mL	Scutellarin	Caco-2 cell monolayers	Bi-directional transport	Decreased ER in a conc. dependent manner.	(Li et al.)
	0.1- 100 µg/mL	Scutellarin	Membrane vesicles of Sf9 MRP2	Uptake transport	Increased the uptake in a conc. dependent manner.	(Li et al.)
	0.1-100 µg/mL	Scutellarin	Caco-2 cell monolayers	Bi-directional transport	Decreased ER.	(Li et al.)
	100 µg/mL	Scutellarin	Membrane vesicles of Sf9 MRP2	Membrane vesicles transport assay	Increased the uptake.	(Li et al.)
	100 µg/mL	Scutellarin	Caco-2 cell monolayers	Bi-directional transport	Decreased ER.	(Li et al.)
	0.1-100 µg/mL	Scutellarin	Membrane vesicles of Sf9 MRP2	Uptake transport	Increased the uptake in a conc. dependent manner.	(Li et al.)
	0.1-10 µg/mL	Scutellarin	Caco-2 cell monolayers	Bi-directional transport	Decreased B-A P_{app} .	(Li et al.)
	100 µg/mL	Scutellarin	Caco-2 cell monolayers	Bi-directional transport	Increased A-B P_{app} .	(Li et al.)
	100 µg/mL	Scutellarin	Membrane vesicles of Sf9 MRP2	Uptake transport	Increased the uptake.	(Li et al.)
	100 µg/mL	Scutellarin	Caco-2 cell	Bi-directional	Decreased ER.	(Li et al.)

	0.1-100 μg/mL	Scutellarin	monolayers Caco-2 cell monolayers	transport Bi-directional transport	Decreased ER.	(Li et al)
	0.1-100 μg/mL	Scutellarin	Sf9 MRP2	Uptake transport	Increased the uptake in a conc. dependent manner.	(Li et al)
	100 μg/mL	Scutellarin	Membrane vesicles of Sf9 MRP2	Uptake transport	Increased the uptake	(Li et al)
	100 μg/mL	Scutellarin	Caco-2 cell monolayers	Bi-directional transport	Decreased ER.	(Li et al)
	0.1-100 μg/mL	Scutellarin	Caco-2 cell monolayers	Bi-directional transport	Decreased ER.	(Li et al)
	10 μg/mL	Baicalcein	MDCKII MRP2	Bi-directional transport	Increased A-B P _{app} and decreased B-A P _{app} .	(Chen)
7, 07, 7	100 μg/mL	Scutellarin	Caco-2 cell monolayers	Bi-directional transport	Decreased ER.	(Li et al)
	0.1-100 μg/mL	Scutellarin	Caco-2 cell monolayers	Bi-directional transport	Decreased ER in a conc. dependent manner.	(Li et al)
	0.1-100 μg/mL	Scutellarin	Membrane vesicles of Sf9 MRP2	Uptake transport	Increased the uptake in a conc. dependent manner.	(Li et al)
	10 μg/mL	Baicalcein	MDCKII MRP2	Bi-directional transport	Decreased B-A P _{app} .	(Chen)
	0.00005- 0.005% (w/w)		MDCKII MRP2	ATP measurements	Decreased ATP levels.	(Batra 2003)
	0.01-0.5% (w/w)	Vincristine	MDCKII MRP2	Uptake transport	Increased intracellular accum. in a conc. dependent manner. Decreased IC ₅₀ value by 6.6 times.	(Batra 2003)
	0.01-0.5% (w/w)	Doxorubicin	MDCKII MRP2	Uptake transport	Increased intracellular accum. in a conc. dependent manner. Decreased IC ₅₀ value by 125 times.	(Batra 2003)
	0.1 % (w/v)		Plasma membranes of MDCKII MRP2	Phosphate release measurements	Decreased V _{max} .	(Batra 2004)
	0.1 % (w/v)	Vincristine	Plasma membranes of MDCKII MRP2	Phosphate release measurements	Decreased V _{max} and increased K _m .	(Batra 2004)
	10 μg/mL	Baicalcein	MDCKII MRP2	Bi-directional transport	Increased A-B P _{app} and decreased B-A P _{app} .	(Chen)
+	10 μg/mL	Baicalcein	MDCKII MRP2	Bi-directional transport	Increased A-B P _{app} and decreased B-A P _{app} .	(Chen)
+	10 μg/mL	Baicalcein	MDCKII MRP2	Bi-directional transport	Increased A-B P _{app} and decreased B-A P _{app} .	(Chen)
+	10 μg/mL	Baicalcein	MDCKII MRP2	Bi-directional	Increased A-B P _{app} and	(Chen)

5				transport	decreased B-A P_{app} .	
5	10 $\mu\text{g/mL}$	Baicalcein	MDCKII MRP2	Bi-directional transport	Increased A-B P_{app} and decreased B-A P_{app} .	(Chen
0	0.05% (v/v)	Calcein-AM	MDCKII MRP2	Bi-directional transport	Decreased B-A P_{app} .	(Hank
	0.01-0.05% (v/v)	Calcein-AM	MDCKII MRP2	Bi-directional transport	Decreased B-A P_{app} .	(Hank
col	100 $\mu\text{g/mL}$	Scutellarin	Membrane vesicles of Sf9 MRP2	Uptake transport	Increased the uptake.	(Li et a
er	100 $\mu\text{g/mL}$	Scutellarin	Caco-2 cell monolayers	Bi-directional transport	Decreased ER.	(Li et a
	0.1 and 1 $\mu\text{g/mL}$	Scutellarin	Caco-2 cell monolayers	Bi-directional transport	Decrease ER.	(Li et a
	0.1 and 100 $\mu\text{g/mL}$	Scutellarin	Membrane vesicles of Sf9 MRP2	Uptake transport	Increased the uptake in a conc. dependent manner.	(Li et a

1517

1518

1519

1520 **Table 6:** Nonionic surfactants inhibited solute carriers (SLCs) *in vitro*.

1521

1522

1523 ~~IC₅₀ were estimated from uptake transport assay. For Regev et al. 2002, impact of surfactant on bi-~~
1524 ~~directional transport assay was shown. 1-methyl-4-phenylpyridinium acetate (MPP⁺), Monocarboxylic~~
1525 ~~acid transporter (MCT, SLC16A1), Organic cation transporter 1 (OCT1, SLC22A1), (OCT2,~~
1526 ~~SLC22A2), (OCT3, SLC22A3), Peptide transporter 1 (PEPT1, SLC15A1), (PEPT2, SLC15A2),~~
1527 ~~Organic anion transporting polypeptide 1A2 (OATP1A2, SLC21A3), (OATP2B1, SLC21A9). Human~~
1528 ~~embryonic kidney cells stably transfected with OATP1A2 (HEK OATP1A2), or with OATP2B1 (HEK~~
1529 ~~OATP2B1), Chinese hamster ovary cells stably transfected with rbOCT1(CHO-K1 rbOCT1), Madin-~~
1530 ~~Darby canine kidney cells stably transfected with OCT1-3 (MDCKII OCT1-3), or with PEP2 (MDCKII~~
1531 ~~PEPT2). Synonyms of surfactants available in Table 2 and 5.~~

1532

1533
1534
1535
1536
1537
1538
1539
1540

1541

Nonionic surfactant	Transporter SLC	Substrate	Cells	IC ₅₀ Impact of surfactant
Solutol® HS 15	OATP1A2	Estrone-3-sulfate	HEK OATP1A2	0.0074%
	OATP1A2	Taurocholate	HEK OATP1A2	0.0041%
	OATP2B1	Estrone-3-sulfate	HEK OATP2B1	0.011%
	OATP2B1	Bromosulfophthalein	HEK OATP2B1	0.00095%
	OCT1	MPP ⁺	MDCKII OCT1	0.008%
	OCT2	MPP ⁺	MDCKII OCT2	0.046%
	OCT3	MPP ⁺	MDCKII OCT3	0.019%
	PEPT2	Glycyl sarcosine	MDCKII PEPT2	0.014%
Cremophor® EL	OATP1A2	Estrone-3-sulfate	HEK OATP1A2	0.00054%
	OATP1A2	Taurocholate	HEK OATP1A2	0.00034%
	OATP2B1	Estrone-3-sulfate	HEK OATP2B1	0.0011%

1542
1543
1544
1545
1546
1547
1548
1549
1550

	OATP2B1	Bromosulfophthalein	HEK OATP2B1	0.0098%
	OCT1	MPP ⁺	MDCKII OCT1	0.019%
	OCT2	MPP ⁺	MDCKII OCT2	0.46%
	OCT3	MPP ⁺	MDCKII OCT3	9.77%
	PEPT2	Glycyl sarcosine	MDCKII-PEPT2	0.16%
	MCT	Benzoic acid	Caco-2 cells	Decreased A-B P _{app} of the substrate in a concentration dependent manner.
Kolliphor® P 188, Poloxamer 188	OCT3	MPP ⁺	MDCKII OCT3	0.024%
Kolliphor® P407	OCT1	MPP ⁺	MDCKII OCT1	1.85 %
Polysorbate 20	OCT1	MPP ⁺	MDCKII OCT1	0.002%
	OCT1	MPP ⁺	CHO-K1 rbOCT1	85 ± 1.12 µg/ml
	OCT2	MPP ⁺	MDCKII OCT2	0.033%
	OCT2	MPP ⁺	CHO-K1 rbOCT2	295 ± 1.48 µg/ml
	OCT3	MPP ⁺	MDCKII OCT3	0.011%
	PEPT2	Glycyl sarcosine	MDCKII-PEPT2	0.005%
Polysorbate 60, Tween® 60, Polyoxyethylene (20) sorbitan stearate	OCT1	MPP ⁺	CHO-K1 rbOCT1	50 ± 1.26 µg/ml
	OCT2	MPP ⁺	CHO-K1 rbOCT2	42 ± 1.15 µg/ml
Polysorbate 80	OCT1	MPP ⁺	MDCKII OCT1	0.0007%
	OCT1	MPP ⁺	CHO-K1 rbOCT1	106 ± 1.20 µg/ml
	OCT2	MPP ⁺	MDCKII OCT2	0.039%
	OCT2	MPP ⁺	CHO-K1 rbOCT2	185 ± 1.20 µg/ml
	OCT3	MPP ⁺	MDCKII OCT3	0.011%
	PEPT2	Glycyl sarcosine	MDCKII PEPT2	0.037%
	PEPT1	Glycyl sarcosine	Caco-2 cells	Decreased A-B P _{app} of the substrate in a concentration dependent manner.

1551

1552

1553 **Declaration of interest**

1554 The authors do not have any conflict of interest to report.


1555

1556 **Author contribution**

1557 Writing - original draft: AAAA, CUN, and RBN. Writing - review & Editing: AAAA, RBN, BS, RH and CUN. Final
1558 approval of the version submitted: AAAA, RBN, BS, RH and CUN.

1559

1560



1561