

The impact of ozone exposure, temperature and CO₂ on the growth and yield of three spring wheat varieties

Hansen, Emilie Marie Øst; Hauggaard-Nielsen, Henrik; Launay, Marie; Rose, Paul; Mikkelsen, Teis Nørgaard

Published in:
Environmental and Experimental Botany

DOI:
[10.1016/j.envexpbot.2019.103868](https://doi.org/10.1016/j.envexpbot.2019.103868)

Publication date:
2019

Document Version
Peer reviewed version

Citation for published version (APA):
Hansen, E. M. Ø., Hauggaard-Nielsen, H., Launay, M., Rose, P., & Mikkelsen, T. N. (2019). The impact of ozone exposure, temperature and CO₂ on the growth and yield of three spring wheat varieties. *Environmental and Experimental Botany*, 168, Article 103868. <https://doi.org/10.1016/j.envexpbot.2019.103868>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

1 The impact of ozone exposure, temperature and CO₂ on the growth and yield of three spring wheat
2 varieties.

3

4 Emilie M. Ø. Hansen^a, Henrik Hauggaard-Nielsen^a, Marie Launay^b, Paul Rose^b, Teis N.
5 Mikkelsen^{c*}

6

7 ^a Department of People and Technology, Research group Environment, Energy, Transport,
8 Regulation, Innovation and Climate Policy, Universitetsvej 1, DK-4000 Roskilde
9 Roskilde University

10 ^b Unité Climat Sol et Environnement, INRA, Site Agroparc, 84914 Avignon Cedex 9, France

11 ^c Department of Environmental Engineering, Section Air, Land and Water Resources, Technical
12 University of Denmark, 2800 Kgs. Lyngby, Denmark

13

14 *Corresponding author: Teis N. Mikkelsen, DTU, temi@env.dtu.dk

15

16

17 ABSTRACT

18 When assessing potentials for crop production under future climatic conditions, multiple
19 environmental parameters need to be included. An increase in carbon dioxide [CO₂], higher
20 temperatures, and regional changes in tropospheric ozone [O₃] will influence the growth responses
21 of existing crop species and varieties. Ozone is phytotoxic and a plant stressor at current
22 concentrations, reducing yields worldwide, but possible interactions with changes in other abiotic
23 factors have been considered very little. In this study, we have used eight combinations: two levels
24 of temperature, two levels of [CO₂], and three [O₃] exposure regimes to assess the impact of
25 medium-to-high ozone concentrations (80-100 ppb) on wheat growth when other abiotic factors
26 change. Two modern spring wheat varieties (KWS Bittern and Lennox) and a landrace variety
27 (Lantvete) were grown to maturity in climate chambers. We examined plant performance during
28 growth as development rate, rate of photosynthesis, stomatal conductance, water use, and at harvest
29 as total aboveground dry matter and grain yield.

30 All three varieties lost yield in all treatments compared to the ambient treatment that had the
31 following settings: Lowest temperature, ambient [CO₂], and very low [O₃]. For episodic ozone
32 exposure in the ambient or high [CO₂] and high-level temperature treatment, the yield losses were
33 18 and 25%, respectively, for KWS Bittern; 44 and 34% for Lennox; and 16 and 37% for Lantvete.
34 The yields of the modern varieties are significantly higher than the landrace variety in two out of
35 eight treatments, although they are higher by weight in seven of the eight treatments. The landrace
36 variety's fraction loss from its highest grain yield in the ambient treatment to the high-[CO₂]-and
37 high-level-temperature-treatments was smaller than the modern varieties', showing a comparably
38 higher degree of plasticity of performance. Current crop varieties might be more sensitive to ozone

39 than older varieties, emphasizing the need of future breeding programs to expand the gene pool to
40 provide more climate robust crops.

41

42 Keywords

43 Climate change, air pollution, ecophysiology, multifactorial design, ozone episodes

44

45

46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70

1. Introduction

The changes in climate relating to the climate scenarios' emission of carbon dioxide include temperature increase, increased frequency of extreme weather events, regional aggravation of current challenges and addition of new challenges (IPCC, 2014) influencing plant growth in natural and agricultural ecosystems (Albert et al., 2011; Pleijel et al., 2018). These factors will not change individually and their impact on ecosystems is increasingly investigated in multifactorial settings (Ingvordsen et al., 2015; Langley and Hungate, 2014; Namazkar et al., 2016). When addressing responsiveness of existing crop species and varieties grown under future climatic conditions, multiple environmental parameters must be included (Frenck et al., 2013; Vázquez et al., 2017). Both temperature and CO₂ are well-known parameters influencing the growth and development of plants. Temperature influence amongst others enzyme activity and water relations of the plant and as a single factor changing, temperature can be both too low and too high to sustain optimal plant growth, the optimal temperature range depend on the genetics of the variety (Albert et al., 2011; Frenck et al., 2011; Gibson and Paulsen, 1999). Being the substrate for photosynthesis, CO₂ availability can be a limiting factor at ambient concentrations meaning that CO₂-increases enhance yields (AbdElgawad et al., 2015; Cure and Acock, 1986). However, simultaneous combinations of changes in climatic factors may not result in additive effects (Clausen et al., 2011; Dieleman et al., 2012; Shaw et al., 2002).

In addition, the air pollutant, ozone, which is causing yield loss under the current climatic conditions, may interact with the changing levels of other abiotic factors, and thus further impact temporal and spatial plant growth patterns and yield (Pleijel et al., 2018). Exposure to elevated ozone typically results in suppressed photosynthesis, accelerated senescence, decreased growth and lower yields (Booker et al., 2009). However, there are major knowledge gaps when assessing the threat that ozone plays (Ainsworth et al., 2012; Fuhrer and Booker, 2003).

71 Ozone is readily formed from available precursors in the presence of UV-radiation. The precursors
72 are natural or anthropogenic volatile organic compounds (VOCs) and nitrogen oxides (NO_xs)
73 (Finlayson-Pitts and Pitts, 1993). The precursors may be carried long distances and result in both
74 peak ozone formations far from the precursor formation/emission and in a regional increase of
75 background concentration (Monks et al., 2015). Efforts to reduce precursor emission contribute to
76 regional control of ozone levels (Derwent et al., 2018). Precursors may also build up during periods
77 of low radiation, i.e. in winter, and result in a peak ozone period in spring when radiation increases
78 (Munir et al., 2013). In stable, high-pressure conditions, ozone concentration may build up to create
79 a peak ozone episode, whilst the ozone molecules may disperse into adjacent areas in more windy
80 conditions (Kleanthous et al., 2014).

81 While everyday ozone concentrations reflect regional and local activities and conditions,
82 background ozone levels have been steadily rising with anthropogenic activity over the last century
83 (Lamarque et al., 2010). The background ozone concentration, which is a result of the mixing of air
84 and the emitted precursors, has also been projected to worsen in the future depending on possible
85 climate scenarios (Vingarzan, 2004) due to emissions from industrialization and other human
86 activities (Wild et al., 2012). Ozone concentration varies with the time of day and season, based on
87 precursor availability and regional weather, and plants may therefore be exposed to periods of
88 ozone peaks as well as the increase in background ozone.

89 The cost of the plant's defense mechanism against ozone is that fewer carbon substances are
90 available for allocation to build biomass, both above and below ground (Calatayud et al., 2011).
91 Thus, ozone limits plant growth and agricultural yields, and Mills et al. (2018) estimate a global
92 yield gap of 7.1% in wheat at the background levels and peak ozone incidents prevailing under
93 current climatic conditions. Effects of changes in climatic factors and levels of ozone concentration

94 can be synergistic or antagonistic interactions with the defense mechanism influencing plant growth
95 and yield in unknown ways (Larsen et al., 2011).

96 In the present study, wheat (*Triticum aestivum*) is used as model crop selected for its global
97 importance as staple food in many countries (Wrigley, 2009). Wheat yields have increased vastly
98 with breeding, research, development, and intensification of production for the last 50 years,
99 however, the rate of yield increase has diminished in the past 30 years (Brisson et al., 2010). The
100 decline in soil carbon content may be one explanation (Brisson et al., 2010), and the combined
101 increase of CO₂-concentration and temperature another (Dieleman et al., 2012). While an increase
102 in [CO₂] has a growth promoting effect on many plants as the substrate of photosynthesis, and
103 higher temperatures increase the rates of activity in many biochemical processes, the combination
104 of CO₂-enrichment and temperature increase may have a less than additive effect on the plant
105 biomass (Dieleman et al., 2012; Shaw et al., 2002). As ozone is known to reduce the yield of cereals
106 (Broberg et al., 2015), the increase in tropospheric ozone with anthropogenic activity (Lamarque et
107 al., 2010) suggests that the increasing ground-level ozone concentration cannot be ignored and
108 might be a threat to future food production through interactions with other climate factors.

109

110 The aim of this study was to investigate eight climate treatments reflecting current conditions and
111 future scenarios. From seed to maturity, the wheat varieties were regularly evaluated for a range of
112 growth indicators to learn how the combinations of abiotic factors influenced the development and
113 yield of the wheat plants. The following four hypotheses were tested: i) Yields of modern varieties
114 are higher than yields of the old variety; ii) Episodic ozone exposure at Zadoks' growth stages
115 (ZS)31-69 is as injurious to wheat yields as full-time ozone exposure; iii) The combined effect of
116 elevated [CO₂] and elevated temperature, which stimulate plant growth less than expected by the
117 elevation of each factor individually is further aggravated by exposure to elevated ozone

118 concentrations; iv) Differences in yields are reflected in the way growth parameters respond to the
119 treatments during growth.

120

121 2. Materials and methods

122 2.1. Climate chamber experiment

123 The study was done in climate chambers that provided a controlled environment and uniform
124 conditions, thus eliminating other potentially interacting parameters. The RERAF phytotron (Risø
125 Environmental Risk Assessment Facility, Technical University of Denmark, Risø, Denmark)
126 consists of six gastight chambers sized 6 x 4 x 3 m, providing detailed control of temperature,
127 [CO₂], air humidity, light, and ozone concentration and exposure duration. Chamber air was
128 exchanged at a rate of 4 m³ h⁻¹, resulting in a complete exchange of chamber air every 18 hours and
129 a wind speed of less than 0.6 m s⁻¹ (measured in earlier experiments). The lighting produced 313-
130 389 μmol depending on chamber. The facility's computational system logged treatment parameters
131 in high resolution. For detailed description of the physical facility see Frenck et al. (2011), Clausen
132 et al. (2011) & Ingvordsen et al., (2015).

133

134 2.2. Common input

135 The studied spring wheat varieties included two modern varieties: *Lennox* (Saaten-Union) used in
136 southern France and *KWS Bittern* (DanishAgro) used in Denmark, and one landrace variety: the
137 Swedish *Lantvete* (Nordic Genetic Resource Center). All with a life span of approx. 3-4 months.
138 Twelve seeds of the spring wheat varieties tested were sown in 11 L pots filled with 4 kg of
139 sphagnum (Pindstrup Substrate No. 4, Pindstrup Mosebrug A/S, Ryomgaard, Denmark) and
140 reduced to eight plants after germination, corresponding to ~165 plants m⁻². Each variety was
141 represented in each treatment with five pots. No additional nutrients were added to the pots as the
142 sphagnum was nutrient enriched. The watering water was tap water.
143 CO₂ was provided by [AGA A/S \(Linde Worldwide, Copenhagen, Denmark\)](#) and ozone was
144 generated by the use of UV Pro 550 A ozone generators ([Crystal Air Products and Services,](#)

145 [Langley, BC, Canada](#)). The UV-lamps in the ozone generators were at times partially shaded with
146 cardboard to reduce the amount of ozone generated.

147 The experiment lasted from March 6th to June 25th 2016.


148

149 2.3. Growth conditions

150 2.3.1. Treatments

151 Eight treatments were tested in total. They were selected among possible combinations of two
152 present and future temperature levels (19/12 °C or 24/17 °C, both levels simulating days (16h) that
153 are warmer than nights (8h)), two concentrations of CO₂ (400 and 700 ppm), and one of three ozone
154 regimes: 1) no ozone enrichment, 2) episodic ozone exposure (.EpO3), and full-time ozone
155 exposure (.O3) (Fig. 1). The ‘no ozone’ ozone-enrichment treatments act as filtered air treatments,
156 as ozone concentrations in the climate chambers ‘background levels’ (5.9 ± 0.5 and 7.2 ± 1.7 ppb,
157 see Table 1) are lower than the outside average ozone concentration near the RERAF phytotron
158 with an average of 40.4 ppb, and maximum ozone one-hour concentrations between 70,9-86,6 ppb
159 (calculated with data from 2013-16 (May-July; 8am - 20pm), from station DK0012R, Danish
160 Center for Environment and Energy, Aarhus University). For both the episodic and full-time ozone
161 exposure, the ozone concentration target was 80-100 ppb during the day, thus resulting in 16 hours
162 of daytime ozone exposure, at night the ozone concentration was as the chamber background
163 measured in the ‘no ozone enrichment’ treatments. Unlike the full-time exposure treatments, which
164 started at sowing, the ozone exposure of the .EpO3 treatments began when the Lennox variety
165 reached Zadoks’ developmental stage 31 (ZS31 - first node detectable) and ended when Lennox
166 reached stage 69 (ZS69 - anthesis complete) (Zadoks and Board, 1974). Thus, the number of days
167 of ozone exposure varied as plant development varied depending on climate. Relative humidity was
168 kept at 70/55% (day/night) for all treatments.

169 The following eight treatments were represented in this experiment: A, A.EpO3, A.O3, C.EpO3,
 170 CT, CT.EpO3, CT.O3, and T.EpO3 as shown in Fig. 1. A denotes ‘ambient conditions’ with
 171 temperatures that correspond to good growing conditions during the growth season of spring wheat
 172 in Denmark, and a CO₂-concentration corresponding to present average atmospheric concentrations;
 173 C denotes an elevated level of [CO₂], and T denotes a higher temperature.



174 Fig. 1: The selected climate combinations are named A, A.EpO3, A.O3 (ambient [CO₂], lower
 175 temperature settings and no addition ()), episodic addition (.EpO3) or chronic addition of ozone
 176 (.O3)) and CT, CT.EpO3, CT.O3 (high [CO₂] (C), higher temperature settings (T), and no addition
 177 ()), episodic addition (.EpO3) or chronic addition of ozone (.O3)). Two treatments, C.EpO3 and
 178 T.EpO3 have episodic addition of ozone (EpO3) and either higher [CO₂] or higher temperature
 179 settings. The aim for concentration of ozone in the chambers was 80-100 ppb. The episodic
 180 exposure started at Zadoks' growth stage ZS31 and ended at stage ZS69. A.EpO3 and CT.EpO3
 181 were transferred to the corresponding full-time ozone exposure chamber for ozone exposure and
 182 returned after exposure, while C.EpO3 and T.EpO3 received ozone exposure in their own chambers
 183 for the episode from ZS31-ZS69.
 184
 185

186 2.3.2. Watering regime

187 Plants were watered three times a week to provide a full supply of water; the warm treatment plants
 188 were, by design and by consumption, given more water than ambient treatment plants and all plants
 189 got increasingly more water as they grew. Pots were weighed before and after watering to ensure

190 the same amount of water was accessible in the treatment regardless of the pot's previous
191 consumption.

192

193 2.4. Potential chamber gradient effects

194 To minimize influences from potential gradients of treatment factors or chamber microclimates, the
195 tables in each chamber were moved to different positions in the chamber following each watering.
196 Also, once a week we moved the treatments between the chambers to reduce the risk of intrinsic
197 traits of one chamber to manifest in any one treatment.

198

199 2.5. Process values and measurements of plant growth parameters

200 Process values of treatment parameters such as relative humidity, CO₂ concentration, and
201 temperature were logged by a data collection system several times per minute. The ozone
202 concentration was monitored sequentially, i.e. twice every hour. Photosynthesis, stomatal
203 conductance, transpiration, and other parameters were measured once or twice a week throughout
204 the experiment using two LI-6400 portable photosynthesis systems (LICOR, 2004). The LiCor light
205 was set to 1500 μmol and the other settings followed the treatment parameters. The gas
206 exchange measurements were done at the second leaf, i.e. at any time the youngest fully expanded
207 leaf, representing throughout the life time of the wheat plants the most productive leaf, and hence
208 the productivity of the plant as a whole as the most ozone-exposed leaves as well as senescence will
209 cause main photosynthesis to take part in other leaves.

210 Water consumption was recorded by weighing the individual pots before and during watering; the
211 plants' heights were measured; the development stage assessed according to Zadok's growth stages
212 (Zadoks and Board, 1974) and noted. Ozone was not switched off during measurements and
213 watering.

214 Leaf discoloration was assessed at four different dates following the initiation of ozone exposure in
215 the episodic ozone exposure treatments (starting at 42 and 43 days after sowing). The ratio of green
216 leaf area to total leaf area on the third leaf was assessed by image processing photos of the
217 harvested leaf with the software ImageJ (Schneider et al., 2012). At sampling one leaf in each
218 treatment was harvested.

219 Grain and aboveground biomass were weighed at the end of the experiment. Gluten index was
220 determined by the ICC 155 procedure by the [Nordic Seed Laboratory Services](#).

221

222 2.6. Ozone treatment recap, fluxes and plant ozone uptake.

223 The ozone flux depends on the ozone concentration in the air, the degree of stomatal opening
224 (which, among other factors, depends on water status and CO₂ availability), and the ozone
225 molecules' resistance to cross the boundary layer of the leaves. As the latter was not measured, the
226 resistance part of the equation is left out and any assumptions would only reflect the similarity of
227 wind conditions in the chambers. Thus, ozone flux is the product of the hourly ozone concentrations
228 (in nmol m⁻³) and the stomatal ozone conductance (H₂O mol m⁻² s⁻¹). The stomatal conductance was
229 measured approximately six times in each variety of every treatment, and the values for the days
230 between measurements were found by linear interpolation. Ozone conductance was found by the
231 ratio of molecule size O₃ to H₂O (0.66) and converted from mol m⁻² s⁻¹ into m s⁻¹ (Monks et al.,
232 2015). From the obtained values for fluxes of ozone into the plant, the accumulated plant uptake of
233 ozone above a threshold of 6 was aggregated, giving a unit of mmol O₃ m⁻² PLA (Projected Leaf
234 Area). The threshold of 6 mmol O₃ reflects the sensitivity of wheat plants (ICP Vegetation, 2017).
235 The data logging system occasionally failed to log ozone data, and thus data gap filling has been
236 necessary for the summation of ozone into the plants. Gap filling for single or a few missing data

points was done using the average of the 12 closest measurements. The average of daytime ozone concentration (ppb) in treatment and as background can be found in Table 1.

Table 1. The average daytime concentration of ozone in the eight different treatments with ozone levels in ppb and i) No ozone treatment equivalent to background concentration in the chambers, ii) daily average of episodic ozone exposure and iii) average of chronic ozone exposure. Averages based on ~32 recordings of ozone concentration per growing day. Further explanation of treatment abbreviations see Fig. 1.

Treatment	Episodic relocation		No ozone	Episodic ozone	Chronic ozone
	Base treatment	Exposure treatment			
A			6.4 ±2.1	-	
A.EpO3	A	A.O3	6.4 ±2.1	84.5 ±28.1	
A.O3					78.8 ±32.4
C.EpO3			6.8 ±1.3	82.7 ±35.2	
CT			5.9 ±0.5	-	
CT.EpO3	CT	CT.O3	5.9 ±0.5	80.1 ±35.9	
CT.O3					88.5 ±22.0
T.EpO3			7.2 ±1.7	98.9 ±22.5	


2.7. Data analysis

Data were statistically analyzed using the statistical software R or Microsoft Excel and tested with Breusch-Pagan test, pairwise t-tests and anovas. Levels of $p \leq 0.05$ were considered significant. Models that were tested followed the formula of e.g. Grain Yield = γ (Treatment_i · Variety_i) + e_i. Different physiological responses, such as biomass or photosynthesis were tested for dependence on the climate treatments or elements thereof, i.e. ozone on/off, climates relating to the CO₂/temperature combinations (e.g. Tables 4 and 6).

255
256
257
258
259
260

3. Results

The results are consequently displayed ‘per pot’ to underline the phytotron experiment design. Nevertheless, the grain yield range of 30.8 ± 1.0 to 55.7 ± 1.1 g pot⁻¹ (Fig. 2) converts to 5.0 ± 0.2 to 12.1 ± 0.2 tons ha⁻¹. Unless otherwise indicated, results are displayed with standard errors.



261
262
263
264
265
266
267
268


Fig. 2. Grain yields with standard errors, sorted from the highest to the lowest; blue squares, purple triangles, and green circles indicate if a variety is KWS Bittern, Lennox or Lantvete, respectively. Colored bars indicate groups of no significant difference. Grey columns indicate how many other treatment*varieties this treatment*variety is significantly different from (p≤0.05). n=5. Further explanation of treatment abbreviations see Fig. 1.

3.1. Experimental levels of treatment parameters.

Treatment parameters are reported with standard deviations, TP = X ±SD, each average is based on 10-15 recordings per growing day or night, the data is not shown. The average temperatures in the

low level temperature treatments (set points 19/12 °C) were 19.3 ± 1.1 and 12.1 ± 0.7 °C, and 24.0 ± 0.3 and 17.1 ± 0.9 °C (for set points 24/17 °C) in the high level temperature treatments. The average CO₂ concentration in the low level CO₂ treatments (set point 400 ppm) were 540 ± 37 , and 724 ± 57 ppm in the high level CO₂ treatments (set point 700 ppm). The average relative humidity with set points of 55/70% day/night, were 54 ± 3 and 69 ± 4 % respectively.

The ambient treatment pots were initially watered to a total pot weight of 5200 g to 7000 g when they consumed the most, and the warmer treatments were watered from 5400 g to 7200 g accordingly. When irrigated based on water consumption, this corresponded to a daily average in all the growth period of 8 ± 1 mm in cold treatments and 9 ± 1 mm in warm treatments. The growth dependent water consumption distribution of the treatment*varieties consuming most (Lantvete T.EpO3) and least (Lennox C.EpO3) can be seen as their daily average in Fig. 3.



284 Fig. 3. Water consumption in mm in the treatments consuming most and the least water, Lantvete
285 T.EpO3 (orange circles) and Lennox C.EpO3 (green triangles). n=5, with standard errors. The
286 horizontal bars indicate the daily average for each treatment.

287

288 3.2. Grain yields

289 Grain yields were significantly influenced by the main effects of treatment, variety and interactions
290 between them (p-values of resp. $2.2\text{E-}16$, $1.11\text{E-}11$, and $9.55\text{E-}05$). Elevated temperatures
291 outcompete other effects (Table 2), whereas at ambient temperatures i) ozone significantly reduces
292 yield (except for Lennox A.EpO3) and ii) elevated CO₂ mitigates effects of episodic ozone
293 exposure EpO3, only in the Lantvete variety. On average, across all treatments, yields of the
294 landrace variety, Lantvete, were significantly lower than the modern varieties, $p = 0.0013$ (KWS
295 Bittern) and $p = 0.00012$ (Lennox). Similarly, across the varieties, treatment A had significantly
296 higher yields than all other treatments ($p < 0.01$), and treatment C.EpO3 had significantly higher
297 yields than treatment A.O3 ($p = 0.0004$) and CT.O3 ($p = 0.02$). There were interactions between the
298 varieties and treatments which led to significant differences between some treatment*variety-
299 combinations, e.g. A Lennox, CT.EpO3 Bittern and A.O3 Lantvete (Fig.2). Within a treatment, the
300 varieties only showed a significant difference in A.EpO3, where Lantvete was different from both
301 Lennox and KWS Bittern, and in CT, where Lantvete was different from Lennox but not from KWS
302 Bittern. All three varieties lost grain yield relative to treatment A, but the losses were differently
303 distributed between the treatments without any patterns between losses of grain yield and biomass
304 to explain loss or gain of harvest index (Table 2). A high harvest index (ratio grain yield mass to
305 total aboveground biomass (grain and vegetative biomass)) indicates a high-yielding variety
306 (Peltonen-Sainio et al., 2008). Modern varieties have been bred to be high-yielding and as such
307 have high harvest indexes, but the production of non-reproductive biomass may serve other
308 purposes, e.g. an increased resilience capacity towards biotic and abiotic stresses (Dolferus, 2014).
309 Lantvete had the highest biomass production, which led to lower harvest indexes than the modern

310 varieties (Table 2). In the colder treatments, Lantvete's harvest index decreased with ozone
311 exposure. In warmer treatments, Lantvete's harvest indexes increased compared to the colder
312 treatments, but ozone exposure diminished that increase. For Lennox and KWS Bittern, the harvest
313 index decreased with full-time ozone in the colder treatments and in general in the warmer
314 treatments (Table 2). Lennox lost the same percentage of biomass in the colder treatments
315 regardless of ozone exposure and elevation of [CO₂]. In the warm T.EpO3, Lennox lost the most
316 biomass of all treatments. However, it lead to an increase in harvest index. KWS Bittern lost the
317 most biomass in the warmer CO₂-enriched treatment with no ozone, CT; however the grain yield
318 loss in that treatment was not the most severe and there was an increase in harvest index.


319 Table 2. Grain yield change, Aboveground DM (grain and vegetative dry matter) and change,
320 Harvest Index and change relative to the best yielding treatment A. For grain yield, see Fig. 2.
321 Harvest Index (HI) is grain yield divided by Aboveground DM. Comparison within same variety,
322 n=5. Star symbol indicate if a change relative to treatment A is statistically significant. Further
323 explanation of treatment abbreviations see Fig. 1.

Variety	Treatment	Grain yield	Aboveground DM		Harvest Index		Change (%)
		Change (%)	g pot ⁻¹	Change (%)	HI		
KWS	A	0 ±1	76.4 ±1.9	0 ±2	0.41 ±0.006		0 ±2
Bittern	A.EpO3	-18 ±1*	62.8 ±1.7	-18 ±2	0.44 ±0.005		0 ±2
	A.O3	-35 ±2*	66.0 ±1.9	-14 ±3	0.35 ±0.005		-16 ±2
	C.EpO3	-14 ±1	65.8 ±1.5	-14 ±2	0.44 ±0.004		0 ±2
	CT	-28 ±5*	52.6 ±5.3	-31 ±7*	0.44 ±0.013		3 ±3
	CT.EpO3	-25 ±3*	62.1 ±6.0	-19 ±8	0.42 ±0.014		-4 ±4
	CT.O3	-32 ±2*	55.3 ±2.0	-28 ±3*	0.4 ±0.008		-3 ±2
	T.EpO3	-32 ±5*	61.6 ±8.1	-19 ±11	0.42 ±0.039		-8 ±9
Lantvete	A	0 ±2	104.1 ±3.7	0 ±4	0.3 ±0.005		0 ±2
	A.EpO3	-44 ±4*	84.6 ±2.6	-19 ±3	0.27 ±0.015		-15 ±5
	A.O3	-45 ±2*	82.5 ±2.9	-21 ±3*	0.27 ±0.003		-14 ±2
	C.EpO3	-26 ±3	92.2 ±4.1	-29 ±18	0.31 ±0.013		-3 ±4
	CT	-39 ±3*	66.6 ±1.6	-36 ±2*	0.34 ±0.011		6 ±4
	CT.EpO3	-34 ±4*	72.9 ±1.4	-30 ±1*	0.34 ±0.011		6 ±4
	CT.O3	-36 ±3*	79.0 ±2.8	-24 ±3*	0.31 ±0.013		-2 ±4
Lennox	T.EpO3	-33 ±3*	70.7 ±1.9	-32 ±2*	0.34 ±0.009		9 ±3
	A	0 ±1	71.8 ±2.7	0 ±4	0.44 ±0.004		0 ±1
	A.EpO3	-16 ±1	60.0 ±0.8	-16 ±1	0.44 ±0.005		1 ±2
	A.O3	-31 ±3*	61.0 ±0.7	-15 ±1	0.39 ±0.01		-12 ±2
	C.EpO3	-22 ±5*	60.5 ±5.1	-16 ±7	0.42 ±0.035		-3 ±8
	CT	-17 ±3	63.1 ±2.7	-12 ±4	0.42 ±0.006		-3 ±2
	CT.EpO3	-37 ±6*	73.6 ±3.4	2 ±5	0.32 ±0.029		-26 ±7*
	CT.O3	-31 ±4*	63.3 ±6.9	-12 ±10	0.39 ±0.034		-11 ±8
	T.EpO3	-22 ±3*	56.9 ±3.8	-21 ±5	0.44 ±0.025		0 ±6

3.3. Ozone uptake and grain yield.

The A and CT treatments contained all three regimes of ozone exposure (Fig. 4). For the A-treatments, the correlation between grain yield and ozone dose was between R^2 0.57 and 0.95, indicating that the increase in ozone dose contributed considerably to the explanation of yield loss. Furthermore, as indicated by the regression equations in Table 3, yield change due to an increased ozone dose was higher in the A-treatments than in the CT-treatments, where the increase in ozone uptake explained only 11 to 55% of the change in yield, due to the detrimental effect of elevated temperature in CT treatments. In both A and CT-treatments, it was the landrace variety's yield change that was least correlated with the increase in ozone dose (R^2 -values of 0.57 (A-settings) and

335 0.11 (CT settings) versus 0.95 and 0.93 in A for KWS Bittern and Lennox resp., and 0.55 and 0.40
 336 in CT). But it may be worth noticing that the Lantvete's $R^2 = 0.11$ in CT-settings covered a yield-
 337 increase in the same order of magnitude as the KWS Bittern's yield loss in those settings. The
 338 anova in Table 4 specifies that in addition to ozone dose ($p= 1.71E-09$), the main effects of variety
 339 and climate as well as interactions between the factors significantly influence grain yield.
 340




341
 342 Fig. 4. Grain yield relating to different amounts of ozone uptake in the two general climate settings;
 343 A (ambient [CO₂] + low temperature) and CT (high [CO₂] + high temperature). The varieties are
 344 presented by color and shape; KWS Bittern (blue squares), Lennox (purple triangles), and Lantvete
 345 (green circles). Solid lines and filled data marks are A treatments and dashed lines with open data
 346 marks are CT treatments. n = 5.
 347

348 Table 2. Regression equations and correlation coefficients of relation between the wheat varieties'
 349 grain yield and plant ozone uptake in ambient temperature and ambient CO₂ treatments (A) and
 350 combined elevated CO₂ and elevated temperature treatments (CT).
 351

	Variety	Regression equation	R ²
A	KWS Bittern	-1.7 + 54.1	0.95
	Lennox	-1.0 + 53.3	0.93
	Lantvete	-1.9 + 43.2	0.57
CT	KWS Bittern	-0.1 + 41.2	0.55
	Lennox	-0.5 + 43.5	0.40
	Lantvete	0.1 + 35.0	0.11

353
354
355
356
357
358
359
360
361

All three varieties had large uptakes of ozone in the T.EpO3 treatment compared to the uptake in the other episodic exposure treatments (not shown). For the episodically exposed treatments with elevated temperature, T.EpO3 and CT.EpO3, the uptake of ozone in the CT.EpO3 treatment appear lower compared to the T.EpO3 for all varieties (Fig. 5). In the colder treatments, the effect of elevated [CO₂] was not as clear and it depended on variety: In the Lennox variety, the ozone uptake was lower in the C.EpO3 treatment than in the A.EpO3, but in the KWS Bittern and Lantvete varieties, the A.EpO3 ozone uptake was the lowest.



362
363
364
365
366
367

Fig. 5. Grain yield of the varieties sorted in treatments. KWS Bittern, Lantvete and Lennox are blue, green and purple, respectively. The plant ozone uptake values of varieties in ozone-exposed treatments are marked with white interruptions on the relevant bar; read at secondary axis. n=5. The marks of no significant difference between the treatments relate to the individual variety (letters = KWS Bittern, numbers = Lantvete, capital letters = Lennox).

Table 3. Anova results for the test of Grain yield as a function of main effects and interactions of Ozone Uptake (OU), in the climate treatments of either ambient [CO₂] + low temperature (A-), or high [CO₂] + high temperature (CT-) and Variety.

Grain Yield	Pr(>F)	
Climate	1.01E-03	**
OU	1.71E-09	***
Variety	8.14E-11	***
Climate : OU	1.06E-07	***
Climate : Variety	2.80E-03	**
OU : Variety	0.915516	
Climate : OU : Variety	0.007374	**

3.4. Plant performance

3.4.1. Photosynthesis versus intercellular carbon dioxide.

The amount of photosynthesis taking place at different availabilities of carbon dioxide is illustrated in Fig. 6 for each variety. If the photosynthetic apparatus is well-functioning, it responds to an increase in CO₂ with an increase in photosynthesis. In the figures, the CO₂-component of the treatments divides the possible increase in photosynthesis depending on treatment. As expected, the CO₂-enrichments resulted in measurements of high rates of photosynthesis, but also measurements of low rates of photosynthesis at high levels of carbon dioxide. The CO₂-concentration in the plant cells of the non-CO₂-enriched treatments was less varied in distribution, and photosynthesis never peaked as high as in the C-treatments but were more often in the lower range (more PS<10 than in C-treatments).

The T.EpO3 Lennox, CT.O3 Lantvete, and A.O3 Lantvete, all had negative relations between intercellular carbon dioxide and photosynthesis (Fig. 7). The majority of the other treatments*varieties have regression slopes with values of 0.05 or less, only six have higher values, among these are all A.EpO3-varieties, modern varieties of A, and Lantvete from the CT-treatment. Table 5 shows the variance analysis for the photosynthesis' dependence on ozone, treatment, intercellular carbon, and variety.

390 Table 4. Anova result of the test for Rate of photosynthesis as a function of main effects of variety
 391 and main effects and interactions of intercellular carbon, ozone (on/off), (high/low) temperature
 392 setting.

Photosynthesis ~ Variety + Ci * ozone * Temp	Pr(>F)	
Variety	2.46E-07	***
Ci	< 2.2e-16	***
Ozone	1.37E-02	*
Temp	9.63E-07	***
Ci : Ozone	0.001023	**
Ci : Temp	6.11E-06	***
Ozone : Temp	1.94E-13	***
Ci : Ozone : Temp	2.24E-03	**

393


394

395 3.4.2. 1000-grain weight, water consumption, protein content, and gluten index.

396 The protein mass was, on average across the treatments, 5.8 ± 0.1 , 5.8 ± 0.2 and 5.2 ± 0.1 g for KWS
 397 Bittern, Lennox and Lantvete, respectively. There was statistically significant interaction between
 398 treatments and varieties (Table 6), although the data for individual treatments are not shown here.

399 Gluten index was on average across the treatments 28.7 ± 1.9 , 29.4 ± 2.5 and 30.8 ± 0.8 (with SD) for
 400 KWS Bittern, Lennox and Lantvete, respectively. There was significant interaction between
 401 treatment and variety. The individual treatment*varieties are illustrated with grain yield in Fig. 8.

402 The 1000-grain weight was also significantly influenced by interactions between treatment and
 403 variety. The total water consumption reflected the different water use efficiencies between the
 404 landrace and the modern varieties, with the modern varieties consuming less water than the landrace
 405 (Fig. 8).



406

407 Fig. 6. Obtained net photosynthesis/ intercellular CO₂-concentration over time. The varieties are
 408 referred to by the data mark shapes; squares, circles, triangles are KWS Bittern, Lantvete, and


409 Lennox, respectively. The colors refer to Fig. 1 of the treatments: lightest blue (A), medium blue
 410 (A.EpO3), dark blue (A.O3,) orange (T.EpO3), yellow with black outline (CT), tan (CT.EpO3),
 411 dark tan (CT.O3), and green (C.EpO3). Above the horizontal axis is marked the duration of ozone
 412 exposure in the colors of the treatments (dashed lines represent carbon enrichment). See Fig. 1 for
 413 further explanation of treatment abbreviations.
 414

415 Table 6. Anova results of test of protein content (g pot⁻¹), gluten index (ICC 155), 1000-grain
 416 weight (g pot⁻¹) and vegetative dry matter (g pot⁻¹) as functions of main effects and interactions of
 417 treatment and variety.


	Protein content	Gluten index	1000-grain weight	Vegetative DM
Treatment	< 2.2E-16 ***	< 2.2E-16 ***	< 2.2E-16 ***	9.50E-12 ***
Variety	0.2898	< 2.2E-16 ***	< 2.2E-16 ***	< 2.2E-16 ***
Treatment : Variety	7.59E-12 ***	< 2.2E-16 ***	0.000188 ***	0.000244 ***

418

419




422 Fig. 7. In descending order, the regression slope values (circles) from Fig. 6 Rate of photosynthesis
423 versus intercellular CO₂-concentration. Correlation coefficient, R² (dashes).



425 Fig. 8. The three diagrams show the grain yield of the treatments presented in descending order but
426 clustered in variety (please note that the treatments do not show in the same order for all varieties).
427 Alongside the yield is the 1000-grain weight, the gluten index and the total water consumption. The
428 data marks are as follow: x (1000 grain weights), \diamond (gluten index) and o (total water consumption).
429 See Fig. 1 for explanation of treatment abbreviations.
430
431

432 3.4.3. The plant development and leaf discoloration.

433 The wheat plant varieties developed at a similar pace with differences induced by treatments (Fig.
434 9). The majority of the differences could be attributed to differences in temperature; however,
435 periods of 5-7 days of delay of development from CO₂-enrichment and ozone exposure could be
436 detected at different times during growth, mostly in the lower temperature treatments.
437 Similar influence from temperature first and ozone second can be seen in the development of leaf
438 discoloration as in Fig. 10 showing also *some* inter-varietal differences. In the Lantvete variety the
439 different treatments induce the largest differences and in the KWS Bittern the smallest.



440

441 Fig. 9. The plant development after sowing, scored as Zadoks' growth stages 2-3 times a week,
 442 exemplified by the Lantvete variety. The colors refer to Fig. 1 of the treatments: lightest blue (A),
 443 medium blue (A.EpO3), dark blue (A.O3,) orange (T.EpO3), yellow (CT), tan (CT.EpO3), dark tan
 444 (CT.O3), and green (C.EpO3). Carbon-enriched treatments (C-) are shown as dashed lines. Above
 445 the horizontal axis is marked the duration of ozone exposure in the colors of the treatments (dashed
 446 lines represent carbon enrichment). Further explanation of treatment abbreviations in Fig. 1.




Fig. 10 Discoloration of second leaf at four dates in the three varieties in the eight treatments. The discoloration is shown as part of total leaf area being green. Square, circle and triangle data marks are KWS Bittern, Lantvete and Lennox respectively. The discoloration is assessed while all treatments were subjected to ozone exposure following the initiation of ozone exposure for the episodic ozone treatments. Further explanation of treatment abbreviations in Fig. 1.

4. Discussion


455 The discussion includes the performance of facilities and settings to realize the chosen climate
456 treatments, after which the discussion considers the results relating to the hypotheses.

457 4.1. Realized climates

458 Apart from the level of CO₂ in treatments with settings of ambient CO₂, the logging of variables by
459 the RERAF facilities show that the experimental values in general were in accordance with the set
460 points of variables (see 3.1). The elevated temperature treatment could possibly have been so high
461 as to induce a moderate chronic heat stress to the wheat plants (Sandhu et al., 2018), contributing to
462 an explanation of the dominating impact of temperature relative to the ozone stress.

463 An unplanned increase in CO₂ level in a facility as RERAF, which cannot remove excess CO₂, may
464 arise from several sources. The use of a growth medium with a high content of organic matter may
465 result in microbial decomposition causing a measurable impact on CO₂-concentration as in
466 Ingvordsen et al. (2018). An increase at nighttime when no photosynthesis occur is similar to
467 natural ecosystems where increases in [CO₂] are seen at low wind speeds (Mikkelsen et al., 2008),
468 and also suggested by Ingvordsen et al. (2018) as a possible explanation for their high average
469 [CO₂]. In the present case, however, the set point was exceeded by approximately 150 ppm, approx
470 100 ppm higher than Ingvordsen et al. (2018) reported. One key difference to the Ingvordsen et al.
471 experiment, was that we did manual watering to obtain knowledge on the water consumption
472 throughout the plants' growth which increased human presence in the chambers. Human activities
473 from measuring and maintaining the plants in the chambers resulted in peaks of [CO₂] of 6-900 ppm
474 depending on the activity, and consequently the impact of human respiration on the CO₂-
475 concentration in the chambers were not adequately accounted for in the experiment set-up. The
476 increase in average chamber CO₂ compared to atmospheric CO₂-concentration minimize the
477 detectable effects of the CO₂-enriched treatments as the difference in plant functioning will be more
478 influenced by the first increase of [CO₂] compared to the influence arising from the increase

479 assigned the CO₂-enrichment-treatments. For future trials of a similar kind, exchanging the chamber
 480 air at a higher rate than 4 m³ h⁻¹ could be explored for better neutralizing such peak CO₂-
 481 enrichments. An extract of measurements of carbon dioxide concentration from the chambers of
 482 treatment A is shown in Fig. 11, where the nighttime respiration accumulation as well as the
 483 attribution of human activity to chamber [CO₂] can be seen.



484
 485 Fig. 11. CO₂ concentration of treatment A, CO₂-setting 400 ppm. Highlighted two days of no
 486 human interference and one with. Arrows 'a' indicate the decrease in daytime when photosynthesis
 487 is activated, 'b' indicate nighttime increase due to plant and soil respiration, and 'c' denote incidents
 488 of human presence in the chamber for a shorter or longer period for watering or measuring reasons.
 489

490
 491 When growing agricultural crops in pots, there are many conditions that are different from field
 492 conditions. In this study, there were fewer plants m⁻² than would be normal in the field, and the
 493 boundary conditions of the potted plants would be far from those of field-grown plants. Albeit
 494 grown in sphagnum in pots and while inside climate chambers, the plants in this study grew to
 495 maturity and produced yields in a range from 5-12 ton ha⁻¹ (see 3. Results), which is better than, but
 496 including, the Danish spring wheat yield average from 2017 of 5.2 ton ha⁻¹ (Lundø, 2017). A higher
 497 yield in climate chambers than in fields is not rare, as many biotic and abiotic yield-reducing factors
 498 are eliminated in climate chambers (Poorter et al., 2016). In the ambient treatments in this study,

499 ozone as a yield-reducing factor is eliminated due to the molecules' reactivity with surfaces through
500 the ventilation system. In the ozone treatments high ozone concentrations were measured in the
501 chambers, but due to low wind speed in the chambers ($<0.6 \text{ m s}^{-1}$) the boundary layer resistance can
502 be assumed to have been considerable and that the plants therefore were less exposed to toxic levels
503 of ozone than what the concentration itself provided. The light supply can be claimed to be
504 insufficient compared to what can be measured in the field, however as it was constant we believe
505 the drawbacks from a suboptimal lighting are evened by the consistency in the supply.

506

507 4.2. Older versus modern wheat varieties.

508 The modern varieties, KWS Bittern and Lennox, which we used in the experiment, were bred to
509 yield well and are used in Denmark and Southern France respectively, and the harvest index (Table
510 2) reveals that their higher yields are based on less biomass than the landrace variety's. Breeding of
511 varieties for modern production take place under current ozone conditions, and as so, modern
512 varieties are supposedly naturally adapted to the current increases in ozone concentration, but the
513 modern wheat varieties tend to be more sensitive to ozone (Biswas et al., 2008; Pleijel et al., 2006).
514 Studies examining the effects of filtered air as well as unfiltered air on the growth and yields of
515 wheat show significant yield loss at current concentrations of ozone, indicating that breeding is not
516 successful in obtaining full ozone resistance (Pleijel et al., 2018, 2006). On the other hand, the
517 landrace variety in this study yielded as poorly when temporarily exposed to ozone as when
518 constantly exposed, where the yield of the modern variety were not significantly different from no
519 ozone to temporary exposure. This indicates perhaps that the modern varieties are resistant to some
520 degree of ozone at the lower temperature settings compared to the resistance of the landrace.
521 The landrace variety used in this study has no commercial use at present, which points to its lack of
522 ability to compete on yields or sales. The Lantvete yield is on the other hand only significantly

523 lower than the modern varieties in two treatments. Its larger biomass may be responsible for its
524 resilience towards the warmer CO₂-enriched treatments (Lopes et al., 2015). The changes in harvest
525 indexes between the treatments reflect that the biomass allocation is a possible actor in the way the
526 varieties deal with the changes in growth related climatic factors (Table 2).

527 As Lennox is a common variety grown in the south of France, it could be expected to resist warmer
528 temperatures (in CT and T.EpO3 treatments in particular) better than KWS Bittern and Lantvete.
529 This seems to be the case: the yield loss in CT treatment is approx. 17% for Lennox, while it's 28%
530 and 39% for KWS Bittern and Lantvete. That is the only non-significant loss ($p = 0.19$) from the A
531 treatment to a warm treatment within the same variety, it is also significantly higher than the yield
532 of CT Lantvete (Fig. 2).

533 The yield losses in T.EpO3 treatment compared to the A.EpO3 are approx. 10% for Lennox and
534 approx. 26% for KWS Bittern, while Lantvete gains 20% from A.EpO3 to T.EpO3. The differences
535 in loss as a temperature response align with the geographic purpose of the two modern varieties,
536 where Lennox is used in the south of Europe and KWS Bitten in more northern regions. The
537 Lantvete gain is probably more related to the severe grain yield loss in A.EpO3 than to a proper
538 acclimatization to elevated temperatures (table 2).

539

540 4.3. Episodic versus full time ozone exposure and effects on wheat yield

541 It was expected that the plants experiencing ozone as a longer-stretched episode at a critical timing
542 lose as much in yields as plants that have adjusted to ozone from growing in a continuously
543 medium-high ozone environment. The Lantvete variety had similar losses in the colder treatment,
544 but the two modern varieties lost more in the A.O3 than in the A.EpO3 treatment (Fig. 4). It could
545 be assumed that intraspecific variations in the ozone acclimation of growth physiological processes
546 exist among wheat cultivars. Such results were shown on thermal acclimation of photosynthesis

among alfalfa cultivars (Zaka et al., 2016). However, it could also refer to the contradictory conditions between ambient air ozone concentration and chamber background concentration of ozone. Treatment A had in fact conditions resembling filtered air treatments, resulting in A reflecting the natural selection for ozone adaptation in the modern varieties' yield. This aligns with literature on ozone diminishing agricultural yields at present conditions (Pleijel et al., 2018). In the warmer treatment, the correlation between the dose of phytotoxic ozone and grain yield was not as strong as in the cold treatments (Table 3). In these wheat varieties, an adaptation to ozone to reduce the impact of continuous ozone to the level of episodic ozone exposure was not found. Linear correlations between grain yield and plant ozone uptake (reported as POD_6) were also reported by Harmens et al. (2018) and Grünhage et al (2012). The weaker correlations between grain yield and ozone uptake in the warmer CO_2 -enriched treatments indicate that the climate components influence the yields more than the ozone dose.

559

4.4. Multiple factors and additional ozone exposure

Increases in $[CO_2]$ have been expected to induce a fertilizing effect on plant growth and the effect of temperature follows a bell curve indicating that it may be too low as well as too high for optimum plant functioning (Norby and Luo, 2004). The best yields in this study were attained in the ambient $[CO_2]$ / low temperature treatment (Table 2), indicating that the additional parameters, increase in $[CO_2]$ and temperature, alone or combined all skewed the plant growth out of its optimum. However, the less than additive positive effect of the combination of changes in climate factors have been documented previously (Frenck et al., 2013; Ingvordsen et al., 2015; Shaw et al., 2002). Broberg et al. (2015) observe in their review that both negative and positive effects are to be expected from ozone exposure.

570 The yields of CT and CT-ozone-treatments were not significantly different. If ozone uptake was the
571 main reason of yield reduction, increased ozone exposure should diminish the yields accordingly.
572 However, the uptake of ozone cannot alone explain the yield decrease of the CT-treatments, and the
573 climate treatment must be contributing.

574 The ozone taken up influenced the varieties' yields differently, for example, very different ozone
575 amounts were taken up in in T.EpO3, but with no comparative yield decline, stressing the
576 interactions of climate and variety on the grain yield (Fig. 5, Table 4). For the episodically exposed
577 treatments with elevated temperature, T.EpO3 and CT.EpO3, a possible effect of the stomatal
578 regulative effect of elevated [CO₂] is visible qua the lower uptake of ozone in the CT.EpO3
579 treatment compared to the T.EpO3.

580

581 Although not significant, we observe that the distribution of Lantvete yields suggests a plasticity of
582 the grain yield from A.EpO3 to CT.EpO3, indicating a trend of not losing additional yields with the
583 future climate scenario. The modern varieties lose yield from A.EpO3 to CT.EpO3; Lennox
584 significantly, KWS Bittern not significantly.

585 From Fig. 4, the impact of ozone is presumably greater in colder temperature treatments, and the
586 addition of ozone to a treatment with elevated [CO₂] and elevated temperature does not significantly
587 increase injury to yields already affected by the higher temperature. However, Tai and Martin
588 (2017) suggest an adaptation of crops to regional ozone and water constraints to induce a stronger
589 ozone tolerance. Elevated [CO₂] may also influence the water relation. In this study, it was decided
590 to keep the plants well-watered. However, had the plants lacked water at times, a better stoma
591 regulation might have reduced the ozone uptake.

592

593 4.5. Growth parameters and responses during growth

594 The photosynthesis vs. intercellular CO₂ depicted in Fig. 6 a, b, c, show that the photosynthetic
595 apparatus of the plants worked as expected on altered input in all varieties and in most treatments.
596 From the sorting of regression slopes in Fig. 7, it seems that the photosynthetic apparatus was
597 affected by the amendment of ozone, and especially full-time ozone treatments resulted in negative
598 regression slopes. This is in accordance with the general perception that ozone has a direct negative
599 influence on photosynthesis (Mikkelsen, 1995; Mikkelsen and Ro-Poulsen, 2013).

600 Protein production could be depending on soil nutrient availability rather than treatment in this case
601 (Thomsen et al., 2008). The obtained amount of protein mass did not vary as much as grain yields,
602 but the baking quality in form of gluten index indicates that the quality and quantity was not tightly
603 linked in this case. Rather, the lower yields represented a better baking quality.

604 The higher total water consumption (Fig. 8c) of the landrace variety is in accordance with the
605 higher biomass of the Lantvete (Table 2) and is a trait of a less efficient variety.

606 As expected, the phenological development of the plants in the treatments (Fig. 9) revealed a
607 stronger dependence on temperature than on any other treatment parameter. However, there was a
608 5-degree difference between colder and warmer treatments, and the strong reflection of temperature
609 influence could be caused by this biologically large leap. The impact of [CO₂]-enrichment could be
610 detected on close observation of the development data (not shown here) and this impact may also
611 relate to the absolute difference between the treatment concentrations.

612 Due to the nature of the sampling of only one leaf in a treatment at a time the results of the
613 development of discoloration of leaves are indicative rather than significant, however, as seen in
614 Fig. 10, the varieties respond differently to the treatments, with the Lantvete variety showing more
615 variation than the modern varieties, and the modern expressing some difference in their response to
616 temperature.

617 5. Conclusion

618 The various combinations of two levels of temperature, two levels of [CO₂], and three ozone
619 exposure regimes affected both growth and yield in spring wheat, while interacting with the studied
620 spring wheat varieties. On average, across all treatments, the landrace variety Lantvete had
621 significantly lower yields than the modern varieties Lennox and KWS Bittern. However, the yields
622 were significantly lower within only two treatments; in the treatment with ambient levels of
623 temperature and [CO₂] and episodically exposure to ozone (A.EpO3), and in the treatment with an
624 elevated level of temperature and [CO₂] (CT). Within the other six treatments, there was no
625 significant difference between the yields of the three tested varieties.

626 The effect on yield with episodic ozone exposure relative to treatment A depended on variety. Yield
627 losses relative to treatment A were significant in all CT.EpO3 and T.EpO3, in A.EpO3 of KWS
628 Bittern and Lantvete, and in C.EpO3 of Lennox. Yield losses due to chronic ozone exposure were
629 significant in all varieties. Increased [CO₂] and higher temperature without ozone exposure induced
630 significant loss of yield in Lantvete and KWS Bittern. Full-time exposure to ozone reduced yields
631 more than ozone exposure during limited time. The combination of climate factors and ozone has
632 shown that the effect of changes in climate factors on wheat yield influence the effect of ozone, and
633 that the influence of temperature can be detrimental regardless of ozone exposure. While an
634 increase in the ozone dose induced yield loss in the low temperature and ambient [CO₂] treatment,
635 the effect in the warm treatments was less clear and yields were lower even at low ozone doses. The
636 different reactions of the varieties to higher ozone doses in the warmer CO₂-enriched treatments
637 compared to the reactions in the treatments with lower temperature and lower [CO₂], lead to the
638 assumption that responses to the combined effects of climate factor changes (temperature, [CO₂],
639 ozone) are due to either more or less additive intraspecific variations and adaptations or one of the
640 factors (e.g. elevated temperature) having an overriding influence on the impact of other factors.

641 Declaration of interest

642 The authors declare that they have no conflict of interest.

643

644 Acknowledgement

645 This research is part of the Joint Programming Initiative on Agriculture, Food Security and Climate

646 Change (FACCE-JPI) and funded by the FACCE-ERA-NET+ project: Climate-CAFÉ.

647 Saaten-Union, Danish AGRO and Nordic Genebank kindly provided the spring wheat seeds for the

648 study.

649

650

651 REFERENCES

- 652 AbdElgawad, H., Farfan-Vignolo, E.R., Vos, D. De, Asard, H., 2015. Elevated CO₂ mitigates
653 drought and temperature-induced oxidative stress differently in grasses and legumes. *Plant Sci.*
654 231, 1–10. <https://doi.org/10.1016/j.plantsci.2014.11.001>
- 655 Ainsworth, E.A., Yendrek, C.R., Sitch, S., Collins, W.J., Emberson, L.D., 2012. The effects of
656 tropospheric ozone on net primary productivity and implications for climate change. *Annu.*
657 *Rev. Plant Biol.* 63. <https://doi.org/10.1146/annurev-arplant-042110-103829>
- 658 Albert, K.R., Mikkelsen, T.N., Michelsen, A., Ro-Poulsen, H., van der Linden, L., 2011. Interactive
659 effects of drought, elevated CO₂ and warming on photosynthetic capacity and photosystem
660 performance in temperate heath plants. *J. Plant Physiol.* 168, 1550–1561.
661 <https://doi.org/10.1016/j.jplph.2011.02.011>
- 662 Biswas, D.K., Xu, H., Li, Y.G., Sun, J.Z., Wang, X.Z., Han, X.G., Jiang, G.M., 2008. Genotypic
663 differences in leaf biochemical, physiological and growth responses to ozone in 20 winter
664 wheat cultivars released over the past 60 years. *Glob. Chang. Biol.* 14, 46–59.
665 <https://doi.org/10.1111/j.1365-2486.2007.01477.x>
- 666 Booker, F., Muntifering, R., Mcgrath, M., Burkey, K., Decoteau, D., Fiscus, E., Manning, W.,
667 Krupa, S., Chappelka, A., Grantz, D., 2009. The ozone component of global change: Potential
668 effects on agricultural and horticultural plant yield, product quality and interactions with
669 invasive species. *J. Integr. Plant Biol.* 51, 337–351. [https://doi.org/10.1111/j.1744-](https://doi.org/10.1111/j.1744-7909.2008.00805.x)
670 [7909.2008.00805.x](https://doi.org/10.1111/j.1744-7909.2008.00805.x)
- 671 Brisson, N., Gate, P., Gouache, D., Charmet, G., Oury, F.X., Huard, F., 2010. Why are wheat yields
672 stagnating in Europe? A comprehensive data analysis for France. *F. Crop. Res.* 119, 201–212.
673 <https://doi.org/10.1016/j.fcr.2010.07.012>
- 674 Broberg, M.C., Feng, Z., Xin, Y., Pleijel, H., 2015. Ozone effects on wheat grain quality - A

summary. Environ. Pollut. 197, 203–213. <https://doi.org/10.1016/j.envpol.2014.12.009>

Calatayud, V., García-Breijo, F.J., Lia Cervero, J., Reig-Armiñ Ana, J., Sanz, M.J., 2011. Physiological, anatomical and biomass partitioning responses to ozone in the Mediterranean endemic plant *Lamottea diana*. Ecotoxicol. Environ. Saf. 74, 1131–1138. <https://doi.org/10.1016/j.ecoenv.2011.02.023>

Clausen, S.K., Frenck, G., Linden, L.G., Mikkelsen, T.N., Lunde, C., Jørgensen, R.B., 2011. Effects of single and multifactor treatments with elevated temperature, CO₂ and ozone on oilseed rape and barley. J. Agron. Crop Sci. 197, 442–453. <https://doi.org/10.1111/j.1439-037X.2011.00478.x>

Cure, J.D., Acock, B., 1986. Crop responses to carbon dioxide doubling: a literature survey. Agric. For. Meteorol. 38, 127–145. [https://doi.org/10.1016/0168-1923\(86\)90054-7](https://doi.org/10.1016/0168-1923(86)90054-7)

Derwent, R.G., Manning, A.J., Simmonds, P.G., Spain, T.G., O'Doherty, S., 2018. Long-term trends in ozone in baseline and European regionally-polluted air at Mace Head, Ireland over a 30-year period. Atmos. Environ. 179, 279–287. <https://doi.org/10.1016/j.atmosenv.2018.02.024>

Dieleman, W.I.J., Vicca, S., Dijkstra, F. a., Hagedorn, F., Hovenden, M.J., Larsen, K.S., Morgan, J. a., Volder, A., Beier, C., Dukes, J.S., King, J., Leuzinger, S., Linder, S., Luo, Y., Oren, R., De Angelis, P., Tingey, D., Hoosbeek, M.R., Janssens, I. a., 2012. Simple additive effects are rare: A quantitative review of plant biomass and soil process responses to combined manipulations of CO₂ and temperature. Glob. Chang. Biol. 18, 2681–2693. <https://doi.org/10.1111/j.1365-2486.2012.02745.x>

Dolferus, R., 2014. To grow or not to grow: A stressful decision for plants. Plant Sci. 229, 247–261. <https://doi.org/10.1016/j.plantsci.2014.10.002>

Finlayson-Pitts, B.J., Pitts, J.N., 1993. Atmospheric chemistry of tropospheric ozone formation:

699 Scientific and regulatory implications. *Air Waste* 43, 1091–1100.

700 Frenck, G., van der Linden, L., Mikkelsen, T.N., Brix, H., Jørgensen, R.B., 2013. Response to
701 multi-generational selection under elevated [CO₂] in two temperature regimes suggests
702 enhanced carbon assimilation and increased reproductive output in *Brassica napus* L. *Ecol.*
703 *Evol.* 3, 1163–1172. <https://doi.org/10.1002/ece3.523>

704 Frenck, G., van der Linden, L., Mikkelsen, T.N., Brix, H., Jørgensen, R.B., 2011. Increased [CO₂]
705 does not compensate for negative effects on yield caused by higher temperature and [O₃] in
706 *Brassica napus* L. *Eur. J. Agron.* 35, 127–134. <https://doi.org/10.1016/j.eja.2011.05.004>

707 Fuhrer, J., Booker, F., 2003. Ecological issues related to ozone: Agricultural issues. *Environ. Int.*
708 29, 141–154. [https://doi.org/10.1016/S0160-4120\(02\)00157-5](https://doi.org/10.1016/S0160-4120(02)00157-5)

709 Gibson, L.R., Paulsen, G.M., 1999. Yield Components of Wheat Grown under High Temperature
710 Stress during Reproductive Growth. *Crop Sci.* 39, 1841.
711 <https://doi.org/10.2135/cropsci1999.3961841x>

712 Grünhage, L., Pleijel, H., Mills, G., Bender, J., Danielsson, H., Lehmann, Y., Castell, J.F.,
713 Bethenod, O., 2012. Updated stomatal flux and flux-effect models for wheat for quantifying
714 effects of ozone on grain yield, grain mass and protein yield. *Environ. Pollut.* 165, 147–157.
715 <https://doi.org/10.1016/j.envpol.2012.02.026>

716 Harmens, H., Hayes, F., Mills, G., Sharps, K., Osborne, S., Pleijel, H., 2018. Wheat yield responses
717 to stomatal uptake of ozone: Peak vs rising background ozone conditions. *Atmos. Environ.*
718 173, 1–5. <https://doi.org/10.1016/j.atmosenv.2017.10.059>

719 ICP Vegetation, 2017. III. Mapping Critical Levels for Vegetation.

720 Ingvordsen, C.H., Backes, G., Lyngkjær, M.F., Peltonen-Sainio, P., Jensen, J.D., Jalli, M., Jahoor,
721 A., Rasmussen, M., Mikkelsen, T.N., Jørgensen, R.B., Stockmarr, A., Jørgensen, R.B., 2015.
722 Significant decrease in yield under future climate conditions: Stability and production of 138

723 spring barley accessions. *Eur. J. Agron.* 63, 105–113. <https://doi.org/10.1016/j.eja.2014.12.003>

724 Ingvordsen, C.H., Lyngkjær, M.F., Peltonen-Sainio, P., Mikkelsen, T.N., Stockmarr, A., Jørgensen,
725 R.B., 2018. How a 10-day heatwave impacts barley grain yield when superimposed onto future
726 levels of temperature and CO₂ as single and combined factors. *Agric. Ecosyst. Environ.* 259,
727 45–52. [https://doi.org/https://doi.org/10.1016/j.agee.2018.01.025](https://doi.org/10.1016/j.agee.2018.01.025)

728 IPCC, 2014. *Climate Change 2014: Synthesis Report, Contribution of working groups I, II and III*
729 *to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Geneva,
730 Switzerland. <https://doi.org/10.1017/CBO9781107415324>

731 Kleanthous, S., Vrekoussis, M., Mihalopoulos, N., Kalabokas, P., Lelieveld, J., 2014. On the
732 temporal and spatial variation of ozone in Cyprus. *Sci. Total Environ.* 476–477, 677–687.
733 <https://doi.org/10.1016/j.scitotenv.2013.12.101>

734 Lamarque, J.-F., Bond, T.C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C.,
735 Mieville, A., Owen, B., Schultz, M.G., Shindell, D., Smith, S.J., Stehfest, E., Van Aardenne,
736 J., Cooper, O.R., Kainuma, M., Mahowald, N., McConnell, J.R., Naik, V., Riahi, K., Van
737 Vuuren, D.P., 2010. Historical (1850–2000) gridded anthropogenic and biomass burning
738 emissions of reactive gases and aerosols: methodology and application. *Atmos. Chem. Phys*
739 10, 7017–7039. <https://doi.org/10.5194/acp-10-7017-2010>

740 Langley, J.A., Hungate, B.A., 2014. Plant community feedbacks and long-term ecosystem
741 responses to multi-factored global change. *AoB Plants* 6, 1–12.
742 <https://doi.org/10.1093/aobpla/plu035>

743 Larsen, K.S., Andresen, L.C., Beier, C., Jonasson, S., Albert, K.R., Ambus, P., Arndal, M.F.,
744 Carter, M.S., Christensen, S., Holmstrup, M., Ibrom, A., Kongstad, J., Van Der Linden, L.,
745 Maraldo, K., Michelsen, A., Mikkelsen, T.N., Pilegaard, K., Priemé, A., Ro-Poulsen, H.,
746 Schmidt, I.K., Selsted, M.B., Stevnbak, K., 2011. Reduced N cycling in response to elevated

747 CO₂, warming, and drought in a Danish heathland: Synthesizing results of the CLIMAITE
748 project after two years of treatments. *Glob. Chang. Biol.* 17, 1884–1899.
749 <https://doi.org/10.1111/j.1365-2486.2010.02351.x>

750 LICOR, 2004. Using the LI-6400 Portable Photosynthesis - OPEN Software version 5.3, 2005th ed,
751 Components. LI-COR Biosciences, Lincoln, Nebraska.

752 Lopes, M.S., El-Basyoni, I., Baenziger, P.S., Singh, S., Royo, C., Ozbek, K., Aktas, H., Ozer, E.,
753 Ozdemir, F., Manickavelu, A., Ban, T., Vikram, P., 2015. Exploiting genetic diversity from
754 landraces in wheat breeding for adaptation to climate change. *J. Exp. Bot.* 66, 3477–3486.
755 <https://doi.org/10.1093/jxb/erv122>

756 Lundø, M., 2017. Statistical report: Harvest of bulk grain, rapeseed and grain legumes 2017 (In
757 Danish).

758 Mikkelsen, T.N., 1995. Physiological responses of *Fagus sylvatica* L. exposed to low levels of
759 ozone in open-top chambers. *Trees* 9, 355–361. <https://doi.org/10.1007/BF00202500>

760 Mikkelsen, T.N., Beier, C., Jonasson, S., Holmstrup, M., Schmidt, I.K., Ambus, P., Pilegaard, K.,
761 Michelsen, A., Albert, K., Andresen, L.C., Arndal, M.F., Bruun, N., Christensen, S., Danbæk,
762 S., Gundersen, P., Jørgensen, P., Linden, L.G., Kongstad, J., Maraldo, K., Priemé, A., Riis-
763 Nielsen, T., Ro-Poulsen, H., Stevnbak, K., Selsted, M.B., Sørensen, P., Larsen, K.S., Carter,
764 M.S., Ibrom, A., Martinussen, T., Miglietta, F., Sverdrup, H., 2008. Experimental design of
765 multifactor climate change experiments with elevated CO₂, warming and drought: The
766 CLIMAITE project. *Funct. Ecol.* 22, 185–195. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2435.2007.01362.x)
767 [2435.2007.01362.x](https://doi.org/10.1111/j.1365-2435.2007.01362.x)

768 Mikkelsen, T.N., Ro-Poulsen, H., 2013. Exposure increases of Norway spruce to current ozone year
769 the to sensitivity of needles photoinhibition. *New Phytol.* 128, 153–163.

770 Mills, G., Sharps, K., Simpson, D., Pleijel, H., Frei, M., Burkey, K., Emberson, L., Uddling, J.,

771 Broberg, M., Feng, Z., Kobayashi, K., Agrawal, M., 2018. Closing the global ozone yield gap:
 772 Quantification and cobenefits for multistress tolerance. *Glob. Chang. Biol.* 24, 4869–4893.
 773 <https://doi.org/10.1111/gcb.14381>

774 Monks, P.S., Archibald, A.T., Colette, A., Cooper, O., Coyle, M., Derwent, R., Fowler, D., Granier,
 775 C., 2015. Tropospheric ozone and its precursors from the urban to the global scale from air
 776 quality to short-lived climate forcer. *Atmos. Chem. Phys* 15, 8889–8973.
 777 <https://doi.org/10.5194/acp-15-8889-2015>

778 Munir, S., Chen, H., Ropkins, K., 2013. Quantifying temporal trends in ground level ozone
 779 concentration in the UK. *Sci. Total Environ.* 458–460, 217–227.
 780 <https://doi.org/10.1016/j.scitotenv.2013.04.045>

781 Namazkar, S., Stockmarr, A., Frenck, G., Egsgaard, H., Terkelsen, T., Mikkelsen, T., Ingvordsen,
 782 C.H., Jørgensen, R.B., 2016. Concurrent elevation of CO₂, O₃ and temperature severely
 783 affects oil quality and quantity in rapeseed. *J. Exp. Bot.* 67, 4117–4125.
 784 <https://doi.org/10.1093/jxb/erw180>

785 Norby, R.J., Luo, Y., 2004. Evaluating ecosystem responses to rising atmospheric CO₂ and global
 786 warming in a multi-factor world. *New Phytol.* 162, 281–293. [https://doi.org/10.1111/j.1469-](https://doi.org/10.1111/j.1469-8137.2004.01047.x)
 787 [8137.2004.01047.x](https://doi.org/10.1111/j.1469-8137.2004.01047.x)

788 Peltonen-Sainio, P., Muurinen, S., Rajala, A., Jauhiainen, L., 2008. Variation in harvest index of
 789 modern spring barley, oat and wheat cultivars adapted to northern growing conditions. *J.*
 790 *Agric. Sci.* 146, 35–47. <https://doi.org/10.1017/S0021859607007368>

791 Pleijel, H., Broberg, M.C., Uddling, J., Mills, G., 2018. Current surface ozone concentrations
 792 significantly decrease wheat growth, yield and quality. *Sci. Total Environ.* 613, 687–692.
 793 <https://doi.org/10.1016/j.scitotenv.2017.09.111>

794 Pleijel, H., Eriksen, A.B., Danielsson, H., Bondesson, N., Selldén, G., 2006. Differential ozone

795 sensitivity in an old and a modern Swedish wheat cultivar - Grain yield and quality, leaf
 796 chlorophyll and stomatal conductance. *Environ. Exp. Bot.* 56, 63–71.
 797 <https://doi.org/10.1016/j.envexpbot.2005.01.004>

798 Poorter, H., Fiorani, F., Pieruschka, R., Wojciechowski, T., van der Putten, W.H., Kleyer, M.,
 799 Schurr, U., Postma, J., 2016. Pampered inside, pestered outside? Differences and similarities
 800 between plants growing in controlled conditions and in the field. *New Phytol.* 212, 838–855.
 801 <https://doi.org/10.1111/nph.14243>

802 Sandhu, S.S., Singh, J., Kaur, P., Gill, K.K., 2018. Heat Stress in Field Crops: Impact and
 803 Management Approaches, in: *Advances in Crop Environment Interaction*. Springer Singapore,
 804 Singapore, pp. 181–204. https://doi.org/10.1007/978-981-13-1861-0_7

805 Schneider, C.A., Rasband, W.S., Eliceiri, K.W., 2012. NIH Image to ImageJ: 25 years of image
 806 analysis.

807 Shaw, M.R., Zavaleta, E.S., Chiariello, N.R., 2002. Grassland Responses to Global Environmental
 808 Changes Suppressed by Elevated CO₂. *Science* (80-.). 298, 1987–1990.

809 Tai, A.P.K.K., Martin, M.V., 2017. Impacts of ozone air pollution and temperature extremes on
 810 crop yields: Spatial variability, adaptation and implications for future food security. *Atmos.*
 811 *Environ.* 169. <https://doi.org/10.1016/j.atmosenv.2017.09.002>

812 Thomsen, I.K., Pedersen, L., Jørgensen, J.R., 2008. Yield and flour quality of spring wheat as
 813 affected by soil tillage and animal manure. *J. Sci. Food Agric.* 88, 2117–2124.
 814 <https://doi.org/10.1002/jsfa.3322>

815 Vázquez, D.P., Gianoli, E., Morris, W.F., Bozinovic, F., 2017. Ecological and evolutionary impacts
 816 of changing climatic variability. *Biol. Rev.* 92, 22–42. <https://doi.org/10.1111/brv.12216>

817 Vingarzan, R., 2004. A review of surface ozone background levels and trends. *Atmos. Environ.* 38,
 818 3431–3442. <https://doi.org/10.1016/j.atmosenv.2004.03.030>

819 Wild, O., Fiore, A.M., Shindell, D.T., Doherty, R.M., Collins, W.J., Dentener, F.J., Schultz, M.G.,
820 Gong, S., Mackenzie, I.A., Zeng, G., Hess, P., Duncan, B.N., Bergmann, D.J., Szopa, S.,
821 Jonson, J.E., Keating, T.J., Zuber, A., 2012. Modelling future changes in surface ozone: a
822 parameterized approach. *Atmos. Chem. Phys* 12, 2037–2054. [https://doi.org/10.5194/acp-12-](https://doi.org/10.5194/acp-12-2037-2012)
823 2037-2012

824 Wrigley, C.W., 2009. Wheat: A unique grain for the world, in: *Wheat: Chemistry and Technology:*
825 *Fourth Edition*. AACCC International Press, pp. 1–17. [https://doi.org/10.1016/B978-1-891127-](https://doi.org/10.1016/B978-1-891127-55-7.50008-2)
826 55-7.50008-2

827 Zadoks, J.C., Board, E., 1974. A decimal code for the growth stages of cereals. *Weed Res.* 14, 415–
828 421. <https://doi.org/10.1111/j.1365-3180.1974.tb01084.x>

829 Zaka, S., Frak, E., Julier, B., Gastal, F., Louarn, G., 2016. Intraspecific variation in thermal
830 acclimation of photosynthesis across a range of temperatures in a perennial crop. *AoB Plants* 8.
831 <https://doi.org/10.1093/aobpla/plw035>

832