

## Clinical/serological outcome in humans bitten by *Babesia* species positive *Ixodes ricinus* ticks in Sweden and on the Åland Islands

Wilhelmsson, P.; Lövmär, M.; Krogfelt, Karen Angeliki; Nielsen, H.V.; Forsberg, Pia; Lindgren, P. E.

*Published in:*  
Ticks and Tick-borne Diseases

*DOI:*  
[10.1016/j.ttbdis.2020.101455](https://doi.org/10.1016/j.ttbdis.2020.101455)

*Publication date:*  
2020

*Document Version*  
Peer reviewed version

*Citation for published version (APA):*  
Wilhelmsson, P., Lövmär, M., Krogfelt, K. A., Nielsen, H. V., Forsberg, P., & Lindgren, P. E. (2020). Clinical/serological outcome in humans bitten by *Babesia* species positive *Ixodes ricinus* ticks in Sweden and on the Åland Islands. *Ticks and Tick-borne Diseases*, 11(4), Article 101455. <https://doi.org/10.1016/j.ttbdis.2020.101455>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

### Take down policy

If you believe that this document breaches copyright please contact [rucforsk@kb.dk](mailto:rucforsk@kb.dk) providing details, and we will remove access to the work immediately and investigate your claim.

## Manuscript Details

<b>Manuscript number</b>	TTBDIS_2018_332_R5
<b>Title</b>	Clinical/serological outcome in humans bitten by Babesia species positive Ixodes ricinus ticks in Sweden and on the Åland Islands
<b>Article type</b>	Research Paper

### Abstract

The risk of contracting babesiosis after a tick bite in Sweden and the Åland Islands, Finland, is unknown. We investigated clinical and serological outcomes in people bitten by Ixodes ricinus ticks positive for Babesia species. Ticks, blood and questionnaires were obtained from study participants in Sweden and on the Åland Islands. Sixty-five of 2098 (3.1%) ticks were positive by real-time PCR. Three Babesia species were detected, Babesia microti (n=33), B. venatorum (n=27) and B. capreoli (n=5), the latter species not known to cause human infection. Half (46%) of the Babesia PCR-positive ticks also contained Borrelia spp. Fifty-three participants bitten by a Babesia PCR-positive tick and a control group bitten by a Babesia PCR-negative tick were tested for B. microti IgG antibodies by IFA. The overall seroprevalence was 4.4%, but there was no significant difference between the groups. None of the participants seroconverted and no participant with a Babesia PCR-positive tick sought medical care or reported symptoms suggestive of babesiosis. Given the prevalence of Babesia in I. ricinus ticks in southern Sweden and on the Åland Islands, babesiosis should be considered a possible diagnosis in symptomatic residents who seek medical care following tick exposure.

**Keywords** Babesia; Babesiosis; Human; Seroconversion; Ixodesricinus; Co-infection

**Corresponding Author** Matilda Lövmär

**Corresponding Author's Institution** Linköpings Universitet

**Order of Authors** Peter Wilhemsson, Matilda Lövmär, Karen Krogfelt, Henrik Vedel Nielsen, Pia Forsberg, Per-Eric Lindgren

## Submission Files Included in this PDF

### File Name [File Type]

Cover letter Ticks and Tick-borne Diseases\_180827.docx [Cover Letter]

Response\_reviewers\_200416.docx [Response to Reviewers]

Additional file 1.docx [Response to Reviewers]

Additional file 2.docx [Response to Reviewers]

Revised\_Manuscript\_20200416.docx [Manuscript File]

Figure1A.jpg [Figure]

Figure1B.jpg [Figure]

Figure2.jpg [Figure]

Credit Author Statement.docx [Author Statement]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.



Linköping University

27<sup>th</sup> August 2018

Dear Editor,

Please find enclosed a manuscript entitled “*Babesia* species present in *Ixodes ricinus* ticks and clinical/serological outcome in humans after an infected tick bite in Sweden and the Åland Islands”.

I and my co-authors would be very pleased if you would consider this manuscript for publication as a Research article in the journal Ticks and Tick-borne Diseases.

Babesiosis is a tick-borne human infection in the temperate regions of North America and Eurasia. In many European countries, cases of babesiosis are not mandatorily notifiable by medical practitioners. Therefore, the number of babesiosis cases are unknown. The risk of developing a *Babesia* infection after a single tick bite is also unknown but depends likely on many factors such as developmental stage of the tick, duration of tick feeding, the *Babesia* species to be transmitted as well as the number of *Babesia* parasites in the tick and possible co-infection with other tick-borne pathogens.

In an effort to elucidate the incidence of babesiosis and to investigate how different factors influence the risk of developing a *Babesia* infection, we collected and analyzed ticks for the presence of *Babesia* spp. that had been found attached to people in Sweden and Åland Islands (Finland). At the time of the tick bite and three months later, we collected and analyzed blood samples for the presence of anti-*Babesia* antibodies from the tick-bitten people. In order to determine if participants were diagnosed with babesiosis within the three-month study period, medical records from participants that visited a health care provider were scrutinized. This study involved 1769 tick-bitten participants.

Our results indicate that the risk of contracting babesiosis after a tick bite is low, even if a *Babesia*-positive tick has been feeding for more than three days and contains up to  $10^7$  *Babesia* spp. genome copies per tick. Our findings of participants with positive serology indicate, however, that human infection with *Babesia* spp. with clinical symptoms occurs in Sweden and in the Åland Islands. Thus, babesiosis should not be neglected as a possible diagnosis in patients experiencing symptoms following a tick bite.

We are convinced that the results of this study should be of great interest to many readers of Ticks and Tick-borne Diseases. We decided to submit this manuscript to Ticks and Tick-borne



**Linköping University**

Diseases also because we consider this medical journal to be a highly appreciated scientific journal of excellent quality.

Being the corresponding author, I – and on behalf of all the authors – hereby certify that this paper is an original work. No part of this manuscript has been published previously. All authors have seen and approved the final version of the manuscript.

We very much appreciate your valuable work and look forward to your response.

Matilda Lövmär, MSc  
Corresponding author

Linköping University  
Department of Clinical and Experimental Medicine  
Division of Medical Microbiology  
SE-581 85 Linköping  
Sweden

Fax: +46(0)10-1034789

E-mail address: [matilda.lovmar@liu.se](mailto:matilda.lovmar@liu.se)

Dear Editor,

We would yet again like to thank the reviewers for the valuable comments on our manuscript TTBDIS\_2018\_332, which have helped us to further substantially improve the quality of the manuscript. We have tried to fulfill all requests and answer all of the questions and give feed-back on the comments raised by the reviewers.

In this letter, point-by-point answers follow. The comments from the reviewers have been copied into this letter and formatted with the Calibri font and put in italics. Our answers follow after each comment in the Times New Roman font.

We hope that this will help with the final decision about the manuscript.

On behalf of all authors,

Matilda Lövmär

*Enclosures:*

1. The revised manuscript with figures
2. The two questionnaires used in the TBD-STING study, translated into English

***Editor and Reviewer Comments:***

*- Section editor:*

*Accept after tidying up the last problems as indicated by reviewer 4!*

*-Managing Editor*

*The questionable 2 paragraphs can stay in the paper.*

*Minor comments:*

*Abstract: Please, write out numbers at the beginning of sentences.*

*L109-110: These references need some copy editing.*

*L161 (similar cases also elsewhere in the text): Borrelia-negative*

*L205: 36 h [space missing]*

*L226: Twenty-nine*

*L227 (also elsewhere in the text): No double full stop.*

L339: Nordström

L360-361: Please, use sentence case (no capitalisation of words) rather than title case.

L404: Bergström

L489: Nyström

All of the above mentioned comments have been corrected.

**-Reviewer 1**

- All of the issues I had raised were already addressed in the previous revision.

**-Reviewer 2**

-

**-Reviewer 3**

-

**-Reviewer 4**

-

**Major Comments**

*Line 162: are the 106 controls made of ANY and ALL Babesia and Borrelia negative samples collected during the entire study period, and matched geographically? If not, the seroprevalence, as reported, would be meaningless.*

We have changed the text slightly to clarify, see lines 157-160. The controls were chosen from all the *Babesia*- and *Borrelia*-negative ticks and geographically matched, twice as many controls were chosen as participants with positive ticks.

*Line 212: Figure 2B appears to indicate that the prevalence of B. venatorum infected ticks is HIGHER in southcentral Sweden when compared with Åland Islands. Here, the text indicates that the prevalence is HIGHER in the Åland Islands. Which is correct? This comment also applies to the sentence on lines 243-244.*

There is no figure 2B but we have assumed that the reviewers comment is regarding figure 1B. We have changed the figure text slightly to make this clearer, the statement in the manuscript is correct regarding higher prevalence in the Åland Islands. See the new figure text for Fig 1B.

**Minor Comments**

*Lines 3-4: the following title "Clinical/serological outcome in humans bitten by Babesia spp. positive Ixodes ricinus ticks in Sweden and on the Åland Islands" may be more attractive to*

*the reader, in particular when browsing PubMed.*

Thank you for this comment, we have changed the title accordingly.

*Lines 36-37: change to "Sixty-five of 2098 ticks were positive (3.1%) by real-time PCR. Three Babesia species ..."*

Changed according to comment.

*Line 39: change to "Half (46%) of the Babesia ...."*

Changed according to comment.

*Line 39: change to "Fifty-three participants ..."*

Changed according to comment.

*Lines 44-47: change to "Given the prevalence of Babesia in I. ricinus ticks in southern Sweden and on the Aland Islands, babesiosis should be considered a possible diagnosis in symptomatic residents who seek medical care following tick exposure."*

Changed according to comment, we have also altered the conclusion where a similar phrasing was used, see lines 318-322.

*Line 53: change to "... transmitted by several tick ..."*

Changed according to comment.

*Line 55: change to "The first documented case of ..."*

Changed according to comment.

*Lines 56-57: change to "... was reported in 1957 from Yugoslavia (Skrabalo and Deanovic, 1957). Other cases followed in Western Europe ..."*

Changed according to comment.

*Line 65: change to "... include fever, malaise, chills, sweats, headache and myalgia ..."*

Changed according to comment.

*Lines 71-73: Add to the citations the report by Moniusko-Malinowska et al. in Infectious Diseases vol 48, pp.537-543, 2016.*

This reference has been added to the citations.

*Line 81: delete "In Sweden,"*

Changed according to comment.

*Line 84: delete "However,"*

Changed according to comment.

*Line 86: delete "previously"*

Changed according to comment.

*Line 87: modify to "Co-infection with Babesia and Borrelia spp. has been documented ..."*

Changed according to comment.

*Line 90: delete the number 3, unless it means something.*

Deleted according to comment.

*Line 92: replace “intensified” with “worse”.*

Changed according to comment.

*Lines 96: delete this part of the sentence “relative to the ..... were used”, and use the edited sentence as the last sentence of the previous paragraph.’*

We deleted the last part of one sentence according to this comment and moved the sentence to the previous paragraph. The last sentence of the paragraph was moved to the Material and Methods section. See lines 93-101.

*Line 104: rephrase as “(PHCs) in the three regions of ...”*

Changed according to comment.

*Line 114: delete “sera from”.*

Changed according to comment.

*Line 115: delete “serologically”.*

Changed according to comment.

*Line 116: delete “and analyzed for the presence of Babesia spp. using real-time PCR” because the current version of the manuscript no longer includes data on Babesia spp. detected by PCR in human blood samples.*

Changed according to comment.

*Lines 119-120: rephrase to “In this study, cDNAs from 2098 ticks detached from 1769 participants were analyzed whereas cDNAs from the remaining 12 ticks were not available for analysis.”*

Changed according to comment.

*Lines 123-124: delete “, in total 5 uL cDNA per well.”*

Changed according to comment.

*Line 141: replace “Amplification” with “Extension”.*

Changed according to comment.

*Line 157: rephrase as “... from 53 of the 61 participants bitten by Babesia positive ticks were ...”*

Changed according to comment.

*Line 165: start sentence as “Samples were ...”.*

Changed according to comment.

*Line 182: replace with “Statistical Analysis”*

Changed according to comment.

*Lines 198-199: delete “, as compared with B. capreoli sequence (AY26009) deposited in Genebank,”. This info should be moved to the Methods section.*



We have left some of this information because it is a result from our study and as such should be presented in the results. Because of this comment we have rephrased it slightly, see lines 193-196.

*Lines 206-207: move the sentence "Different ticks ...Babesia spp." to after the next sentence.*  
Changed according to comment.

*Lines 213-214: rephrase to "No other differences in species composition between regions were significant".*  
Changed according to comment.

*Line 224: rephrase to "Seven participants ...".*  
Changed according to comment.

*Line 252: rephrase to "... in 0.6% of ticks (Jensen et al. 2017)".*  
Changed according to comment.

*Line 256: change to "...positive larvae among questing ticks, ...".*  
Changed according to comment.

*Line 259: change to "... B. capreoli is known to be ...".*  
Changed according to comment.

*Line 280: delete "on seroprevalence"*  
Changed according to comment.

*Line 282: remove "antibodies".*  
Changed according to comment.

## To participants of the STING-study

Please answer all questions!

### When did you notice that you had been tick-bitten?

Year-Month-Day: \_\_\_\_\_ Time \_\_\_\_\_

### When do you think you were tick-bitten?

Year-Month-Day: \_\_\_\_\_ Time \_\_\_\_\_

### Where do you think you were when you were tick-bitten? Please state the name of the municipality.

\_\_\_\_\_

### What kind of habitat (vegetation type) had you visited?

Lake/Sea  Forest  Garden  Lawn

Other: \_\_\_\_\_

### When was the tick removed?

Year-Month-Day: \_\_\_\_\_ Time \_\_\_\_\_

### Where on the body was the tick attached?

\_\_\_\_\_

Did you remove the whole tick? Yes  No  Do not know

Have you had any other tick bites this season? Yes  No  Do not know

If Yes, how many? 1-4  5-9  >10

**Have you ever been treated for the tick-borne infection Borrelia?**

Yes  No  Do not know  If Yes; Year–Month–Day \_\_\_\_\_

**Did you receive any medicine?**

Yes  No  Do not know  If Yes; what kind of medicine did you get?  
\_\_\_\_\_

**Have you ever been treated for “Erythema migrans”?**

(Erythema migrans = red ring-like or homogenous expanding rash.)

Yes  No  Do not know  If Yes; Year–Month–Day \_\_\_\_\_

**Did you then receive any medicine to treat the infection?**

Yes  No  Do not know  If Yes; what kind? \_\_\_\_\_

**Have you ever been treated for the tick-borne infection “Ehrlichia”  
(= Ehrlichiosis, also called “Anaplasma” or anaplasmosis)?**

Yes  No  Do not know  If Yes; Year–Month–Day \_\_\_\_\_

**Did you receive any medicine to cure the Ehrlichia (Anaplasma) infection?**

Yes  No  Do not know  If Yes; what kind? \_\_\_\_\_

**Have you ever been treated for the tick-borne infection TBE?**

(TBE is a viral disease which sometimes causes disease in the central nervous system.)

Yes  No  Do not know

If Yes; Year-Month-Day \_\_\_\_\_

**Did you receive any medicine?**

Yes  No  Do not know

If Yes; what kind? \_\_\_\_\_

**Do you have any of the following diseases?**

**Asthma**

Yes  No  Do not know

**Allergy**

Yes  No  Do not know

**Diabetes**

Yes  No  Do not know

**Tumour-related**

Yes  No  Do not know

**Are you on medication?**

Yes  No

If Yes; what kind of medicine?

\_\_\_\_\_

**Do you smoke?**

Yes  No

Stopped smoking  Year \_\_\_\_\_

If Yes, how many cigarettes per week? \_\_\_\_\_

How many years have you smoked? \_\_\_\_\_

**Do you have any pets?**

Yes  No

**Dog**

Yes  No

**Cat**

Yes  No

**Bunny (rabbit)**

Yes  No

**Other:** \_\_\_\_\_

**Have you been vaccinated against TBE?**

Yes  No  Do not know

If Yes; Year-Month-Day\_\_\_\_\_

**Have you been vaccinated against Yellow fever?**

Yes  No  Do not know

If Yes; Year-Month-Day\_\_\_\_\_

**Have you been vaccinated against Japanese encephalitis?**

Yes  No  Do not know

If Yes; Year-Month-Day\_\_\_\_\_

**Thank you for your answers!**

**Dear STING participant!**

Three months have passed since you initiated your participation in the Tick-Borne Diseases STING-study. We previously received blood samples from you and a filled in questionnaire. Now, we need a follow-up blood sample. Therefore, you are requested to visit your primary health care centre at \_\_\_\_\_, week \_\_\_\_, Monday, Tuesday, Wednesday, or Thursday, between \_\_\_\_\_ and \_\_\_\_\_ a clock.

If you had any additional tick-bites since your study initiation and if you have collected the ticks in the tube with yellow cork, the please take that tube with you to the blood-sampling.

We would also like to know if you have had any symptoms related to tick-borne diseases during the study period. Please answer the following three questions and write your name, birth date and telephone number on the next page. We might contact you if you reported symptoms. Take this paper to your primary health care centre when you go for the sample-taking.

**1) Have you had any additional tick-bites since the first sample-taking?**

Yes  No  Do not know

If Yes; when? Year-Month-Day: \_\_\_\_\_

**2) How have you been feeling in general since the first sample-taking?**

Have you been feeling good/as usual?

Yes  No  Do not know

If No; please report if you had any of the following symptoms:

Headache Yes  No

Fatigue Yes  No

Fever, 38° or higher Yes  No

Neck pain Yes  No

Loss of appetite Yes  No

Nausea Yes  No

Weight loss Yes  No

Vertigo Yes  No

Concentration difficulties Yes  No

Radiating pain Yes  No

Muscle or joint pain Yes  No

Numbness Yes  No

**3) If you reported any symptoms in question 2, did the symptoms appear before or after any additional tick-bites?**

Before additional tick-bite Yes  No  Do not know

After additional tick-bite Yes  No  Do not know

**4) If you reported any symptoms in question 2, did you visit your primary health care centre due to the symptoms?**

Yes  No

**5) If you reported any symptoms in question 2, how many days did the symptoms last?**

\_\_\_\_\_

**Thanks for your answers!**

**Please make sure you answered every question!  
Bring this paper to your new sample-taking!**

**Name:** \_\_\_\_\_

**Date of birth:** \_\_\_\_\_

**Telephone number:**

**Home** \_\_\_\_\_

**Work** \_\_\_\_\_

**Mobile** \_\_\_\_\_

**Best regards**

Lotta Lindvall

Forskningssköterska

Infektionskliniken

Universitetssjukhuset

581 85 Linköpings

Tfn xxx/xxx xxxx

2020-04-16

1  
2  
3  
4 1 **TITLE PAGE**

5  
6 2 **Title**

7  
8  
9 3 Clinical/serological outcome in humans bitten by *Babesia* species positive *Ixodes*  
10  
11 4 *ricinus* ticks in Sweden and on the Åland Islands

12  
13 5 **AUTHORS**

14  
15  
16 6 Wilhelmsson P<sup>1,2\*</sup>

17  
18 7 Lövmär M<sup>1\*</sup>

19  
20  
21 8 Krogfelt KA<sup>3,4</sup>

22  
23 9 Nielsen HV<sup>3</sup>

24  
25 10 Forsberg P<sup>5</sup>

26  
27 11 Lindgren PE<sup>1,2</sup>

28  
29  
30 12 <sup>1</sup>Division of Medical Microbiology, Department of Biomedical and Clinical Sciences,  
31  
32 Linköping University, Linköping, Sweden

33  
34 14 matilda.lovmar@liu.se

35  
36  
37 15 <sup>2</sup> Division of Medical Services, Department of Clinical Microbiology, Ryhov County  
38  
39 Hospital, Jönköping, Sweden

40  
41 17 per-eric.lindgren@liu.se

42  
43 18 <sup>3</sup>Department of Bacteria Parasites and Fungi, Statens Serum Institute, Copenhagen, Denmark

44  
45 19 [kak@ssi.dk](mailto:kak@ssi.dk), [hvn@ssi.dk](mailto:hvn@ssi.dk)

46  
47 20 <sup>4</sup> Department of Science and Environmental, Roskilde University, Denmark

48  
49  
50 21 <sup>5</sup>Division of Infectious Diseases, Department of Biomedical and Clinical Sciences, Linköping  
51  
52 University, Linköping, Sweden

53  
54 23 pia.forsberg@liu.se



24 \* these authors contributed equally to this article

25 \*Corresponding author: Matilda Lövmär, Email: matilda.lovmar@liu.se

26 Linköping University, Department of Biomedical and Clinical Sciences, Division of Medical

27 Microbiology, 581 85 Linköping Phone: (+46) 101032054

28

29

30

### 31 **ABSTRACT**

32 The risk of contracting babesiosis after a tick bite in Sweden and the Åland Islands, Finland,  
33 is unknown. We investigated clinical and serological outcomes in people bitten by *Ixodes*  
34 *ricinus* ticks positive for *Babesia* species. Ticks, blood and questionnaires were obtained  
35 from study participants in Sweden and on the Åland Islands. Sixty-five of 2098 (3.1%) ticks  
36 were positive by real-time PCR. Three *Babesia* species were detected, *Babesia microti*  
37 (n=33), *B. venatorum* (n=27) and *B. capreoli* (n=5), the latter species not known to cause  
38 human infection. Half (46%) of the *Babesia* PCR-positive ticks also contained *Borrelia* spp.  
39 Fifty-three participants bitten by a *Babesia* PCR-positive tick and a control group bitten by a  
40 *Babesia* PCR-negative tick were tested for *B. microti* IgG antibodies by IFA. The overall  
41 seroprevalence was 4.4%, but there was no significant difference between the groups. None  
42 of the participants seroconverted and no participant with a *Babesia* PCR-positive tick sought  
43 medical care or reported symptoms suggestive of babesiosis. Given the prevalence of *Babesia*  
44 in *I. ricinus* ticks in southern Sweden and on the Åland Islands, babesiosis should be  
45 considered a possible diagnosis in symptomatic residents who seek medical care following  
46 tick exposure.

47

## 48 Keywords

49 *Babesia*; Babesiosis; Human; Seroconversion; *Ixodes ricinus*; Co-infection

51 **INTRODUCTION**

52 Human babesiosis is caused by parasites of the genus *Babesia* and transmitted by several tick  
53 species (Vannier and Krause, 2012). There are more than 100 known *Babesia* spp. that infect  
54 animals but only a few are known to infect humans. The first documented case of human  
55 babesiosis in Europe was reported in 1957 from Yugoslavia (Skrabalo and Deanovic, 1957).  
56 Other cases followed in Western Europe, including Scandinavia (Haapasalo et al., 2010;  
57 Morch et al., 2015) and a travel-associated case in Denmark (Holler et al., 2013). In Sweden  
58 there have been two reported cases of human babesiosis, both in splenectomized patients  
59 (Bläckberg et al., 2018; Uhnoo et al., 1992). In North America, babesiosis is considered an  
60 emerging health threat that is expanding into new geographical areas and may be overlooked  
61 by clinicians in regions not previously considered endemic (Gray and Herwaldt, 2019). Most  
62 cases in the United States have been reported in immunocompetent patients (Vannier et al.,  
63 2015). The first case in the United States was reported in 1966 (Scholtens et al., 1968).

64           Symptoms of babesiosis include fever, malaise, chills, sweats, headache and  
65 myalgia accompanied by anemia, leukopenia or leukocytosis, thrombocytopenia and elevated  
66 hepatic enzymes (Vannier and Krause, 2012). In Europe the most common cause of human  
67 babesiosis is *Babesia divergens*, which typically is diagnosed in immunocompromised  
68 individuals and gives rise to a severe illness (Vannier et al., 2015; Vannier and Krause, 2012).  
69 Infection with *B. divergens* has also been reported in immunocompetent patients (Martinot et  
70 al., 2011). A few cases of *B. microti* and *B. venatorum* infection have been reported in Europe  
71 (Bläckberg et al., 2018; Blum et al., 2011; Haselbarth et al., 2007; Herwaldt et al., 2003;

178  
179  
180 72 Hildebrandt et al., 2007; Moniuszko-Malinowska et al., 2016). Infection with *B. venatorum*,  
181  
182 73 giving mild to severe symptoms in splenectomized patients, has been reported (Haselbarth et  
183  
184 74 al., 2007; Herwaldt et al., 2003). In the United States, *B. microti* is the most common causative  
185  
186 75 agent of babesiosis; it causes mild to moderate symptoms and subclinical infections in  
187  
188 76 immunocompetent persons (Vannier et al., 2015; Vannier and Krause, 2012). However, severe  
189  
190 77 babesiosis may also occur, even in previously apparently healthy individuals (Gray and  
191  
192 78 Herwaldt, 2019; Hatcher et al., 2001).

195  
196 79 In Sweden, the prevalence of *Babesia* spp. in questing *I. ricinus* ticks was  
197  
198 80 recently estimated to be 4.4% and included *B. microti* (3.2%), *B. venatorum* (1.0%) and *B.*  
199  
200 81 *divergens* (0.2%) (Karlsson and Andersson, 2015). *B. capreoli* has been reported in roe deer,  
201  
202 82 but this *Babesia* sp. is not known to cause human infections (Andersson et al., 2016;  
203  
204 83 Malandrin et al., 2010). In Italy, the prevalence of *B. venatorum* in ticks that have bitten  
205  
206 84 humans was estimated to be 0.6% (Otranto et al., 2014). To our knowledge, the clinical  
207  
208 85 outcome and the rate of seroconversion after a bite by a *Babesia* containing tick or a tick co-  
209  
210 86 infected with *Borrelia*, have not been investigated.

212  
213 87 Co-infection with *Babesia* and *Borrelia* spp. has been documented several times  
214  
215 88 (Diuk-Wasser et al., 2016; Knapp and Rice., 2015). There are differing opinions on the  
216  
217 89 consequences of co-infection for severity of symptoms and prognosis in humans. According  
218  
219 90 to Diuk-Wasser et al., (2016), the severity and duration of symptoms are greater in co-  
220  
221 91 infected patients and there are indications that co-infection may result in altered or  
222  
223 92 suppressed immune response, which in turn leads to worse pathogenesis; more research is,  
224  
225 93 however, needed. The aims of the present study were to investigate the prevalence of *Babesia*  
226  
227 94 spp. in ticks that had bitten humans and to evaluate the concomitant risk of clinical babesiosis  
228  
229 95 or subclinical seroconversion against *Babesia* spp.

## 96 MATERIAL AND METHODS

### 97 TBD STING-study

98 Ticks, serum and questionnaires from the Tick-Borne Diseases (TBD) STING-study were  
99 used (Fryland et al., 2011; Grankvist et al., 2015; Henningson et al., 2015; Henningson et  
100 al., 2016; Lindblom et al., 2014; Wilhelmsson et al., 2010; Wilhelmsson et al., 2013a;  
101 Wilhelmsson et al., 2013b).

102 Ticks and blood samples were collected during 2008–2009 at 34 primary healthcare centers  
103 (PHCs) in the three regions of Sweden (Northern Sweden, South Central Sweden and  
104 Southernmost Sweden) and on the Åland Islands, Finland (Figure 1A). Only  
105 immunocompetent tick-bitten individuals  $\geq 18$  years were included. Questionnaires, ethical  
106 approval, collection, transport and storage of ticks and blood samples, determination of  
107 developmental stage and feeding time of ticks, and extraction and treatment of nucleic acids  
108 have been described previously (Andersson et al., 2016; Wilhelmsson et al., 2013a;  
109 Wilhelmsson et al., 2013b). *Babesia* spp. were detected using real-time PCR-based  
110 amplification of reverse-transcribed total nucleic acids.

111 Blood samples were collected from participants within three days of the tick  
112 bite and three months later (Figure 2). Serum samples from participants with *Babesia* PCR-  
113 positive ticks and a negative control group, consisting of participants bitten by a *Babesia*  
114 PCR-negative tick, geographically matched, were analyzed by IFA.

### 115 Detection of *Babesia* spp. in ticks using SYBR-green real-time PCR assay

116 A total of 2110 ticks detached from 1770 participants were delivered from PHCs during  
117 2008–2009. In this study, cDNAs from 2098 ticks detached from 1769 participants were  
118 analyzed whereas cDNAs from the remaining 12 ticks were not available for analysis.

296  
297  
298  
299 119 The 2098 cDNA samples of the individually extracted ticks were grouped into  
300 120 pools of five, i.e. one  $\mu\text{l}$  of cDNA from each tick was used per well. Samples from the  
302 121 positive pools were individually analyzed using 2  $\mu\text{l}$  cDNA. The PCR mixture contained 10  
304 122  $\mu\text{l}$  SYBR-green (Thermo Scientific, Helsingborg, Sweden), 0.4  $\mu\text{l}$  of each primer (10  $\mu\text{M}$ ),  
307 123 BJ1: 5'-GTC TTG TAA TTG GAA TGA TGG-3' (Invitrogen™, Thermo Scientific) and  
309 124 BN2: 5'-TAG TTT ATG GTT AGG ACT ACG-3' (Casati et al., 2006), 4.2  $\mu\text{l}$  RNase free  
311 125 water. Primers BJ1 and BN2 were designed to target the *Babesia 18S* rRNA gene to amplify  
313 126 a 411 to 452 bp long amplicon depending on the species of *Babesia*. For species  
315 127 determination, sequencing of the amplicon was carried out (see below). As a positive control,  
317 128 2  $\mu\text{l}$  *B. microti* DNA (~10 ng/ $\mu\text{l}$ ), extracted from *I. ricinus* ticks collected in Slovakia (kindly  
320 129 provided by Dr Bronislava Vichová, through Dr Martin Andersson), and 2  $\mu\text{l}$  of a synthetic  
322 130 plasmid preparation was used. The plasmid contained the target sequence of the SYBR green  
324 131 real-time PCR assay, spanning the nucleotides 467-955 of the *B. divergens 18S* rRNA gene  
326 132 (acc. no AJ439713), synthesized and cloned in a pUC57 vector (Genscript USA Inc, NJ). The  
328 133 non-template control consisted of 10.8  $\mu\text{l}$  PCR mixture and 9.2  $\mu\text{l}$  RNase free water. The  
330 134 SYBR-green real-time PCR on *Babesia* were performed using C1000™ Thermal Cycler,  
332 135 CFX96™ system (BioRad Laboratories, Inc., Hercules, CA). The PCR run was initiated by a  
334 136 denaturation step at 94°C for 10 min followed by 35 cycles of denaturation at 94°C for 1 min,  
336 137 annealing at 55°C for 1 min and elongation at 72°C for 2 min. Extension was completed by a  
338 138 further step at 72°C for 5 min, and melt curve analysis was performed (Casati et al., 2006).

341 139

342

343

344 140 ***Babesia* spp. identification by sequencing**

345

346 141 Samples positive for *Babesia* spp. in the real-time PCR assay were sent to Macrogen Inc.

347

348 142 (Amsterdam, the Netherlands) for nucleotide sequencing. The reactions were based on

349

350 143 BigDye chemistry. Chromatograms were edited using BioEdit Sequence Alignment Editor

351

352

353

354

355  
356  
357 144 version 7.2.5 (Tom Hall, Ibis Therapeutics, Carlsbad, CA) and sequences examined using the  
358  
359 145 Basic Local Alignment Tool (BLAST). Sequences obtained have been deposited in GenBank  
360  
361  
362 146 with accession numbers ranging from MH351680 to MH351744.

363  
364 147 Species determination for *B. microti* and *B. venatorum* is possible by sequencing the  
365  
366 148 amplicon from the real-time PCR assay. To fully distinguish between the two genetically  
367  
368 149 similar *Babesia* spp., *B. divergens* and *B. capreoli*, three sets of primers were used to amplify  
369  
370 150 and sequence the complete 18S rRNA gene from all samples positive for these species, as  
371  
372  
373 151 earlier described (Malandrin et al., 2010).

### 374 375 376 152 **Detection of *Babesia microti* IgG antibodies in human serum**

377  
378 153 The first and second serum samples (collected at recruitment of participants and three months  
379  
380 154 later, respectively) from 53 of the 61 participants bitten by *Babesia*-positive ticks were  
381  
382 155 analyzed for the presence of *B. microti* IgG antibodies, using an indirect immunofluorescence  
383  
384 156 assay (IFA); (Focus Diagnostics, Cypress, CA). For the remaining eight participants, serum  
385  
386 157 was not available for analysis, since it had been used for other analyses. Twice as many  
387  
388 158 controls were selected; first sample sera from participants bitten by *Babesia*- and *Borrelia*-  
389  
390 159 negative ticks were matched geographically and used as controls (n=106) for analysis  
391  
392 160 regarding IgG antibodies against *B. microti*. For *B. venatorum* there were no commercial kits  
393  
394 161 available at the time of the study. IFA titers  $\geq 1:64$  were defined as positive. A cut-off value  
395  
396 162 of 1:64 was used in accordance with previous research (Johnson et al., 2009). Samples were  
397  
398 163 diluted to determine the highest positive titer. The IFA-slides were analyzed by two  
399  
400 164 researchers independently, samples were defined as positive when both researchers found  
401  
402 165 them positive. For the diagnosis of on-going or recent *Babesia* spp. infection, at least a four-  
403  
404 166 fold rise of the IFA titer (Krause, 2003; Vannier and Krause, 2012) was required when first  
405  
406 167 and second sera were tested simultaneously.  
407  
408  
409  
410  
411  
412  
413

414  
415  
416 **168 Self-reported symptoms in the questionnaires of the TBD STING-study and medical**  
417  
418 **169 records**

420  
421 170 The questionnaires from participants with *Babesia* PCR-positive ticks were scrutinized for  
422  
423 171 symptoms suggestive of babesiosis, i.e. chills, fever, headache, nausea, myalgia, malaise,  
424  
425 172 weight loss, arthralgia and lack of appetite (Vannier and Krause, 2012). Other symptoms  
426  
427 173 indicating TBD in the questionnaires were neck pain, vertigo, concentration difficulties,  
428  
429 174 numbness, radiating pain. If participants sought medical care during the three-month study  
430  
431 175 period, the medical records were obtained and scrutinized for symptoms of babesiosis and/or  
432  
433 176 if they were diagnosed with babesiosis or another TBD.

434  
435  
436 **177 Co-infection of tick-borne pathogens in ticks**

437  
438  
439 178 *Borrelia*-data from the TBD STING-study were used to determine which ticks were co-  
440  
441 179 infected with both *Babesia* spp. and *Borrelia* spp. (Wilhelmsson et al., 2013a).

442  
443  
444 **180 Statistical analysis**

445  
446  
447 181 The Chi square test was applied to compare prevalence of *Babesia* spp. between regions and  
448  
449 182 between tick developmental stages, but when the expected frequency was  $< 5$  in at least one  
450  
451 183 of the cells of the contingency table, we used Fisher's exact test. Statistical analyses were  
452  
453 184 performed using GraphPad Prism version 6.04 for Windows (GraphPad Software, San Diego,  
454  
455 185 CA). P-values  $< 0.05$  were considered significant.

456  
457  
458 **186 RESULTS**

459  
460  
461 **187 *Babesia* species in ticks detached from humans**

462  
463 188 A total of 2098 ticks that had bitten humans were analyzed (Table 1). Sixty (2.9%) of the  
464  
465 189 ticks were damaged to the extent that neither developmental stage nor the species could be  
466  
467 190 determined. The remaining 2038 ticks were identified as *I. ricinus*; 86 (4.1%) larvae, 1466

473  
474  
475 191 (70%) nymphs, and 486 (23%) adults. 65 of 2098 ticks (3.1%) were positive for *Babesia* spp.  
476  
477 192 by real-time PCR assay. Based on nucleotide sequencing of the PCR products, three *Babesia*  
478  
479 193 species were detected; *B. microti* (n=33), *B. venatorum* (n=27), and *B. capreoli* (n=5). Our  
480  
481 194 analysis of the complete 18S rDNA sequences for the five samples initially determined as  
482  
483 195 either *B. divergens* or *B. capreoli* revealed a signature typical of *B. capreoli* at positions 631,  
484  
485 196 663 and 1637 (GTT), as compared with *B. capreoli* sequence (GeneBank: AY26009). In our  
486  
487 197 study there was no significant difference in *Babesia* prevalence between adult ticks and  
488  
489 198 nymphs or between nymphs and larvae. However, the prevalence of *B. microti* was  
490  
491 199 significantly higher in adult ticks than in nymphs ( $p < 0.05$ ).  
492  
493

494 200 The 65 PCR-positive ticks were collected from 61 participants and the duration  
495  
496 201 of tick feeding could be estimated for 58 of the 65 ticks (Appendix). 21 of these ticks had  
497  
498 202 been feeding for  $> 36$  h and the remaining 37 ticks  $< 36$  h. There were three participants  
499  
500 203 bitten by more than one infected tick. One participant from the Åland Islands was bitten by  
501  
502 204 three infected ticks, one from Southcentral Sweden was bitten by two infected ticks, and one  
503  
504 205 from Southernmost Sweden was bitten by two infected ticks (Appendix). Different ticks from  
505  
506 206 the same participant contained different *Babesia* spp. There was no significant difference in  
507  
508 207 prevalence of infected ticks between the geographic regions (Figure 1B). However, there was  
509  
510 208 a significant difference in species composition of *Babesia* between the geographic regions,  
511  
512 209 with *B. venatorum* more prevalent in the Åland Islands than in Southcentral Sweden  
513  
514 210 ( $p < 0.03$ ). No other differences in species composition between the regions were significant.  
515  
516  
517

#### 518 211 **Co-infection with *Babesia* spp. and *Borrelia* spp. in ticks**

519  
520  
521 212 To find which ticks contained *Borrelia* spp., data from the TBD STING-study were used  
522  
523 213 (Wilhelmsson et al., 2013a). Thirty out of 65 (46%) *Babesia* spp. positive ticks contained  
524  
525 214 *Borrelia* spp. (Appendix). There was a significant difference in the frequency of co-infections  
526  
527  
528  
529  
530  
531



215 involving *B. microti* and *Borrelia* spp. (60%) and the frequency of co-infections involving *B.*  
216 *venatorum* and *Borrelia* spp. (30%) ( $p < 0.01$ ).

### 217 **Seroprevalence, seroconversion and reported symptoms of the tick-bitten participants**

218 The overall seroprevalence for *B. microti* was 7 out of 159 (4.4%) with no significant  
219 difference between the participants bitten by *Babesia* PCR-positive ticks and participants  
220 bitten by *Babesia* PCR-negative ticks. Seven participants were found seropositive both in  
221 first and second sample sera (Table 2).

222           Twenty-nine participants were bitten by ticks containing both *Babesia* spp. and  
223 *Borrelia* spp. One of these participants was bitten by two ticks positive for *Babesia* spp. and  
224 *Borrelia* spp. Data regarding seroconversion for *Borrelia* spp. showed that one of the  
225 participants who seroconverted to *Borrelia* had a *Babesia* PCR-positive tick. However, this  
226 participant had a negative *B. microti* serology.

227           There were ten participants with *Babesia* PCR-positive ticks who reported  
228 symptoms in their questionnaires. The symptoms included headache, muscle pain, fatigue,  
229 neck pain, dizziness, concentration difficulties, numbness, radiating pain, joint pain and  
230 nausea. Only one of these participants had *B. microti* IgG antibodies. This participant sought  
231 medical care at the PHC during the study period but according to notes found in the medical  
232 records, symptoms were deemed not relevant to the tick bite.

## 233 **DISCUSSION**

234 To our knowledge this is the first time *Babesia* has been found in ticks that have bitten  
235 humans in Sweden. In total, 3.1% of ticks collected from four regions in Sweden and in the  
236 Åland Islands contained *Babesia* spp. *Babesia* infected ticks were found in three of the four  
237 regions studied (Fig. 1B). A previous study on the prevalence of *Babesia* spp., in questing  
238 ticks in southern Sweden, found a prevalence of 4.4% which is in line with our findings

591  
592  
593 239 (Karlsson and Andersson, 2015). Comparing prevalence of *Babesia* spp. in the different  
594  
595 240 regions, *B. venatorum* was more prevalent on the Åland Islands than in Southcentral Sweden.  
596  
597 241 One could speculate the difference is related to different *Babesia* reservoir host composition.  
598  
599  
600 242 We found that the most prevalent species in ticks collected from  
601  
602 243 humans was *B. microti*, followed by *B. venatorum* and *B. capreoli*. In Sweden *B. capreoli* has  
603  
604 244 been found in 44% of roe deer (Andersson et al., 2016), but this species is not known to cause  
605  
606 245 human infections (Malandrin et al., 2010). We did not find any samples positive for *B.*  
607  
608 246 *divergens*. A study conducted in Norway found that 0.1% of field-collected ticks are infected  
609  
610 247 with *B. divergens*, 0.1% with *B. capreoli* and 0.6% with *B. venatorum* (Øines et al., 2012). In  
611  
612 248 Denmark, *B. divergens* was found in 1.9% of ticks and *B. venatorum* in 0.6% of ticks (Jensen  
613  
614 249 et al., 2017). Karlsson and Andersson (2015) found 0.2% questing ticks containing *B.*  
615  
616 250 *divergens* in Sweden and found no significant difference in prevalence of *Babesia* spp. in  
617  
618 251 adult ticks compared to nymphs. They found a higher prevalence in nymphs than in adults  
619  
620 252 and they found no positive larvae among questing ticks, but we analyzed four times as many  
621  
622 253 ticks and they were collected from humans. We found only one positive larva (Table 1) that  
623  
624 254 contained *B. capreoli*. *B. capreoli* has not previously been found in larvae, however, *B.*  
625  
626 255 *divergens* which is genetically similar to *B. capreoli* is known to be transovarially transmitted  
627  
628 256 (Bonnet et al., 2007).

631  
632 257 Nearly one-half of the ticks positive for *Babesia* spp. were positive for *Borrelia*  
633  
634 258 spp. Sixty percent of ticks containing *B. microti* were co-infected with *Borrelia* spp.  
635  
636 259 compared to 30% of ticks containing *B. venatorum*. This may reflect the genus ratio of  
637  
638 260 *Babesia* spp. and *Borrelia* spp. in reservoir hosts. It has been observed in an experimental  
639  
640 261 model that mice co-infected with a strain of *B. microti* and an invasive strain of *B.*  
641  
642 262 *burgdorferi* s.s. increased *B. microti* frequency in *I. scapularis* that fed from them compared  
643  
644  
645  
646  
647  
648  
649

650  
651  
652 263 to mice infected with *B. microti* alone (Dunn et al., 2014). It remains to be confirmed if this  
653  
654 264 finding applies to *I. ricinus* and one or several *Babesia* spp. it can transmit.  
655  
656

657 265                 Seven participants were positive for *B. microti* antibodies. Since  
658  
659 266 none of them seroconverted and the antibody titers were low, we did not suspect an ongoing  
660  
661 267 infection (Vannier and Krause, 2012). It is probable that they had a previous or subclinical  
662  
663 268 infection and still carried antibodies. In a previous study, Lempereur et al. (2015) found no  
664  
665 269 antibody cross-reactivity between *B. microti* and *B. venatorum* used as IFA antigens, but IgM  
666  
667 270 cross reactivity between *B. microti* and *B. divergens* has been observed in another study  
668  
669 271 (Haselbarth et al., 2007). Bläckberg et al. (2018) reported that a patient with *B. venatorum*-  
670  
671 272 infection had *B. divergens* IgG antibodies. For the participants with a positive *B. microti*  
672  
673 273 serology, only three had been bitten by *Babesia* PCR-positive ticks during the study period.  
674  
675 274 Because they had the same antibody titers in the first serum sample as in the second, it is not  
676  
677 275 likely that the tick-bite during the study period was the cause of the positive serology or that  
678  
679 276 the participants had an ongoing infection. One study in southern Sweden has revealed a  
680  
681 277 prevalence of 16.3% for *B. microti* and *B. divergens* antibodies in a geographically selected  
682  
683 278 cohort of seropositive *Borrelia* s.l. patients; and a 2.5% prevalence in a healthy control group  
684  
685 279 (Svensson et al., 2019). However, comparing this study to ours is complicated, since we have  
686  
687 280 used another serological assay for *Babesia* antibodies and the study populations differ from  
688  
689 281 each other.  
690  
691  
692  
693

694 282                 In total, three participants were bitten by more than one *Babesia* PCR-positive  
695  
696 283 tick, none of them developed antibodies against *B. microti* during the study period. One of  
697  
698 284 these participants was bitten by ticks containing *B. capreoli*, not confirmed to be human  
699  
700 285 pathogenic. Since we only analyzed for *B. microti* IgG we cannot draw any general  
701  
702 286 conclusions regarding *Babesia* spp. antibodies. However, it is an interesting future  
703  
704 287 prospective. None of the participants with more than one positive tick reported symptoms in  
705  
706  
707  
708

709  
710  
711 288 their questionnaires or sought medical care during the study period. This suggests a low risk  
712  
713 289 of transmission despite being bitten by several positive ticks and different species.  
714  
715 290 Furthermore, the efficacy of transmission has been shown to correlate with the duration of  
716  
717 291 tick feeding (for *I. scapularis*) in hamsters and white-footed mice, with infection rates close  
718  
719 292 to 100% if the tick is allowed to feed to repletion (Piesman and Spielman, 1980). However, it  
720  
721 293 is not known if this is true for *I. ricinus* and/or human hosts. In our study ticks were removed  
722  
723 294 by the participant before repletion.  
724  
725

726 295 Ten participants with *Babesia* PCR-positive ticks reported symptoms in their  
727  
728 296 questionnaires. These symptoms were nonspecific and might indicate different conditions,  
729  
730 297 babesiosis included. Only one participant bitten by a *Babesia* PCR-positive tick sought  
731  
732 298 medical care during the study period. According to medical records, symptoms were  
733  
734 299 unrelated to the tick bite. Thus, we conclude that none of the participants suffered from  
735  
736 300 symptoms of babesiosis.  
737  
738

739 301 Twenty-nine of 61 participants with *Babesia* PCR-positive ticks were bitten by  
740  
741 302 ticks co-infected with *Borrelia*. Of the ten participants who reported symptoms, five had been  
742  
743 303 bitten by ticks positive for both *Babesia* spp. and *Borrelia* spp. The symptoms reported could  
744  
745 304 be attributed either to babesiosis, borreliosis or other infections. None of the participants who  
746  
747 305 reported symptoms, with the exception of the participant mentioned above, had sought  
748  
749 306 medical care. This suggests that no participant suffered from severe illness.  
750  
751

752 307 One of the potential limitations of this study is that we did not test the serum for  
753  
754 308 *B. venatorum* antibodies since we did not have available commercial kits for these analyses.  
755  
756 309 The conclusions that can be drawn from the serological analyses are further limited by the  
757  
758 310 lack of information regarding travel history, since this was not included in the questionnaires,  
759  
760 311 designed for the TBD STING-study. Unfortunately, whole blood samples from the TBD  
761  
762 312 STING-study were not available for real-time PCR analysis.  
763  
764  
765  
766  
767

**313 Conclusions**

314 In conclusion our results indicate that immunocompetent individuals have a low risk of  
315 developing severe babesiosis after an *I. ricinus* tick bite in Sweden and on the Åland Islands,  
316 particularly when the tick has been feeding for less than 36 hours. Our findings of  
317 participants with positive serology suggests that human infection with *B. microti* occurs in  
318 Sweden, although we do not know about the travel history of these participants. Given the  
319 prevalence of *Babesia* in *I. ricinus* ticks as well as the seroprevalence of *Babesia* antibodies  
320 among residents in southern Sweden and on the Åland Islands, babesiosis should be  
321 considered a possible diagnosis in symptomatic residents who seek medical care following  
322 tick exposure.

**324 ACKNOWLEDGEMENTS**

325 The authors would like to thank all participants of the study, as well as the staff at the PHCs  
326 participating in the TBD STING study. We would like to thank the members of the TBD  
327 STING study group for all the valuable work and advice on the study design and collection  
328 logistics. We would also like to thank Dr Andrew Jenkins, Telemark University College,  
329 Norway, for excellent linguistic revision. This study was supported by The Swedish  
330 Research Council (Branch of Medicine, no. K2008-58X-14631-06-3), The Medical Research  
331 Council of Southeast Sweden (FORSS, 657881), Region Östergötland, the Interreg IVA  
332 Programme ScandTick, Interreg V ÖKS programme ScandTick Innovation, the Wilhelm and  
333 Else Stockmann Foundation, and the Foundation for Åland Medical Research of the Åland  
334 Culture Foundation.

**335 REFERENCES**

827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885

- 336 Andersson, M.O., Bergvall, U.A., Chirico, J., Christensson, M., Lindgren, P.E., Nordström, J.,  
337 Kjellander, P. 2016. Molecular detection of *Babesia capreoli* and *Babesia venatorum* in wild  
338 Swedish roe deer, *Capreolus capreolus*. Parasit. Vectors. 9, 221.  
339 <https://doi.org/10.1186/s130710161503-8>.
- 340  
341 Blum, S., Gattringer, R., Haschke, E., Walochnik, J., Tschurtschenthaler, G., Lang, F., Oberbauer, R.  
342 2011. The case: hemolysis and acute renal failure. Babesiosis. Kidney Int. 80, 681-83.  
343 <https://doi.org/10.1038/ki.2011.184>.
- 344  
345 Bläckberg, J., Lazarevic, V.L., Hunfeld, K.P., Persson, K.E.M. 2018. Low-virulent *Babesia venatorum*  
346 infection masquerading as hemophagocytic syndrome. Ann. Hematol. 97, 731-733.  
347 <https://doi.org/10.1007/s00277-017-3220-6>.
- 348  
349 Bonnet, S., Jouglin, M., Malandrin, L., Becker, C., Agoulon, A., L'Hostis, M., Chauvin, A. 2007.  
350 Transstadial and transovarial persistence of *Babesia divergens* DNA in *Ixodes ricinus* ticks fed on  
351 infected blood in a new skin-feeding technique. Parasitology. 134, 197-207.  
352 <https://doi.org/10.1017/S0031182006001545>
- 353  
354 Casati, S., Sager, H., Gern, L., Piffaretti, J.C. 2006. Presence of potentially pathogenic *Babesia* sp. for  
355 human in *Ixodes ricinus* in Switzerland. Ann Agric Environ Med. 13, 65-70.  
356
- 357 Diuk-Wasser, M.A., Vannier, E., Krause, P.J. 2016. Coinfection by *Ixodes* tick-borne pathogens:  
358 ecological, epidemiological, and clinical consequences. Trends Parasitol. 32, 30-42.  
359 <https://doi.org/10.1016/j.pt.2015.09.008>
- 360  
361 Dunn, J.M., Krause, P.J., Davis, S., Vannier, E.G., Fitzpatrick, M.C., Rollend, L., Belperron, A.A., States,  
362 S.L., Stacey, A., Bockenstedt, L.K., Fish, D., Diuk-Wasser, M.A. 2014. *Borrelia burgdorferi* promotes  
363 the establishment of *Babesia microti* in the northeastern United States. PLoS One. 9, e115494.  
364 <https://doi.org/10.1371/journal.pone.0115494>
- 365  
366 Fryland, L., Wilhelmsson, P., Lindgren, P.E., Nyman, D., Ekerfelt, C., Forsberg, P. 2011. Low risk of  
367 developing *Borrelia burgdorferi* infection in the south-east of Sweden after being bitten by a *Borrelia*  
368 *burgdorferi*-infected tick. Int J Infect Dis. 15, e174-81. <https://doi.org/10.1016/j.ijid.2010.10.006>
- 369  
370 Grankvist, A., Sandelin, L.L., Andersson, J., Fryland, L., Wilhelmsson, P., Lindgren, P.E., Forsberg, P.,  
371 Wenneras, C. 2015. Infections with *Candidatus Neohhrlichia mikurensis* and cytokine responses in 2  
372 persons bitten by ticks, Sweden. Emerg Infect Dis. 21, 1462-65.  
373 <https://doi.org/10.3201/eid2108.150060>
- 374  
375 Gray, E.B., Herwaldt, B.L. 2019. Babesiosis Surveillance - United States, 2011-2015. MMWR Surveill  
376 Summ. 68, 1-11.  
377 <https://doi.org/10.15585/mmwr.ss6806a1>.
- 378  
379 Gray, J., Stanek, G., Kundi, M., Kocianova, E. 2005. Dimensions of engorging *Ixodes ricinus* as a  
380 measure of feeding duration. Int J Med Microbiol. 295, 567-72.

886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944

381 <https://doi.org/10.1016/j.ijmm.2005.05.008>  
382

383 Haapasalo, K., Suomalainen, P., Sukura, A., Siikamaki, H., Jokiranta, T.S. 2010. Fatal babesiosis in  
384 man, Finland, 2004. *Emerg Infect Dis.* 16, 1116-18.  
385 <https://doi.org/10.3201/eid1607.091905>  
386

387 Haselbarth, K., Tenter, A.M., Brade, V., Krieger, G., Hunfeld, K.P. 2007. First case of human babesiosis  
388 in Germany - clinical presentation and molecular characterisation of the pathogen. *Int J Med*  
389 *Microbiol.* 297, 197-204.  
390 <https://doi.org/10.1016/j.ijmm.2007.01.002>  
391

392 Hatcher, J.C., Greenberg, P.D., Antique, J., Jimenez-Lucho, V.E. 2001. Severe babesiosis in Long  
393 Island: review of 34 cases and their complications. *Clin Infect Dis.* 32, 1117-25.  
394 <https://doi.org/10.1086/319742>  
395

396 Henningsson, A.J., Wilhelmsson, P., Gyllemark, P., Kozak, M., Matussek, A., Nyman, D., Ekerfelt, C.,  
397 Lindgren, P.E., Forsberg, P. 2015. Low risk of seroconversion or clinical disease in humans after a bite  
398 by an *Anaplasma phagocytophilum*-infected tick. *Ticks Tick Borne Dis.* 6, 787-92.  
399 <https://doi.org/10.1016/j.ttbdis.2015.07.005>  
400

401 Henningsson, A.J., Lindqvist, R., Norberg, P., Lindblom, P., Roth, A., Forsberg, P., Bergström, T.,  
402 Överby, A.K., Lindgren, P.E. 2016. Human tick-borne encephalitis and characterization of virus from  
403 biting tick. *Emerg Infect Dis.* 22, 1485-87.  
404 <https://doi.org/10.3201/eid2208.151962>  
405

406 Herwaldt, B.L., Caccio, S., Gherlinzoni, F., Aspöck, H., Slemenda, S.B., Piccaluga, P., Martinelli, G.,  
407 Edelhofer, R., Hollenstein, U., Poletti, G., Pampiglione, S., Loschenberger, K., Tura, S., Pieniazek, N.J.  
408 2003. Molecular characterization of a non-*Babesia divergens* organism causing zoonotic babesiosis  
409 in Europe. *Emerg Infect Dis.* 9, 942-48.  
410 <https://doi.org/10.3201/eid0908.020748>  
411

412 Hildebrandt, A., Hunfeld, K.P., Baier, M., Krumbholz, A., Sachse, S., Lorenzen, T., Kiehntopf, M.,  
413 Fricke, H.J., Straube, E. 2007. First confirmed autochthonous case of human *Babesia microti* infection  
414 in Europe. *Eur J Clin Microbiol Infect Dis.* 26, 595-601.  
415 <https://doi.org/10.1007/s10096-007-0333-1>  
416

417 Holler, J.G., Roser, D., Nielsen, H.V., Eickhardt, S., Chen, M., Lester, A., Bang, D., Frandsen, C., David,  
418 K.P. 2013. A case of human babesiosis in Denmark. *Travel Med Infect Dis.* 11, 324-28.  
419 <https://doi.org/10.1016/j.tmaid.2013.06.003>  
420

421 Jensen, P.M., Christoffersen, C.S., Moutailler, S., Michelet, L., Klitgaard, K., Bodker, R. 2017.  
422 Transmission differentials for multiple pathogens as inferred from their prevalence in larva, nymph  
423 and adult of *Ixodes ricinus* (Acari: Ixodidae). *Exp Appl Acarol.* 71, 171-182.

- 945  
946  
947 424 <https://doi.org/10.1007/s10493-017-0110-5>  
948 425  
949  
950 426 Johnson, S.T., Cable, R.G., Tonnetti, L., Spencer, B., Rios, J., Leiby, D.A. 2009. Seroprevalence of  
951 427 *Babesia microti* in blood donors from *Babesia*-endemic areas of the northeastern United States:  
952 428 2000 through 2007. *Transfusion*. 49, 2574-82.  
953 429 <https://doi.org/10.1111/j.1537-2995.2009.02430.x>  
954 430  
955  
956 431 Karlsson, M.E., Andersson, M.O. 2015. *Babesia* species in questing *Ixodes ricinus*, Sweden. *Ticks Tick*  
957 432 *Borne Dis*. 7, 10-12.  
958 433 <https://doi.org/10.1016/j.ttbdis.2015.07.016>  
959 434  
960 435 Knapp, K.L., Rice, N.A. 2015. Human coinfection with *Borrelia burgdorferi* and *Babesia microti* in the  
961 436 United States. *J Parasitol Res*. 2015, 587131. <https://doi.org/10.1155/2015/587131>  
962 437  
963  
964 438 Krause, P.J. 2003. Babesiosis diagnosis and treatment. *Vector Borne Zoonotic Dis*. 3, 45-51.  
965 439 <https://doi.org/10.1089/153036603765627451>  
966 440  
967  
968 441 Lempereur, L., Shiels, B., Heyman, P., Moreau, E., Saegerman, C., Losson, B., Malandrin, L. 2015. A  
969 442 retrospective serological survey on human babesiosis in Belgium. *Clin Microbiol Infect*. 21, 96.e1-7.  
970 443 <https://doi.org/10.1016/j.cmi.2014.07.004>  
971 444  
972  
973 445 Lindblom, P., Wilhelmsson, P., Fryland, L., Sjowall, J., Haglund, M., Matussek, A., Haglund, M.,  
974 446 Sjowall, J., Vene, S., Nyman, D., Forsberg, P., Lindgren, P.E. 2014. Tick-borne encephalitis virus in  
975 447 ticks detached from humans and follow-up of serological and clinical response. *Ticks Tick Borne Dis*.  
976 448 5, 21-28. <https://doi.org/10.1016/j.ttbdis.2013.07.009>  
977 449  
978  
979 450 Malandrin, L., Jouglin, M., Sun, Y., Brisseau, N., Chauvin, A. 2010. Redescription of *Babesia capreoli*  
980 451 (Enigk and Friedhoff, 1962) from roe deer (*Capreolus capreolus*): isolation, cultivation, host  
981 452 specificity, molecular characterisation and differentiation from *Babesia divergens*. *Int J Parasitol*. 40,  
982 453 277-84. <https://doi.org/10.1016/j.ijpara.2009.08.008>  
983 454  
984 455 Martinot, M., Zadeh, M.M., Hansmann, Y., Grawey, I., Christmann, D., Aguillon, S., Jouglin, M.,  
985 456 Chauvin, A., De Briel, D. 2011. Babesiosis in immunocompetent patients, Europe. *Emerg Infect Dis*.  
986 457 17, 114-16. <https://doi.org/10.3201/eid1701.100737>  
987 458  
988  
989 459 Moniuszko-Malinowska, A., Swiecicka, I., Dunaj, J., Zajkowska, J., Czupryna, P., Zambrowski,  
990 460 G., Chmielewska-Badora, J., Żukiewicz-Sobczak, W., Swierzbinska, R., Rutkowski, K., Garkowski,  
991 461 A., Pancewicz, S. 2016. Infection with *Babesia microti* in humans with non-specific symptoms in  
992 462 North East Poland. *Infect Dis (Lond)*. 48, 537-43. <https://doi.org/10.3109/23744235.2016>  
993 463  
994 464 Mørch, K., Holmaas, G., Frolander, P.S., Kristoffersen, E.K. 2015. Severe human *Babesia divergens*  
995 465 infection in Norway. *Int J Infect Dis*. 33, 37-38. <https://doi.org/10.1016/j.ijid.2014.12.034>  
996 466  
997  
998 467 Øines, Ø., Radziejvskaja, J., Paulauskas, A., Rosef, O. 2012 Prevalence and diversity of *Babesia* spp. in  
999 468 questing *Ixodes ricinus* ticks from Norway. *Parasit Vectors*. 5, 156.  
1000  
1001  
1002  
1003



2020-04-16

- 1004  
1005  
1006 469 <https://doi.org/10.1186/1756-3305-5-156>  
1007 470  
1008 471 Otranto, D., Dantas-Torres, F., Giannelli, A., Latrofa, M.S., Cascio, A., Cazzin, S., Ravagnan, S.,  
1009 472 Montarsi, F., Zanzani, S.A., Manfredi, M.T., Capelli, G. 2014. Ticks infesting humans in Italy and  
1010 473 associated pathogens. *Parasit Vectors*. 7, 328.  
1011 474 <https://doi.org/10.1186/1756-3305-7-328>  
1012 475  
1013  
1014 476 Piesman, J., Spielman, A. Human babesiosis on Nantucket Island: prevalence of *Babesia microti* in  
1015 477 ticks. 1980. *Am J Trop Med Hyg*. 29, 742-46.  
1016 478 <https://doi.org/10.4269/ajtmh.1980.29.742>  
1017 479  
1018  
1019 480 Scholtens, R.G., Braff, E.H., Healey, G.A., Gleason, N. 1968. A case of babesiosis in man in the United  
1020 481 States. *Am J Trop Med Hyg*. 17, 810-13.  
1021 482 <https://doi.org/10.4269/ajtmh.1968.17.810>  
1022 483  
1023  
1024 484 Skrabalo, Z., Deanovic, Z. 1957. Piroplasmiasis in man; report of a case. *Doc Med Geogr Trop*. 9, 11-  
1025 485 16.  
1026 486  
1027 487 Svensson, J., Hunfeld, K.P., Persson, K.E.M. 2019. High seroprevalence of *Babesia* antibodies among  
1028 488 *Borrelia burgdorferi*-infected humans in Sweden. *Ticks Tick Borne Dis*. 10, 186-190.  
1029 489 <https://doi.org/10.1016/j.ttbdis.2018.10.007>  
1030 490  
1031  
1032 491 Uhnoo, I., Cars, O., Christensson, D., Nyström-Rosander, C. 1992. First documented case of human  
1033 492 babesiosis in Sweden. *Scand J Infect Dis*. 24, 541-47.  
1034 493 <https://doi.org/10.3109/00365549209052642>  
1035 494  
1036 495 Vannier, E., Krause, P.J. 2012. Human babesiosis. *N Engl J Med*. 366, 2397-407.  
1037 496 <https://doi.org/10.1056/NEJMra1202018>  
1038 497  
1039  
1040 498 Vannier, E.G., Diuk-Wasser, M.A., Ben Mamoun, C., Krause, P.J. 2015. Babesiosis. *Infect Dis Clin*  
1041 499 *North Am*. 29, 357-70.  
1042 500 <https://doi.org/10.1016/j.idc.2015.02.008>  
1043 501  
1044  
1045 502 Wilhelmsson, P., Fryland, L., Borjesson, S., Nordgren, J., Bergström, S., Ernerudh, J., Forsberg, P.,  
1046 503 Lindgren, P.E. 2010. Prevalence and diversity of *Borrelia* species in ticks that have bitten humans in  
1047 504 Sweden. *J Clin Microbiol*. 48, 4169-76.  
1048 505 <https://doi.org/10.1128/JCM.01061-10>  
1049 506  
1050  
1051 507 Wilhelmsson, P., Lindblom, P., Fryland, L., Ernerudh, J., Forsberg, P., Lindgren, P.E. 2013a.  
1052 508 Prevalence, diversity, and load of *Borrelia* species in ticks that have fed on humans in regions of  
1053 509 Sweden and Aland Islands, Finland with different Lyme borreliosis incidences. *PLoS One*. 8, e81433.  
1054 510 <https://doi.org/10.1371/journal.pone.0081433>  
1055 511  
1056 512 Wilhelmsson, P., Lindblom, P., Fryland, L., Nyman, D., Jaenson, T. G., Forsberg, P., Lindgren, P. E.  
1057 513 2013b. *Ixodes ricinus* ticks removed from humans in Northern Europe: seasonal pattern of

infestation, attachment sites and duration of feeding. *Parasit Vectors*. 6, 362.  
<https://doi.org/10.1186/1756-3305-6-362>

Table 1. Analyzed ticks and distribution of developmental stages.

Developmental stage of the tick	Total no. of ticks analyzed	No. of <i>Babesia</i> PCR-positive ticks (%)	No. of ticks PCR-positive for <i>B. microti</i>	No. of ticks PCR-positive for <i>B. venatorum</i>	No. of ticks PCR-positive for <i>B. capreoli</i>
Adult females	478	20 (4.2)	13	6	1
Adult males	8	1 (12.5)	0	1	0
Nymphs	1466	43 (2.9)	20	20	3
Larvae	86	1 (1.2)	0	0	1
ND*	60	0 (0)	0	0	0
<b>Total</b>	<b>2098</b>	<b>65 (3.1)</b>	<b>33</b>	<b>27</b>	<b>5</b>

\*Developmental stage could not be determined due to damaged tick

Table 2. *B. microti* IgG antibody titers for samples positive in the serological analysis (n=7).

Participant Id. code.*	Antibody titers in 1 <sup>st</sup> sample <sup>‡</sup>	Antibody titers in 2 <sup>nd</sup> sample <sup>‡</sup>	<i>Babesia</i> spp. in the tick <sup>§</sup>	Tick feeding duration (h)	Developmental stage of the tick <sup>#</sup>
Afa 89	1:128	1:128	<i>B. venatorum</i>	25	N
Vofa 15	1:256	1:256	<i>B. microti</i>	35	A
Vifa 25	1:64	1:64	<i>B. microti</i>	<24	A
†Kafa 6	1:256	1:256	Neg.	<24	N
†Kafa 52	1:64	1:64	Neg.	25	N
†Kfa 11	1:128	1:64	Neg.	ND <sup>¶</sup>	N
†Kfa 13	1:64	1:64	Neg.	<24	A

\*Participant Id. code. Letters representing primary healthcare center where tick was collected followed by serial number.

†Sample from control group

‡1<sup>st</sup> sample collected at inclusion, 2<sup>nd</sup> sample after three months

§ *Babesia* spp. found in the tick collected at inclusion

¶ND = not determined due to deformed tick, making scutal and coxal indices impossible to determine

# A = Adult female, N = nymph

529

530

**Figure 1. A.** Map, showing the four regions (Northern Sweden, South Central Sweden,

Southernmost Sweden, Åland Islands) where the 34 primary health care centers (PHCs, black

dots) are located. **B.** Map showing PHCs where ticks positive for different *Babesia* species

were collected, *Babesia microti* (red circles), *Babesia venatorum* (black crosses) and *Babesia*

*capreoli* (filled green circles). Numbers (X/Y) next to region showing number of positive

ticks in each region (X) with total number of ticks collected (Y). Maps modified from

Wilhelmsson et al. 2013b.

**Figure 2.** Flow-chart showing the study design and methods. Green for information regarding ticks, red for blood samples and blue for questionnaires.

\*Excluded because no samples were available, they had been used for previous analyses.

† Excluded since sera were not available, it had been used for previous analyses.

‡ Denoted as the negative control group.

**Appendix.** 65 ticks positive for *Babesia* in real-time PCR

Region	Id. Number	Developmental stage of the tick <sup>§</sup>	Feeding time (h)	<i>Babesia</i> spp. in the tick	<i>Borrelia</i> spp. in the tick	Estimated no. of spirochetes
Åland Islands	Afa 22	A	<24	<i>B. microti</i>	<i>B. afzelii</i>	3.3 x 10 <sup>2</sup>
	Afa 89	N	25	<i>B. venatorum</i>	<i>B. afzelii</i>	3.1 x 10 <sup>3</sup>
	Afa 115B*	N	<24	<i>B. venatorum</i>		
	Afa 132B	N	46	<i>B. venatorum</i>	ND <sup>‡</sup>	1
	Afa 132C	N	<24	<i>B. microti</i>		
	Afa 132D	A	30	<i>B. venatorum</i>		
	Afa 144	N	50	<i>B. venatorum</i>		
	Afa 156A	A(male)	-†	<i>B. venatorum</i>		
	Afa 178	N	42	<i>B. venatorum</i>		
	Afa 185	A	-	<i>B. microti</i>		
	Afa 190	N	<24	<i>B. venatorum</i>	<i>B. afzelii</i>	2.3 x 10 <sup>4</sup>
	Afa 217	A	<24	<i>B. microti</i>	<i>B. afzelii</i>	1.6 x 10 <sup>4</sup>
	Afa 237	A	34	<i>B. venatorum</i>		
	Afa 338	A	<24	<i>B. microti</i>		
	Afa 350	N	<24	<i>B. microti</i>	<i>B. afzelii</i>	1.1 x 10 <sup>4</sup>
	Afa 380	N	51	<i>B. capreoli</i>		
	Afa 381	N	30	<i>B. microti</i>		
	Afa 412	N	45	<i>B. venatorum</i>		
	Afa 465	N	<24	<i>B. venatorum</i>		
	Afa 466	N	42	<i>B. venatorum</i>	<i>B. afzelii</i>	6.7 x 10 <sup>4</sup>
Afa 476	N	<24	<i>B. microti</i>	<i>B. afzelii</i>	5.3 x 10 <sup>4</sup>	
Afa 498D	N	<24	<i>B. venatorum</i>			
Afa 499	N	<24	<i>B. venatorum</i>			
Afa 518A	N	<24	<i>B. microti</i>			
Afa 537	N	<24	<i>B. venatorum</i>			
Afa 560	N	<24	<i>B. microti</i>			
Afa 615B	N	33	<i>B. microti</i>	<i>B. valaisiana</i>	3	

1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239

<b>Southcentral Sweden</b>	Bafa 6	A	<24	<i>B. microti</i>	<i>B. afzelii</i>	3.4 x 10 <sup>2</sup>
	Bafa 73A	A	<24	<i>B. microti</i>		
	Bafa 86	A	<24	<i>B. capreoli</i>	<i>B. burgdorferi</i> sensu stricto	3.1 x 10 <sup>4</sup>
	Ekfa 84	A	-	<i>B. microti</i>	<i>B. afzelii</i>	2.7 x 10 <sup>4</sup>
	Grfa 28	N	56	<i>B. microti</i>	<i>B. garinii</i>	1.3 x 10 <sup>3</sup>
	Hafa 8	N	<24	<i>B. venatorum</i>		
	Hafa 16	N	25	<i>B. microti</i>	<i>B. afzelii</i>	3.1 x 10 <sup>4</sup>
	Hafa 44B	A	<24	<i>B. venatorum</i>	<i>B. miyamotoi</i>	1.9 x 10 <sup>6</sup>
	Hafa 115	A	61	<i>B. venatorum</i>	<i>B. afzelii</i>	1.7 x 10 <sup>1</sup>
	Jofa 7	N	47	<i>B. microti</i>	<i>B. afzelii</i>	6.3 x 10 <sup>2</sup>
	Lidfa 14	N	55	<i>B. venatorum</i>	<i>B. garinii</i>	1.4x 10 <sup>3</sup>
	Lidfa 30	A	<24	<i>B. microti</i>	ND	1.1 x 10 <sup>2</sup>
	Lidfa 39A	L	-	<i>B. capreoli</i>		
	Lidfa 39B	N	52	<i>B. capreoli</i>		
	Lidfa 46	N	26	<i>B. venatorum</i>		
	Lidfa 92B	N	<24	<i>B. microti</i>	<i>B. afzelii</i>	3.9 x 10 <sup>4</sup>
	Mekfa 16	A	-	<i>B. venatorum</i>		
	Sofa 76	A	166	<i>B. microti</i>	ND	1.1 x 10 <sup>2</sup>
	Vvfa 66	A	-	<i>B. microti</i>		
	Vifa 20	N	70	<i>B. microti</i>		
	Vifa 25	A	<24	<i>B. microti</i>	<i>B. afzelii</i>	1.9 x 10 <sup>5</sup>
<b>Southernmost Sweden</b>	Blefa 13	N	<24	<i>B. microti</i>	<i>B. afzelii</i>	7.9 x 10 <sup>3</sup>
	Blefa 19	N	37	<i>B. microti</i>		
	Blefa 38B	N	<24	<i>B. microti</i>	<i>B. afzelii</i>	8.8 x 10 <sup>3</sup>
	Blefa 38D	N	<24	<i>B. venatorum</i>	<i>B. afzelii</i>	9.5 x 10 <sup>2</sup>
	Kafa 7	N	57	<i>B. venatorum</i>		
	Kafa 34	N	59	<i>B. venatorum</i>		
	Kafa 36	N	<24	<i>B. microti</i>	<i>B. afzelii</i>	3.5 x 10 <sup>3</sup>
	Kafa 75	N	49	<i>B. microti</i>		
	Kafa 84A	N	60	<i>B. microti</i>	<i>B. afzelii</i>	7.0 x 10 <sup>1</sup>
	Kafa 100A	N	25	<i>B. microti</i>	<i>B. afzelii</i>	1.8 x 10 <sup>4</sup>
	Kfa 9	N	<24	<i>B. venatorum</i>		
	Kfa 18	N	-	<i>B. venatorum</i>		
	Kfa 25	N	37	<i>B. capreoli</i>		
	Lafa 13	A	51	<i>B. venatorum</i>		
	Ofa 15C	N	35	<i>B. microti</i>	<i>B. afzelii</i>	1.4 x 10 <sup>4</sup>
	Vofa 15	A	35	<i>B. microti</i>		
	Ysfa 9	A	38	<i>B. microti</i>	<i>B. afzelii</i>	1.5 x 10 <sup>1</sup>

\*Letter after Id. Code for participants who turned in more than one tick to the PHC, the first tick XXfaA, the second XXfaB etc.


\*Feeding time could not be determined


\*ND = Species could not be determined

§A = Adult female, N = nymph

549







## **CREDIT AUTHOR STATEMENT**

Wilhelmsson, Peter:

Conceptualization  
Formal Analysis  
Investigation  
Writing – Review & Editing  
Supervision

Lövmar, Matilda:

Investigation  
Data Curation  
Writing – Original Draft  
Writing – Review & Editing  
Visualization

Krogfelt, Karen Angeliki:

Conceptualization  
Project Administration  
Writing – Review & Editing

Nielsen, Henrik Vedel:

Investigation  
Resources  
Writing – Review & Editing

Forsberg, Pia:

Conceptualization  
Resources  
Writing – Review & Editing  
Supervision  
Project Administration  
Funding Acquisition

Lindgren, Per-Eric:

Conceptualization  
Writing – Review & Editing  
Supervision



Project Administration  
Funding Acquisition