Changes in Lyme neuroborreliosis incidence in Denmark, 1996 to 2015

Tetens, Malte M; Haahr, Rasmus; Dessau, Ram B; Krogfelt, Karen A; Bodilsen, Jacob; Andersen, Nanna S; Møller, Jens K; Roed, Casper; Christiansen, Claus B; Ellermann-Eriksen, Svend; Bangsbo, Jette M.; Hansen, Klaus; Benfield, Thomas L; Andersen, Christian Ø.; Obel, Niels; Omland, Lars H; Lebech, Anne-Mette

Published in:
Ticks and Tick-borne Diseases

DOI:
[10.1016/j.ttbdis.2020.101549](https://doi.org/10.1016/j.ttbdis.2020.101549)

Publication date:
2020

Document Version
Peer reviewed version

Citation for published version (APA):
Tetens, M. M., Haahr, R., Dessau, R. B., Krogfelt, K. A., Bodilsen, J., Andersen, N. S., Møller, J. K., Roed, C., Christiansen, C. B., Ellermann-Eriksen, S., Bangsbo, J. M., Hansen, K., Benfield, T. L., Andersen, C. Ø., Obel, N., Omland, L. H., & Lebech, A-M. (2020). Changes in Lyme neuroborreliosis incidence in Denmark, 1996 to 2015. *Ticks and Tick-borne Diseases*, 11(6), [101549]. <https://doi.org/10.1016/j.ttbdis.2020.101549>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

1 **Title:** Changes in Lyme neuroborreliosis incidence in Denmark, 1996 to 2015

2

3 **Authors:** Malte M. Tetens¹, Rasmus Haahr¹, Ram B. Dessau, M.D.², Karen A. Krogfelt, Ph.D.^{3,4},
4 Jacob Bodilsen, M.D.^{5,6}, Nanna S. Andersen, Ph.D.⁷, Jens K. Møller, D.M.Sc.⁸, Casper Roed,
5 Ph.D.¹, Claus B. Christiansen, Ph.D.⁹, Svend Ellermann-Eriksen, D.M.Sc.¹⁰, Jette M. Bangsborg,
6 D.M.Sc.¹¹, Klaus Hansen, D.M.Sc.¹², Thomas L. Benfield, D.M.Sc.^{13,14}, Christian Østergaard
7 Andersen, D.M.Sc.¹⁵, Niels Obel, D.M.Sc.^{1,14}, Lars H. Omland, D.M.Sc.¹, Anne-Mette Lebech,
8 D.M.Sc.^{1,14}

9

10 **Affiliations**

11 1. Department of Infectious Diseases, Copenhagen University Hospital, Rigshospitalet,
12 Copenhagen, Denmark

13 2. Department of Clinical Microbiology, Slagelse Hospital, Slagelse, Denmark

14 3. Department of Virus and Microbiological Special Diagnostics, Statens Serum Institut, Denmark

15 4. Department of Natural Sciences and Environment, Roskilde University, Denmark

16 5. Departments of Clinical Microbiology, Aalborg University hospital, Aalborg, Denmark

17 6. Departments of and Infectious Diseases, Aalborg University hospital, Aalborg, Denmark

18 7. Clinical Centre for Emerging and Vector-borne Infections, Odense University Hospital, Odense,
19 Denmark

20 8. Department of Clinical Microbiology, Vejle Hospital, Vejle, Denmark

21 9. Department of Clinical Microbiology, Copenhagen University Hospital, Rigshospitalet,
22 Copenhagen, Denmark

23 10. Department of Clinical Microbiology, Aarhus University Hospital, Aarhus, Denmark

24 11. Department of Clinical Microbiology, Herlev University Hospital, Copenhagen, Denmark

25 12. Department of Neurology, Copenhagen University Hospital, Rigshospitalet, Copenhagen,
26 Denmark

27 13. Department of Infectious Diseases, Hvidovre University Hospital, Copenhagen, Denmark

28 14. Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of
29 Copenhagen, Copenhagen, Denmark

30 15. Department of Clinical Microbiology, Hvidovre University Hospital, Copenhagen, Denmark

31

32

33 **Corresponding author**

34 Malte Mose Tetens, Bachelor of Medical Sciences

35 Department of Infectious Diseases

36 Copenhagen University hospital

37 Blegdamsvej 9

38 DK-2100 Copenhagen Ø

39 Denmark

40 Phone: +45 25 68 72 03

41 E-mail: malte.mose.tetens.01@regionh.dk

42

43 Word count (abstract): 237

44 Word count (manuscript): 3,071

45 Tables: 1

46 Figures: 3

47

48 **Keywords (max 6):** surveillance; epidemiology; *Borrelia burgdorferi* sensu lato.

49 **Declaration of Competing Interest:** K. Hansen has received royalties from Thermo Fisher; R.

50 Dessau participated in advisory board meeting Roche Diagnostics 2018 outside this work; all other

51 authors declare no conflicts of interests and no support from any organization for the submitted

52 work.

53 **Local Ethics Committee approval:** The study was approved by the Danish Data Protection

54 Agency and the National Board of Health (RH-2015-285, I-Suite no.: 04297). An approval from the

55 local Ethics Committee is not needed for this type of study in Denmark.

56

57 **ABSTRACT**

58 Lyme neuroborreliosis (LNB) has recently been added to the list of diseases under the European
59 Union epidemiological surveillance in order to obtain updated information on incidence. The goal
60 of this study was to identify temporal (yearly) variation, high risk geographical regions and risk
61 groups, and seasonal variation for LNB in Denmark.

62 This cohort-study investigated Danish patients (n= 2,791) diagnosed with LNB
63 (defined as a positive *Borrelia burgdorferi* sensu lato (s.l.) intrathecal antibody test) between 1996-
64 2015. We calculated incidence and incidence ratios of LNB by comparing 4-yr groups of calendar-
65 years, area of residency, sex and age, income and education groups, and the number of new LNB
66 cases per month.

67 The incidence of LNB was 2.2 per 100,000 individuals and year in 1996-1999, 2.7 in
68 2004-2007 and 1.1 per 100,000 individuals in 2012-2015. Yearly variations in LNB incidence were
69 similar for most calendar-year groups. LNB incidence was highest in Eastern Denmark and among
70 males and individuals who were 0-14 yrs old, who had a yearly income of >449,000 DKK, and who
71 had a Master's degree or higher education. The number of LNB cases was highest from July to
72 November ($p < 0.001$).

73 In conclusion, based on Danish nationwide data of patients with positive *B.*
74 *burgdorferi* s.l. intrathecal antibody index (1996-2015) the incidence of LNB was found to increase
75 until 2004-2007 but thereafter to decline. European surveillance studies of Lyme borreliosis should
76 be encouraged to monitor the incidence trend.

77

1. INTRODUCTION

Lyme borreliosis, a tick-borne disease caused by the spirochetes of the *Borrelia burgdorferi* sensu lato (s.l.) complex is the most prevalent vector-borne infection in Europe (Stanek et al., 2012). Due to climatic and environmental changes, the incidence of tick-borne disease is expected to increase, and it has been suggested that Lyme borreliosis will become a more prominent health concern. However, as it is recognized that surveillance across European countries is heterogenous Lyme neuroborreliosis (LNB) has since 2018 been included on the list of diseases under the European Union epidemiological surveillance by the European Commission (The Lancet, 2018) in order to achieve more comprehensive information of the incidence of Lyme borreliosis at the European level.

In Denmark LNB has been a mandatory clinical notifiable disease since 1991. However, laboratory-based surveillance based on positive tests for *B. burgdorferi* s.l. intrathecal antibody index with electronic data-transfer has been shown to be more complete than surveillance based on manually processed notification and the hospital discharge databases registers (Dessau et al., 2015; Septfons et al., 2019). Furthermore, studies of surveillance of LNB based on positive test for *B. burgdorferi* s.l. intrathecal antibody index only reported a low risk of including misclassified cases (Dahl et al., 2019; Hansen and Lebech, 1992).

Potential changes in LNB incidence over time (years) in Denmark remains uninvestigated. Further identification of groups at risk for acquiring LNB will be of interest to the public and health care providers. We used an established Danish nationwide cohort of patients with LNB to investigate temporal changes in LNB incidence over time and to investigate whether LNB incidence differ according to geography, age, sex, or socioeconomic factors. Lastly, we investigated the seasonal variations of LNB incidence.

101 2. METHODS

102 2.1 Setting and data sources

103 In the years of study inclusion, Denmark had a population of 5.2 million to 5.7 million
104 individuals (*Statbank Denmark*, 2019). Tax-supported health care is provided free of charge to all
105 Danish residents (Schmidt et al., 2019). The unique 10-digit personal identification number
106 assigned to all Danish residents at birth or upon immigration was used to track individuals in the
107 Danish national health and administrative registries (Schmidt et al., 2019). Data on *B. burgdorferi*
108 s.l. intrathecal antibody index were extracted from data files obtained from the Danish Departments
109 of Microbiology laboratories that performed this test (see Supplementary Appendix). Additional
110 data were extracted from the Danish Civil Registration system, the Building and Housing Register,
111 the Income Statistics Register and the Danish Educational Attainment Registry (see Supplementary
112 Appendix). We extracted data on the general Danish population numbers according to calendar-
113 year, municipality of residence, sex, age, yearly income and highest educational attainment from
114 Statistics Denmark (see Supplementary Appendix).

116 2.2 Study population

117 *LNB patient cohort:* We identified all Danish residents with a positive test for *B.*
118 *burgdorferi* s.l. intrathecal antibody index during the period between 1 January 1996 and 31
119 December 2015, based on data files obtained from all Danish Departments of Microbiology that
120 performed the test during the time period. *B. burgdorferi* s.l. intrathecal antibody index was
121 measured by capture enzyme-linked immunosorbent assays (ELISA) that uses native purified
122 flagellum from *Borrelia* (strain DK1) as antigen (Hansen and Lebech, 1991). The antibody index
123 had been calculated using the formula: $(OD_{csf}/OD_{serum}) \times (OD_{csf} - OD_{serum})$, with OD_{csf} and
124 OD_{serum} representing the optical density in cerebrospinal fluid (CSF) and sera, respectively. An
125 antibody index ≥ 0.3 is regarded as positive.

126 Inclusion date for LNB patients was the date of lumbar puncture. The Danish cohort
127 of LNB patients and methodology for testing has been described previously (Haahr et al., 2019;
128 Obel et al., 2018).

130 2.3 Statistical analysis

131 *Incidence – temporal changes*

132 To investigate LNB changes over time (years) we grouped Danish resident and LNB
133 patients according to calendar-year (1996-1999, 2000-2003, 2004-2007, 2008-2011, and 2012-
134 2015). For each of these calendar year-periods, we divided the number of LNB cases with the
135 number of inhabitants at risk at 1 January to estimate the average yearly incidence and incidence
136 ratio (IR) and corresponding 95% confidence interval (CI), with the calendar year-period with the
137 lowest incidence serving as reference. We further examined whether LNB changed over time in the
138 demographic subgroups represented by geographical area of residence (East Zealand, North
139 Zealand, Southwest Zealand, Funen, South Jutland, Mid Jutland, Northwest Jutland, North Jutland
140 or Bornholm), sex (male or female), age (0-<15 years, 15-<30 years, 30-<45 years, 45-<60 years or
141 ≥ 60 years), yearly income (<150,000 DKK, 150,000-<250,000 DKK, 250,000-<450,000 DKK or $\geq$
142 450,000 DKK) and highest educational attainment (less than Bachelor's degree, Bachelor's degree
143 or higher than Bachelor's degree). The geographical areas were defined according to municipalities
144 (see Supplementary Table 1 and Supplementary Figure 1).

145

146 Incidence - demographic

147 Danish residents and LNB patients were grouped according to geographical area of
148 residence, sex, age, yearly income and highest educational attainment. We identified the total
149 number of Danish residents at risk at 1 January each year between 1996-2015 according to
150 geographical area of residency, sex, age, yearly income and educational level. We divided the total
151 number of LNB cases between 1996-2015 with the total number of inhabitants at risk at 1 January
152 every year between 1996-2015 according to geographical area of residency, sex, age, yearly income
153 and highest educational attainment to estimate average yearly incidence and IR and corresponding
154 95% CI with the category with the lowest incidence serving as reference.

155

156 Incidence – seasonal variation

157 We calculated the number of LNB patients with inclusion dates defined as the date of
158 lumbar puncture with a positive intrathecal antibody index test for each calendar-month to estimate
159 the number of new LNB cases per calendar-month. We ascertained difference in LNB incidence
160 between calendar-months and performed a chi-square test to investigate for statistical differences
161 with a significance level of $p < 0.05$. We used SPSS Statistics, version 25 (SPSS, Inc., Chicago,
162 Illinois, USA) and R version 3.5.1 for all analysis.

163

164 **2.4 Regulatory compliance**

165 The study was approved by the Danish Data Protection Agency and the National
166 Board of Health (RH-2015-285, I-Suite no.: 04297). An approval from the local Ethics Committee
167 is not needed for this type of study in Denmark.

168

3. RESULTS

We identified a total of 2,791 LNB patients with a first-time positive test for *B. burgdorferi* s.l intrathecal antibody index between 1 January 1996 and 31 December 2015. The average incidence for the entire study period was 2.6 per 100,000 individuals per year. The median age of LNB patients was 45.8 years and the proportion of males was 56.8 % (Table 1).

3.1 Incidence – temporal changes

The incidence of LNB increased nationwide from the calendar year-period 1996-1999 (2.2 LNB cases per 100,000 individuals per year) to 2004-2007 (3.3 LNB cases per 100,000 individuals per year), but thereafter declined to 1.8 LNB cases per 100,000 individuals per year during 2012-2015 (Table 1 and Figure 1). The incidence of LNB increased until 2004-2007 but thereafter declined with time for most geographical areas of residency and irrespective of sex, age, yearly income or educational level (Figure 2, Supplementary Figure 2, Supplementary Figure 3, Supplementary Figure 4 and Supplementary Figure 5).

3.2 Incidence – demographic

As shown in Table 1, higher average incidence of LNB was observed in North Zealand, Southwest Zealand, Funen and Bornholm compared with the area with the lowest incidence of LNB (South Jutland). The average incidence of LNB was higher in males (3.0 LNB cases per 100,000 individuals per year) compared with females (2.2 LNB cases per 100,000 individuals per year) corresponding to an IR of 1.3, 95% CI: 1.2 to 1.4 (Table 1). Moreover, the average incidence of LNB was higher in individuals aged 0-<15 years (4.2 LNB cases per 100,000 individuals per year, IR 5.4, 95% CI: 4.5 to 6.3), 45-<60 years (2.9 LNB cases 100,000 individuals per year, IR 3.7, 95% CI: 3.1 to 4.3) and 60 years or older (3.3 LNB cases per 100.00 individuals per year, IR 4.2, 95% CI: 3.5 to 5.0) compared to people aged 15-<30 years (0.8 LNB cases per 100,000 individuals per year) (Table 1).

With regards to annual income, the average incidence of LNB was highest in individuals with a yearly income of 450,000 DKK or more (3.3 LNB cases 100,000 individuals per year) compared with individuals with a yearly income between 150,000-<250,000 DKK (2.1 LNB cases per 100,000 individuals per year), corresponding to an IR of 1.6, 95% CI: 1.3 to 1.8 (Table 1). Finally, the average incidence of LNB was higher in individuals with a higher educational attainment than a bachelor's degree (3.3 cases per 100,000 individuals per year) compared with

201 individuals with less than a bachelor's degree (2.1 LNB cases per 100,000 individuals per year),
202 corresponding to an IR of 1.6, 95% CI: 1.3 to 1.9 (Table 1).

203

204 3.3 Incidence - seasonal variation

205 We observed a monthly variation with the lowest number of new cases of LNB in
206 March (2.8 cases of LNB/month) and the highest number of new cases in August (26.6 cases of
207 LNB/month) ($p < 0.0001$) (Figure 3).

208

209

4. DISCUSSION

Our study on LNB using nationwide data of positive *B. burgdorferi* s.l. intrathecal antibody index (1996-2015) provides an updated overview of the epidemiology of LNB in Denmark and documents that the incidence of LNB has increased until 2004-2007 but thereafter declined.

The overall incidence of LNB in Denmark between 1996-2015 was 2.6 per 100,000 individuals per year. This was of the same magnitude as has been estimated in earlier studies from Denmark as well as in Belgium but higher than in Germany and France (Enkelmann et al., 2018; Geebelen et al., 2019; Septfons et al., 2019). The incidence was however half the IR described from the neighboring country Sweden that reported an overall incidence of 6.3 per 100,000 for 2014. As the Swedish incidence data also was based on positive *B. burgdorferi* s.l. intrathecal antibody index cases with the national microbiology database our data seems comparable to their data. (Dahl et al., 2019; Knudtzen et al., 2017). The variation in incidences could be due to differences in prevalence of *B. burgdorferi* s.l. in the tick *Ixodes ricinus*, climate and biomes as well as number of people residing or working in areas endemic for Lyme borreliosis (Strnad et al., 2017). However, comparisons between countries must in general be interpreted with caution due to heterogeneity among surveillance systems which impact the estimates.

The incidence of LNB is likely to differ across European countries, possibly depending on differences in geographical factors, presence and abundance of ticks, distribution of the neurotropic genospecies *Borrelia garinii* as well as differences in human behavior influencing risk of tick exposure. Comparison of incidences of LNB between European countries is however difficult as surveillance of LNB is based on different methods of data collection: physician reporting, hospital diagnoses or laboratory surveillance. We used laboratory data of positive *B. burgdorferi* s.l. intrathecal antibody index as measure for LNB cases, as this previously has been shown to be more accurate as physician reporting or hospital discharge diagnosis (Dessau et al., 2015; Septfons et al., 2019).

Using this measure, we observed an increasing incidence of LNB in Denmark from 1996-1999 to 2004-2007, but thereafter the incidence declined until the last study period 2012-2015 except for in two geographical areas. In agreement, no increase in LNB incidence was observed in France between 2005-2016 (Septfons et al., 2019) or Sweden between 2002-2014, but there were increases in a specific Swedish region between 2000-2005 (Henningsson et al., 2010; Södermark et al., 2017). Furthermore, the incidence of Lyme borreliosis was reported to increase in eastern Germany

242 between 2002-2006 but decreased overall between 2009-2012 and did not increase between 2013-
243 2017 (Enkelmann et al., 2018; Fülöp and Poggensee, 2008; Wilking and Stark, 2014). Changes in
244 Lyme borreliosis incidence may also be influenced by improved awareness which could lead to a
245 decrease in the number of patients that develop LNB and other disseminated manifestations of
246 Lyme borreliosis. The Danish physicians may be increasingly aware of early symptoms of Lyme
247 borreliosis and therefore promptly initiate antibiotic therapy. Also, the Danish population may have
248 become increasingly informed of the importance of daily checks for ticks, prompt removal of ticks
249 after exposure to avoid infection and recognizing erythema migrans especially in highly endemic
250 areas (Jepsen et al., 2019). Therefore, our results due not necessarily reflect the overall national
251 trend of Lyme borreliosis manifestations during the study period.

252 Identification of possible high incidence areas for acquiring LNB will be of interest to
253 the public and health care professionals. We observed a significant variation in LNB incidence
254 according to geographical area. The geographical distribution of LNB was in agreement with an
255 estimated distribution of *I. ricinus* in Denmark as well as an estimate of incidence of *Borrelia*
256 seropositivity among roe deer (Skarphéðinsson et al., 2005).

257 In agreement with other studies we found an increased incidence of LNB in males
258 (Dahl et al., 2019; Enkelmann et al., 2018; Hansen and Lebech, 1992; Södermark et al., 2017).
259 However, erythema migrans was reported to be more common in females than males (Enkelmann et
260 al., 2018). Females have been observed to use protective practices against ticks more often than
261 males (Jepsen et al., 2019). Males could also be less likely to notice early signs of Lyme disease
262 compared with females. This would lead males to develop late-stage manifestations of *B.*
263 *burgdorferi* s.l. infection such as LNB more often than females.

264 We observed a U-shaped incidence distribution with incidence being highest in
265 children younger than 15 yr and individuals 60 yr or older as also described by others (Dahl et al.,
266 2019; Dessau et al., 2015; Septfons et al., 2019; Wilking and Stark, 2014). The observation may
267 partly be explained by more intense radicular pains in middle-age and elderly patients compared
268 with younger adults leading to hospital admission (Hansen and Lebech, 1992). It has also been
269 suggested that young adult individuals are more likely to have a subclinical infection with the
270 neurotropic genospecies *B. garinii* (Carlsson et al., 2018) with less prominent radiculitic pain and
271 rarely signs of meningism. Therefore, it is possible that the actual incidence of individuals infected
272 with *B. burgdorferi* s.l. especially in this age-group is under-recognized. Explanations for the
273 differences between age groups could also be related to differences in outdoor activity and leisure

time activities between age groups. Our estimated LNB incidence for children was slightly lower than incidence of LNB in children in the two Scandinavian countries Norway and Sweden (Henningsson et al., 2010; Øymar and Tveitnes, 2009) but comparable to another study from Sweden including 548 children with LNB from Gothenburg and surrounding municipalities (Södermark et al., 2017).

We observed a proportional increase in LNB incidence with increasing income with the highest yearly earners having the highest incidence of LNB. This is in agreement with American studies on other socioeconomic factors such as race and education (Moon et al., 2019; Springer and Johnson, 2018). An increased LNB incidence for individuals with higher attained education was observed in agreement with studies from North America (Springer and Johnson, 2018). Income and educational level would be expected to be correlated, and the high incidence of LNB in groups with either high income or high education level, could likely be explained by this correlation. Since infection is correlated with tick exposure, this variability in incidence rates among age groups and education achievements could very likely be due to differences in outdoor activities.

A seasonal variation in the monthly incidence of LNB with the incidence being highest between July and November was found and thus a close association of the seasonal activity of *I. ricinus* and the onset of LNB. This agrees with other studies on the seasonal variation of LNB in Denmark (Dessau et al., 2015; Hansen and Lebech, 1992) and other European countries (Enkelmann et al., 2018; Septfons et al., 2019) as well as studies on seasonal and climatic variation in *I. ricinus* activity (Brugger et al., 2018; Lin et al., 2019; Lindgren et al., 2000).

The major strengths of the study are the large sample size and our ability to include all Danish citizens with a proven positive *B. burgdorferi* s.l. intrathecal antibody index test over a 20-yr-period. The registry-based design was hampered by lacking access to data on cerebrospinal fluid (CSF) leucocytes counts, as the presence of CSF pleocytosis would have substantiated the diagnosis of LNB further. Thus, we may have overestimated the LNB incidence. However, as a positive *B. burgdorferi* s.l. intrathecal antibody index has a high diagnostic sensitivity for LNB we assume a low rate of misclassification of LNB cases and thus this effect is likely to be very limited (Dessau et al., 2015; Hansen, 1994; Hansen and Lebech, 1992, 1991; Henningsson et al., 2014). Factors that may have led to underestimation of LNB incidence are the lack of inclusion of (i) patients with an early LNB that have not yet have developed specific *B. burgdorferi* s.l. intrathecal antibodies and (ii) patients diagnosed and treated on clinical presentation alone without CSF investigation. The observed changes in LNB incidence over the study period could be due to changes in clinical testing

practices as only positive *B. burgdorferi* s.l. intrathecal antibody index tests were used to estimate incidence of LNB. However, because the national clinical guidelines for testing and diagnosing LNB in Denmark recommend testing all suspected cases of LNB with a *B. burgdorferi* s.l. antibody index test and these guidelines have not changed during the time period covered by this study, it is unlikely that the changes in LNB incidence are due to changes in how LNB is diagnosed in Denmark (Dessau et al., 2014). Our analyses of geographical variation may be limited by the fact, that we only have access to information on place of residence, which is not necessarily the place of exposure to tick bites.

314

315 **5. CONCLUSION**

Based on a Danish nationwide cohort of patients with LNB defined by positive *B. burgdorferi* s.l. intrathecal antibody index and data from the Danish National registries, LNB incidence in Denmark increased from the time period 1996-1999 to 2004-2007 but thereafter declined until the last study time period in 2011-2015. The incidence of LNB was highest for individuals with residency in Eastern Denmark, males, children and individuals with high income and high educational attainment.

322

323 **Conflict of interest**

K. Hansen has received royalties from Thermo Fisher; R. Dessau participated in advisory board meeting Roche Diagnostics 2018 outside this work; all other authors declare no conflicts of interests and no support from any organization for the submitted work.

327

328 **Funding sources**

The study was sponsored by the Danish Council for Independent Research (grant number: 6110-0173B).

331

332

333

334 References

- 335 Brugger, K., Walter, M., Chitimia-Dobler, L., Dobler, G., Rubel, F., 2018. Forecasting next
336 season's *Ixodes ricinus* nymphal density: the example of southern Germany 2018. Exp.
337 Appl. Acarol. 75, 281–288. <https://doi.org/10.1007/s10493-018-0267-6>
- 338 Carlsson, H., Ekerfelt, C., Henningsson, A.J., Brudin, L., Tjernberg, I., 2018. Subclinical Lyme
339 borreliosis is common in south-eastern Sweden and may be distinguished from Lyme
340 neuroborreliosis by sex, age and specific immune marker patterns. Ticks Tick Borne Dis. 9,
341 742–748. <https://doi.org/10.1016/j.ttbdis.2018.02.011>
- 342 Dahl, V., Wisell, K.T., Giske, C.G., Tegnell, A., Wallensten, A., 2019. Lyme neuroborreliosis
343 epidemiology in Sweden 2010 to 2014: clinical microbiology laboratories are a better data
344 source than the hospital discharge diagnosis register. Euro Surveill. 24.
345 <https://doi.org/10.2807/1560-7917.ES.2019.24.20.1800453>
- 346 Dessau, R.B., Bangsberg, J., Hansen, K., Lebech, A., Sellebjerg, F., Skarphedinsson, S.,
347 Østergaard, C., 2014. Lyme Borreliose: Klinik, diagnostik og behandling i Danmark. The
348 Danish Microbiological Society, The Danish Society of Infectious Diseases & the Danish
349 Neurological Society.
- 350 Dessau, R.B., Espenhain, L., Mølbak, K., Krause, T.G., Voldstedlund, M., 2015. Improving
351 national surveillance of Lyme neuroborreliosis in Denmark through electronic reporting of
352 specific antibody index testing from 2010 to 2012. Euro Surveill. 20.
353 <https://doi.org/10.2807/1560-7917.es2015.20.28.21184>
- 354 Enkelmann, J., Böhmer, M., Fingerle, V., Siffczyk, C., Werber, D., Littmann, M., Merbecks, S.-S.,
355 Helmeke, C., Schroeder, S., Hell, S., Schlotthauer, U., Burckhardt, F., Stark, K., Schielke,
356 A., Wilking, H., 2018. Incidence of notified Lyme borreliosis in Germany, 2013-2017. Sci
357 Rep 8, 14976. <https://doi.org/10.1038/s41598-018-33136-0>
- 358 Fülöp, B., Poggensee, G., 2008. Epidemiological situation of Lyme borreliosis in Germany:
359 surveillance data from six Eastern German States, 2002 to 2006. Parasitol. Res. 103 Suppl 1,
360 S117-120. <https://doi.org/10.1007/s00436-008-1060-y>
- 361 Geebelen, L., Van Cauteren, D., Devleeschauwer, B., Moreels, S., Tersago, K., Van Oyen, H.,
362 Speybroeck, N., Lernout, T., 2019. Combining primary care surveillance and a meta-
363 analysis to estimate the incidence of the clinical manifestations of Lyme borreliosis in
364 Belgium, 2015-2017. Ticks Tick Borne Dis. 10, 598–605.
365 <https://doi.org/10.1016/j.ttbdis.2018.12.007>
- 366 Haahr, R., Tetens, M.M., Dessau, R.B., Krogfelt, K.A., Bodilsen, J., Andersen, N.S., Møller, J.K.,
367 Roed, C., Christiansen, C.B., Ellermann-Eriksen, S., Bangsberg, J.M., Hansen, K., Benfield,
368 T.L., Østergaard Andersen, C., Obel, N., Lebech, A.-M., Omland, L.H., 2019. Risk of
369 neurological disorders in patients with European Lyme neuroborreliosis. A nationwide
370 population-based cohort study. Clin. Infect. Dis. <https://doi.org/10.1093/cid/ciz997>
- 371 Hansen, K., 1994. Lyme neuroborreliosis: improvements of the laboratory diagnosis and a survey of
372 epidemiological and clinical features in Denmark 1985-1990. Acta Neurol. Scand., Suppl.
373 151, 1–44.
- 374 Hansen, K., Lebech, A.M., 1992. The clinical and epidemiological profile of Lyme neuroborreliosis
375 in Denmark 1985-1990. A prospective study of 187 patients with *Borrelia burgdorferi*
376 specific intrathecal antibody production. Brain 115 (Pt 2), 399–423.
377 <https://doi.org/10.1093/brain/115.2.399>
- 378 Hansen, K., Lebech, A.M., 1991. Lyme neuroborreliosis: a new sensitive diagnostic assay for
379 intrathecal synthesis of *Borrelia burgdorferi*-specific immunoglobulin G, A, and M. Ann.
380 Neurol. 30, 197–205. <https://doi.org/10.1002/ana.410300212>

381 Henningsson, A.J., Christiansson, M., Tjernberg, I., Löfgren, S., Matussek, A., 2014. Laboratory
382 diagnosis of Lyme neuroborreliosis: a comparison of three CSF anti-*Borrelia* antibody
383 assays. *Eur. J. Clin. Microbiol. Infect. Dis.* 33, 797–803. [https://doi.org/10.1007/s10096-](https://doi.org/10.1007/s10096-013-2014-6)
384 013-2014-6

385 Henningsson, A.J., Malmvall, B.-E., Ernerudh, J., Matussek, A., Forsberg, P., 2010.
386 Neuroborreliosis--an epidemiological, clinical and healthcare cost study from an endemic
387 area in the south-east of Sweden. *Clin. Microbiol. Infect.* 16, 1245–1251.
388 <https://doi.org/10.1111/j.1469-0691.2009.03059.x>

389 Jepsen, M.T., Jokelainen, P., Jore, S., Boman, A., Slunge, D., Krogfelt, K.A., 2019. Protective
390 practices against tick bites in Denmark, Norway and Sweden: a questionnaire-based study.
391 *BMC Publ. Health* 19, 1344. <https://doi.org/10.1186/s12889-019-7613-4>

392 Knudtzen, F.C., Andersen, N.S., Jensen, T.G., Skarphéðinsson, S., 2017. Characteristics and
393 clinical outcome of Lyme neuroborreliosis in a high endemic area, 1995-2014: A
394 retrospective cohort study in Denmark. *Clin. Infect. Dis.* 65, 1489–1495.
395 <https://doi.org/10.1093/cid/cix568>

396 Lin, S., Shrestha, S., Prusinski, M.A., White, J.L., Lukacik, G., Smith, M., Lu, J., Backenson, B.,
397 2019. The effects of multiyear and seasonal weather factors on incidence of Lyme disease
398 and its vector in New York State. *Sci. Total Environ.* 665, 1182–1188.
399 <https://doi.org/10.1016/j.scitotenv.2019.02.123>

400 Lindgren, E., Tälleklint, L., Polfeldt, T., 2000. Impact of climatic change on the northern latitude
401 limit and population density of the disease-transmitting European tick *Ixodes ricinus*.
402 *Environ. Health Perspect.* 108, 119–123. <https://doi.org/10.1289/ehp.00108119>

403 Moon, K.A., Pollak, J., Hirsch, A.G., Aucott, J.N., Nordberg, C., Heaney, C.D., Schwartz, B.S.,
404 2019. Epidemiology of Lyme disease in Pennsylvania 2006-2014 using electronic health
405 records. *Ticks Tick Borne Dis.* 10, 241–250. <https://doi.org/10.1016/j.ttbdis.2018.10.010>

406 Obel, N., Dessau, R.B., Krogfelt, K.A., Bodilsen, J., Andersen, N.S., Møller, J.K., Roed, C.,
407 Omland, L.H., Christiansen, C.B., Ellermann-Eriksen, S., Bangsbo, J.M., Hansen, K.,
408 Benfield, T.L., Rothman, K.J., Sørensen, H.T., Andersen, C.Ø., Lebech, A.-M., 2018. Long
409 term survival, health, social functioning, and education in patients with European Lyme
410 neuroborreliosis: nationwide population based cohort study. *BMJ* 361, k1998.
411 <https://doi.org/10.1136/bmj.k1998>

412 Øymar, K., Tveitnes, D., 2009. Clinical characteristics of childhood Lyme neuroborreliosis in an
413 endemic area of northern Europe. *Scand. J. Infect. Dis.* 41, 88–94.
414 <https://doi.org/10.1080/00365540802593453>

415 Schmidt, M., Schmidt, S.A.J., Adelborg, K., Sundbøll, J., Laugesen, K., Ehrenstein, V., Sørensen,
416 H.T., 2019. The Danish health care system and epidemiological research: from health care
417 contacts to database records. *Clin. Epidemiol.* 11, 563–591.
418 <https://doi.org/10.2147/CLEP.S179083>

419 Septfons, A., Goronflot, T., Jaulhac, B., Roussel, V., De Martino, S., Guerreiro, S., Launay, T.,
420 Fournier, L., De Valk, H., Figoni, J., Blanchon, T., Couturier, E., 2019. Epidemiology of
421 Lyme borreliosis through two surveillance systems: the national Sentinelles GP network and
422 the national hospital discharge database, France, 2005 to 2016. *Euro Surveill.* 24.
423 <https://doi.org/10.2807/1560-7917.ES.2019.24.11.1800134>

424 Skarphéðinsson, S., Jensen, P.M., Kristiansen, K., 2005. Survey of tickborne infections in
425 Denmark. *Emerg. Infect. Dis.* 11, 1055–1061. <https://doi.org/10.3201/eid1107.041265>

426 Södermark, L., Sigurdsson, V., Näs, W., Wall, P., Trollfors, B., 2017. Neuroborreliosis in Swedish
427 children: A population-based study on incidence and clinical characteristics. *Pediatr. Infect.*
428 *Dis. J.* 36, 1052–1056. <https://doi.org/10.1097/INF.0000000000001653>

429 Springer, Y.P., Johnson, P.T.J., 2018. Large-scale health disparities associated with Lyme disease
 430 and human monocytic ehrlichiosis in the United States, 2007-2013. PLoS ONE 13,
 431 e0204609. <https://doi.org/10.1371/journal.pone.0204609>
 432 Stanek, G., Wormser, G.P., Gray, J., Strle, F., 2012. Lyme borreliosis. Lancet 379, 461–473.
 433 [https://doi.org/10.1016/S0140-6736\(11\)60103-7](https://doi.org/10.1016/S0140-6736(11)60103-7)
 434 Statbank Denmark, <https://www.statistikbanken.dk/statbank5a/default.asp?w=1920> (Accessed 11
 435 October 2019).
 436 Strnad, M., Hönig, V., Růžek, D., Grubhoffer, L., Rego, R.O.M., 2017. Europe-wide meta-analysis
 437 of *Borrelia burgdorferi* sensu lato prevalence in questing *Ixodes ricinus* ticks. Appl.
 438 Environ. Microbiol. 83. <https://doi.org/10.1128/AEM.00609-17>
 439 The Lancet, 2018. Introducing EU-wide surveillance of Lyme neuroborreliosis [editorial]. Lancet
 440 392, 452. [https://doi.org/10.1016/S0140-6736\(18\)31738-0](https://doi.org/10.1016/S0140-6736(18)31738-0)
 441 Wilking, H., Stark, K., 2014. Trends in surveillance data of human Lyme borreliosis from six
 442 federal states in eastern Germany, 2009-2012. Ticks Tick Borne Dis. 5, 219–224.
 443 <https://doi.org/10.1016/j.ttbdis.2013.10.010>
 444
 445
 446
 447

448 **TABLE 1:** Incidence and incidence ratio of Lyme neuroborreliosis (LNB) stratified on 4-yr time
449 periods, geography, sex, age, yearly income and highest attained education in Denmark, 1996-2015.

	Number of LNB cases	Incidence of LNB per 100,000 individuals per year	Incidence rate ratio (95% confidence interval)
Calendar year-period			
<i>1996-1999</i>	480	2.2	1.3 (1.1 to 1.4)
<i>2000-2003</i>	579	2.7	1.5 (1.3 to 1.7)
<i>2004-2007</i>	714	3.3	1.8 (1.6 to 2.1)
<i>2008-2011</i>	616	2.8	1.6 (1.4 to 1.8)
<i>2012-2015</i>	402	1.8	1 (Ref.)
Geographical area			
<i>East Zealand</i>	683	2.3	3.9 (2.7 to 5.7)
<i>North Zealand</i>	266	3.9	6.5 (4.4 to 9.7)
<i>Southwest Zealand</i>	420	3.6	6.2 (4.2 to 9.1)
<i>Funen</i>	409	4.2	7.1 (4.9 to 10.5)
<i>South Jutland</i>	28	0.6	1 (Ref.)
<i>Mid Jutland</i>	463	2.8	4.7 (3.2 to 6.9)
<i>Northwest Jutland</i>	26	0.6	1.1 (0.6 to 1.7)
<i>North Jutland</i>	425	1.8	3.0 (2.1 to 4.5)
<i>Bornholm</i>	71	8.2	14.0 (9.0 to 21.7)
Sex			
<i>Men</i>	1,584	3.0	1.3 (1.2 to 1.4)
<i>Women</i>	1,207	2.2	1.0 (Ref.)

450

451

452 **TABLE 1:** Continued

Age (years)			
<i>0-<15</i>	826	4.2	5.4 (4.5 to 6.3)
<i>15-<30</i>	157	0.8	1.0 (Ref.)
<i>30-<45</i>	400	1.7	2.2 (1.8 to 2.6)
<i>45-<60</i>	636	2.9	3.7 (3.1 to 4.3)
<i>≥60</i>	772	3.3	4.2 (3.5 to 5.0)
Yearly income*			
<i>< 150,000 DKK</i>	474	2.2	1.0 (0.9 to 1.2)
<i>150,000-<250,000 DKK</i>	526	2.1	1.0 (Ref.)
<i>250,000-<450,000 DKK</i>	662	2.4	1.1 (1.0 to 1.3)
<i>≥450,000 DKK</i>	257	3.3	1.6 (1.3 to 1.8)
Highest attained education**			
<i>Less than bachelor's degree</i>	1,179	2.1	1.0 (Ref.)
<i>Bachelor's degree</i>	322	3.2	1.5 (1.3 to 1.7)
<i>Higher than bachelor's degree</i>	148	3.3	1.6 (1.3 to 1.9)

453 *Only individuals older than 20 years

454 **Only individuals older than 20 years and younger than 70 years

455

456 **FIGURE 1:** Nationwide yearly Lyme neuroborreliosis (LNB) incidence per 100,000 individuals in
457 Denmark

458
459 **FIGURE 2:** Yearly Lyme neuroborreliosis (LNB) incidence per 100,000 individuals by
460 geographical area

461
462 **FIGURE 3:** Average number of new cases of Lyme neuroborreliosis (LNB) nationwide per month

463
464