

CDX2 regulates interleukin-33 gene expression in intestinal epithelial cells (LS174T)

Larsen, Sylvester; Seidelin, Jakob Benedict; Davidsen, Johanne; Dahlgaard, Katja; Nielsen, Claus Henrik; Bennett, Eric Paul; Pedersen, Ole Birger; Coskun, Mehmet; Troelsen, Jesper Thorvald

Published in:
FEBS Open Bio

DOI:
[10.1002/2211-5463.13161](https://doi.org/10.1002/2211-5463.13161)

Publication date:
2021

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Larsen, S., Seidelin, J. B., Davidsen, J., Dahlgaard, K., Nielsen, C. H., Bennett, E. P., Pedersen, O. B., Coskun, M., & Troelsen, J. T. (2021). CDX2 regulates interleukin-33 gene expression in intestinal epithelial cells (LS174T). *FEBS Open Bio*, 11(6), 1638-1644. <https://doi.org/10.1002/2211-5463.13161>

General rights



Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

CDX2 regulates interleukin-33 gene expression in intestinal epithelial cells (LS174T)

Sylvester Larsen^{1,2} , Jakob Benedict Seidelin³, Johanne Davidsen^{1,4}, Katja Dahlgaard¹, Claus Henrik Nielsen⁵, Eric Paul Bennett⁶, Ole Birger Pedersen², Mehmet Coskun^{3,7} and Jesper Thorvald Troelsen¹ 

¹ Department of Science and Environment, Roskilde University, Denmark

² Department of Clinical Immunology, Næstved Hospital, Denmark

³ Department of Gastroenterology, Medical Section, Herlev Hospital, University of Copenhagen, Herlev, Denmark

⁴ Department of Surgical Gastroenterology, Zealand University Hospital, Køge, Denmark

⁵ Institute for Inflammation Research, Center for Rheumatology and Spine Diseases, Section 7521, Copenhagen University Hospital Rigshospitalet, Denmark

⁶ Department of Odontology, Copenhagen Center for Glycomics, Faculty of Health Sciences, University of Copenhagen, Denmark

⁷ Department of Biology, The Bioinformatics Centre, Biotech Research and Innovation Centre (BRIC), University of Copenhagen, Denmark

Keywords

Caudal-related homeobox transcription factor 2; gene regulation; inflammatory bowel disease; interleukin-33; intestinal regulation; transcriptional control

Correspondence

J. T. Troelsen, Department of Science and Environment, Roskilde University, Universitetsvej 1, Roskilde 4000, Denmark
 Tel: +45 4674 2728
 E-mail: troelsen@ruc.dk Mehmet Coskun and Jesper Thorvald Troelsen shared last authors

(Received 14 January 2021, revised 18 March 2021, accepted 8 April 2021)

doi:10.1002/2211-5463.13161

Dysregulation of interleukin-33 (IL-33) has been implicated in the pathogenesis of several autoimmune and inflammatory diseases, but few studies have examined transcriptional regulation of the *IL33* gene. In the intestines, gene regulation is controlled by a transcription factor network of which the intestinal-specific transcription factor CDX2 is a key component. In this study, we investigated whether CDX2 regulates *IL33* mRNA expression. We examined *IL33* mRNA expression in primary colonic epithelial cells from healthy humans and epithelial cell lines, revealing high expression levels in primary colonic and LS174T cells. Combining genomics data (ChIP-seq, RNA-seq) and *IL33* promoter analyses in LS174T cells revealed intronic enhancer activity in the *IL33* gene that is dependent on CDX2 expression. Western blotting and qRT-PCR confirmed that *IL33* expression is upregulated in a CDX2 concentration-dependent manner, thereby providing the first evidence that CDX2 regulates the expression of *IL33*.

Interleukin (IL)-33 is a pleiotropic member of the IL-1 family of cytokines and is constitutively expressed by a diverse range of cells including epithelial, endothelial, and fibroblast-like cells [1]. It regulates metabolic homeostasis and intestinal inflammation through its ability to increase wound healing and goblet cell numbers and plays a role in inflammatory bowel disease (IBD) [1,2]. IL-33 is synthesized as a full-length 30 kDa bioactive protein, but it can be cleaved or inactivated by caspases or proteases to modify its function [3].

Full-length IL-33 can be released to the extracellular environment by necrotic or damaged cells in response to injury, infections, or inflammation, where it acts as an ‘alarmin’ that amplifies immune responses [4]. However, intracellular full-length IL-33 may also act as a nuclear factor with transcriptional activity, as it was found that it binds to the p65 subunit of NF-κB and reduces p65-induced transcriptional activation [5].

Few studies have addressed the transcriptional regulation of the *IL33* gene. It has been shown that the IL-

Abbreviations

CDX2, *Caudal*-related homeobox transcription factor 2; HA, hemagglutinin; IBD, inflammatory bowel disease; IL-33, interleukin-33.

33 promoter activity can be increased by interferon- γ and that the IL-33 protein expression can be stimulated by growth factors, transcription factors, and other proteins [6,7]. In the intestines, gene regulation is controlled by a transcription factor network of which *Caudal*-related homeobox transcription factor 2 (CDX2) is a key component. CDX2 is specifically expressed in the adult intestine, and it is essential in maintaining intestinal homeostasis to control the balance between cell proliferation and differentiation [8].

Materials and methods

Cultivation of cell lines and isolation of primary colonic epithelial cells

The human colon cancer cell lines Caco-2, HT-29, LS174T, SW480 and DLD-1 (American Type Culture Collection, Rockville, MD, USA) and CDX2-inducible LS174T cells [9] were cultured in Dulbecco's modified Eagle's medium supplemented with 10% FBS and 1% glutamine and maintained at 37 °C and 5% CO₂. Primary colonic epithelial cells were isolated from four nonpregnant healthy people over the age of 18, by pooling six biopsies from either colon transversum or descendens taken at the same time during routine colonoscopy where all investigations subsequently turned out to be normal. The method was as described previously [10]. The biopsies were collected with informed written consent at the Department of Gastroenterology, Herlev Hospital, Denmark. The study methodologies conformed to the standards set by the Declaration of Helsinki and were approved by the Capital Region of Denmark Committee on Health Research Ethics.

CDX2 knockdown and induction with doxycycline

The CDX2-inducible LS174T cells are described in Ref. [9]. In short, zinc finger nucleases were utilized to generate LS174T cells with a biallelic *CDX2* knockout. Subsequently, a cassette containing the *CDX2* gene under the control of a doxycycline-activated Tet-3G induction system was inserted into the AAVS1 safe-harbor locus of one allele. Using this system, it is possible to control the expression of *CDX2* by stimulation with doxycycline without any leakiness [9].

IL33 qRT-PCR and Luminex analysis

Total RNA was extracted (NucleoSpin RNA kit; Macherey-Nagel, Düren, Germany) from Caco-2, HT29, LS174T, CDX2-inducible LS174T, SW480, DLD1, and isolated colonic epithelial cells. 200 ng RNA was used for cDNA with SuperScript III Reverse Transcriptase

(Invitrogen, Paisley, UK). Quantitative RT-PCR was performed using the Maxima SYBR Green qPCR Master Mix (Thermo Fisher Scientific, Waltham, MA, USA). All experiments were normalized to the housekeeping gene Ribosomal Protein Large P0 (RPLP0).

CDX2 expression in wild-type and CDX2-inducible LS174T cells was induced with 0, 2, 4, 6, 8, or 10 ng·mL⁻¹ doxycycline for 24 h in quadruplicates. Growth media were stored at -80 °C before measuring IL-33 protein using the Bio-Plex Pro Human Th17 assay for IL-33 (171-BA012M).

Construction of luciferase reporter constructs

The *IL33* promoter and enhancer sequences were PCR-amplified and subsequently extended with In-Fusion primers and gel-purified (Table 1A). pGL4.10 vector (Promega, Madison, WI, USA) was digested with *Hind*III and gel-purified. The pGL4.10-*IL33*-promoter plasmid was constructed by In-Fusion cloning (Clontech, Fremont, CA, USA) using purified promoter DNA and digested vector in a 1 : 2 molar ratio. 2.5 μ L cloning reaction was used for transformation of One Shot TOP10 chemically competent *Escherichia coli* (Thermo Fisher Scientific). DNA from colonies was isolated and Sanger-sequenced (Beckman Coulter Genomics, High Wycombe, UK). For the construction of the pGL4.10-*IL33*-promoter+enhancer plasmid, the pGL4.10-*IL33*-promoter plasmid was digested with *Sal*I, gel-purified, and cloned with enhancer insert in a 1 : 2 molar ratio using the In-Fusion cloning kit.

Transfection with reporter constructs

LS174T and CDX2-inducible LS174T cells were seeded in 24-well plates at 100 000 cells/well and transfected after 24 h. For each well, a transfection mixture of 50 μ L DNA/PEI mix was prepared, consisting of 2 μ M polyethylenimine (PEI) (Alfa Aesar, Heysham, UK) diluted in 150 mM NaCl to 25 μ L and 300 ng plasmid DNA diluted with 150 mM NaCl to 25 μ L. The transfection mix contained 50 ng CMV promoter-driven expression plasmid for either the *IL33* promoter or both the promoter and enhancer. Fifty nanogram of an empty pGL4.10 CMV expression plasmid was used as a no-insert control. Twenty-five nanogram CMV-LacZ plasmid was added to the mix as an internal control of transfection efficiency, and 225 ng pBluescript SK+ II was used as inactive DNA to reach 300 ng DNA/well. Volume was adjusted with 150 mM NaCl, and the transfection mix was incubated for 60 min at room temperature. The transfection mix was added in drops to the wells, followed by light shaking. After the transfection, the plates were centrifuged at 390 g for 5 min and incubated overnight at 37 °C and 5% CO₂. The media were changed after 24 h, and cells were grown for an additional 24 h before assayed for luciferase and β -galactosidase activity.

Table 1. (A) Primers used to produce the luciferase reporter constructs. Primers for In-Fusion cloning have the tail sequences underlined, and bold text denotes the sequence of the restriction sites used for fusing the insert and vector. (B) Primers used for real-time PCR in the analysis of chromatin immune-precipitated DNA from LS174T wild-type nuclear extract and in the analysis of mRNA from intestinal cell lines and primary colonic epithelial cells.

Primer name	Primer sequence	Restriction site
(A)		
Primers used for reporter constructs		
<i>IL33</i> promoter FW	GCCCCTTTGTTTCAAGGTCAC	
<i>IL33</i> promoter RW	TCATGGAGGGGCGAGAATTTAC	
Infusion <i>IL33</i> promoter FW	CTCGGCGGCC AAGCTT	HindIII
Infusion <i>IL33</i> promoter RW	CCGGATTGCC AAGCTT	HindIII
<i>IL33</i> enhancer FW	ATGGTGGGACTGGCATAAC	
<i>IL33</i> enhancer RW	AAAGAAAGATCTGTTAAGGGCCA	
Infusion <i>IL33</i> enhancer FW	ATAAGGATCC GTCGAC	SalI
Infusion <i>IL33</i> enhancer RW	AAGGGCATCG GTCGAC	Sall
(B)		
Primers used for ChIP analysis		
<i>IL33</i> ChIP FW	CAACCAGGAAATCACAGCAG	
<i>IL33</i> ChIP RW	GAGGCAATTTAGGGATGCAA	
Primers used for qPCR		
<i>IL33</i> FW	GTGGAAGAACACAGCAAGCA	
<i>IL33</i> RW	AAGCAAAGCACTCCACAGT	
<i>RPLP0</i> FW	GCTTCCTGGAGGGTGTCC	
<i>RPLP0</i> RW	GGACTCGTTTGTACCCGTG	

Reporter construct activity

The cells were rinsed three times with 1× PBS and lysed with 130 µL TROPIC lysis solution (Thermo Fisher Scientific) added 0.5 mM DTT by incubation for 10 min on ice. Ten microlitre of lysate was transferred to a 96-well light

plate, whereafter luciferase activity and β-galactosidase activity were measured using 5-s integration time and 2-s delay using the Dual-Light™ Luciferase & β-Galactosidase Reporter Gene Assay System (Thermo Fisher Scientific) in GloMax luminometer.

Chromatin immunoprecipitation assays

LS174T wild-type cells were cultured until 5 days postconfluence in a 30 × 30 cm culture dish and were cross-linked and sonicated as described previously to generate fragments of ~0.2 to 1.2 kb [11]. Briefly, four replicate immunoprecipitations (IP) were performed overnight at 4 °C with an antibody against human CDX2 (α-CDX2 clone CDX2-88; #MU392A-UC, BioGenex, Fremont, CA, USA) or, as a negative control, an antibody against influenza hemagglutinin (HA) (rabbit polyclonal α-HA, Y-11 X, #SC-805 X; Santa Cruz Biotechnology Inc, Heidelberg, Germany). Immune complexes were recovered with 50 µL protein A/G beads (Invitrogen, Grand Island, NY, USA). Real-time PCR was carried out using DNA immunoprecipitated with CDX2 and HA or on non-IP material from LS174T wild-type cells (Input; representing 1% of the total amount used in IP) using primers designed for the peak in the *IL33* region (Table 1B).

Protein extraction and immunoblotting

Wild-type and CDX2-inducible LS174T cells were seeded in 6-well plates at 300 000 cells/well. After 48 h, the media were changed to media with varying concentrations of doxycycline. At 72 h, cells were rinsed with cold PBS and lysed in RP1 lysis buffer (Macherey-Nagel, Düren, Germany) and proteins were purified. Protein concentrations were determined by the Bradford analysis (Bio-Rad, Hercules, CA, USA). 20 µg protein was mixed 1 : 4 (v/v) with Bolt loading buffer and 1 : 10 (v/v) with Bolt sample reducing agent (Thermo Fisher Scientific). Samples were incubated at 95 °C for 10 min and loaded on a Bolt 4–12% Bis-Tris Plus SDS gel, and electrophoresis was performed in 1× Bolt MOPS running buffer (Thermo Fisher Scientific) for 90 min at 100 V. Gels were transferred by wet electro-transfer to PVDF membranes for 60 min at 25 V and max 100 mA in 1× NuPAGE transfer buffer (Thermo Fisher Scientific). Membranes were blocked with 5% dry skim milk in wash buffer (1× PBS with 0.1% Tween-20) for 1 h at room temperature, washed with wash buffer 3 × 7 min, and incubated ON at 4 °C with primary antibody diluted in dilution buffer (2.5% skim milk in wash buffer). Membranes were washed 3 × 7 min and incubated with secondary antibodies for 1 h at room temperature and washed 3 × 7 min. Bands were visualized with SuperSignal™ West Dura Extended Duration Substrate (Thermo Fisher Scientific). Primary antibodies were rabbit anti-IL-33 (1 : 1000; #ab207737; Abcam, Cambridge,




Fig. 2. (A) Expression of *IL33* mRNA in intestinal cell lines and primary human colonic epithelial cells. Total RNA from SW480, Caco-2, DLD-1, HT29, LS174T cells, and colonic epithelial cells isolated from healthy humans. *IL33* mRNA levels were measured by qRT-PCR. Data were normalized to the reference gene RPLP0 and represented as mean values with SD error bars ($n = 4$) using one-way ANOVA with multiple comparisons. (B) The relative luciferase activity of the *IL33* promoter and enhancer in LS174T wild-type and CDX2-inducible cells. + indicates that either the *IL33* promoter or enhancer sequence was inserted into the luciferase construct sequence. The final concentration of doxycycline present in the growth media is shown below. The activity was normalized to the constructs with the *IL33* promoter. Bars represent mean values with SD error bars. **** $P < 0.0001$, $n = 4$ using one-way ANOVA with multiple comparisons.

cells, and partially recovered in cells where CDX2 was induced back to the endogenous level (Fig. 1A).

We then measured the expression of *IL33* transcripts by qRT-PCR in several well-characterized intestinal epithelial cell lines and in primary colonic epithelial cells isolated from healthy subjects. To characterize the epithelial-specific expression of the primary cells, we used an isolation method that minimizes fibroblast contamination [10]. The highest expression level of *IL33* was found in primary colonic epithelial cells, closely followed by LS174T and HT29 cells, while the *IL33* expression was either undetected or very low in SW480, DLD-1, and Caco-2 cell lines (Fig. 2A).

To further investigate the regulatory effect of CDX2 on *IL33* expression, we analyzed the *IL-33* promoter activity by constructing luciferase reporter plasmids containing the *IL33* promoter with or without the potential enhancer region from intron 2 and transfected them into LS174T wild-type and the CDX2-inducible LS174T clone. In the wild-type cells, the potential enhancer region increased the *IL33* promoter activity more than 45-fold from the promoter level (Fig. 2B), which demonstrates that it is a powerful enhancer of *IL33* in LS174T cells. In the unstimulated CDX2-inducible LS174T cells expressing no CDX2, the relative enhancer only increased reporter gene expression by eightfold. This indicates that the enhancer activity, to a certain degree, is regulated by other

factors than CDX2. However, when CDX2 was induced with 4 or 6 ng·mL⁻¹ doxycycline for 24 h, the effect of the enhancer increased by 19-fold and 32-fold, respectively, demonstrating a CDX2-dependent upregulation of the *IL33* promoter activity.

Subsequently, we treated CDX2-inducible LS174T cells with increasing concentrations of doxycycline and measured the *IL33* mRNA expression using quantitative RT-PCR and *IL-33* and CDX2 protein expression using western blotting analysis (Fig. 3). Similar to the RNA-seq, we observed a relatively high *IL33* mRNA and protein expression in LS174T wild-type cells and much lower expression in CDX2 knockout cells. Furthermore, we observed a doxycycline concentration-dependent increase in *IL33* mRNA and protein expression. Interestingly, another study has found a significantly reduced *IL-33* mRNA expression (GEO: GSE24633) in the intestines of mice where CDX2 was conditionally knocked out compared with CDX2-positive control mice [12].

To determine whether *IL-33* protein was excreted from the cells during CDX2 induction, we analyzed supernatants from LS174T cells induced with 0–14 ng·mL⁻¹ doxycycline for 24 h using Luminex analysis but found no detectable level of *IL-33* protein in the media (data not shown). Thus, our data suggest that the *IL-33* protein is retained intracellularly in the intestinal epithelial cells under unstimulated conditions.




Fig. 3. (Top) Quantitative RT-PCR of *IL33* mRNA expression in LS174T wild-type and CDX2-inducible LS174T cells exposed to increasing concentrations of doxycycline, normalized to the reference gene RPLP0. CDX2 expression was induced by incubation with doxycycline for 24 h. Data are represented as mean values with SD error bars, ($n = 4$), **** $P < 0.0001$ using one-way ANOVA with multiple comparisons. (Bottom) Western blot of the CDX2 and IL-33 protein level in LS174T wild-type and CDX2-inducible LS174T cells after exposure to different concentrations of doxycycline for 24 h. GAPDH and vinculin were used as loading controls.

This study provides the first evidence that the intestinal-specific transcription factor CDX2 regulates the expression of *IL33*. Furthermore, we identified an intronic enhancer of *IL33* gene expression in LS174T cells. We suggest that CDX2 adds an intestinal-specific layer of transcriptional control to the expression of IL-33, potentially placing CDX2 as a key factor controlling the intestinal expression of an important cytokine with ‘alarmin’ function. Further validation from *in vivo* studies in mice or human organoid systems could strengthen this conclusion.

Acknowledgements

This study was funded by the Independent Research Fund Denmark (grant ref 4004-00140B), the Lundbeck Foundation, Fonden til Lægevidenskabens Fremme (the A. P. Møller Foundation), the Foundation of Aase and Ejnar Danielsen, Læge Sofus Carl Emil Friis & Hustru Olga Doris Friis’ Legat, the Region Zealand Foundation for Health and Research, and the Danish National Research Foundation (DNRF107). The funders had no role in study design, data collection,

analysis, decision to publish, or preparation of the manuscript.

Conflict of interest

The authors declare no conflict of interest.

Data accessibility

The RNA-seq and ChIP-seq data shown from Pinto *et al.* are uploaded at the NIH GEO server: <http://www.ncbi.nlm.nih.gov/geo/> and are accessible using the following accession number GSE97273.

Author contributions

SL designed the study, performed experiments, analyzed data, obtained funding, and drafted the paper. JBS obtained patient samples, ran experiments, and drafted the paper. JD performed data analysis and drafted the paper. KD performed data analysis and drafted the paper. CHN performed Luminex assays, performed data analysis and drafted the paper. EPB cocreated the inducible cell line, performed data analysis, and drafted the paper. OBP performed data analysis and drafted the paper. MC designed the study, performed experiments, analyzed data, obtained funding, and drafted the paper obtained patient samples. JTT designed the study, obtained funding, performed data analysis, and drafted the paper. All authors discussed the results and contributed to the final manuscript.

References

- Liew FY, Girard JP and Turnquist HR (2016) Interleukin-33 in health and disease. *Nat Rev Immunol* **16**, 676–689.
- Seo DH, Che X, Kwak MS, Kim S, Kim JH, Ma HW, Kim DH, Kim TI, Kim WH, Kim SW *et al.* (2017) Interleukin-33 regulates intestinal inflammation by modulating macrophages in inflammatory bowel disease. *Sci Rep* **7**, 851.
- Afonina IS, Muller C, Martin SJ and Beyaert R (2015) Proteolytic processing of interleukin-1 family cytokines: variations on a common theme. *Immunity* **42**, 991–1004.
- Martin NT and Martin MU (2016) Interleukin 33 is a guardian of barriers and a local alarmin. *Nat Immunol* **17**, 122–131.
- Ali S, Mohs A, Thomas M, Klare J, Ross R, Schmitz ML and Martin MU (2011) The dual function cytokine IL-33 interacts with the transcription factor NF-kappaB to dampen NF-kappaB-stimulated gene transcription. *J Immunol* **187**, 1609–1616.

- 6 Polumuri SK, Jayakar GG, Shirey KA, Roberts ZJ, Perkins DJ, Pitha PM and Vogel SN (2012) Transcriptional regulation of murine IL-33 by TLR and non-TLR agonists. *J Immunol* **189**, 50–60.
- 7 Tsuda H, Komine M, Tominaga SI and Ohtsuki M (2017) Identification of the promoter region of human IL-33 responsive to induction by IFN γ . *J Dermatol Sci* **85**, 137–140.
- 8 Freund JN, Duluc I, Reimund JM, Gross I and Domon-Dell C (2015) Extending the functions of the homeotic transcription factor Cdx2 in the digestive system through nontranscriptional activities. *World J Gastroenterol* **21**, 1436–1443.
- 9 Pinto R, Hansen L, Hintze J, Almeida R, Larsen S, Coskun M, Davidsen J, Mitchelmore C, David L, Troelsen JT *et al.* (2017) Precise integration of inducible transcriptional elements (PrITE) enables absolute control of gene expression. *Nucleic Acids Res* **40**, 439–452.
- 10 Seidelin JB, Horn T and Nielsen OH (2003) Simple and efficient method for isolation and cultivation of endoscopically obtained human colonocytes. *Am J Physiol Gastrointest Liver Physiol* **285**, G1122–G1128.
- 11 Boyd M, Bressendorff S, Møller J, Olsen J and Troelsen JT (2009) Mapping of HNF4 α target genes in intestinal epithelial cells. *BMC Gastroenterol* **9**, 68.
- 12 Verzi MP, Shin H, Ho LL, Liu XS and Shivdasani RA (2011) Essential and redundant functions of caudal family proteins in activating adult intestinal genes. *Mol Cell Biol* **31**, 2026–2039.
- 13 Mathelier A, Fornes O, Arenillas DJ, Chen C-Y, Denay G, Lee J, Shi W, Shyr C, Tan GE, Worsley-Hunt R *et al.* (2016) JASPAR 2016: a major expansion and update of the open-access database of transcription factor binding profiles. *Nucleic Acids Res* **44**, D110–D115.