

## A Spectroscopic Overview of Intramolecular Hydrogen Bonds of NH...O,S,N Type

Hansen, Poul Erik

*Published in:*  
Molecules*DOI:*  
[10.3390/molecules26092409](https://doi.org/10.3390/molecules26092409)*Publication date:*  
2021*Document Version*  
Publisher's PDF, also known as Version of record*Citation for published version (APA):*  
Hansen, P. E. (2021). A Spectroscopic Overview of Intramolecular Hydrogen Bonds of NH...O,S,N Type. *Molecules*, 26(9), Article 2409. <https://doi.org/10.3390/molecules26092409>**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

**Take down policy**

If you believe that this document breaches copyright please contact [rucforsk@kb.dk](mailto:rucforsk@kb.dk) providing details, and we will remove access to the work immediately and investigate your claim.

## Review

# A Spectroscopic Overview of Intramolecular Hydrogen Bonds of NH ... O,S,N Type

Poul Erik Hansen 

Department of Science and Environment, Roskilde University, Universitetsvej 1, DK-4000 Roskilde, Denmark; poulerik@ruc.dk

**Abstract:** Intramolecular NH ... O,S,N interactions in non-tautomeric systems are reviewed in a broad range of compounds covering a variety of NH donors and hydrogen bond acceptors.  $^1\text{H}$  chemical shifts of NH donors are good tools to study intramolecular hydrogen bonding. However in some cases they have to be corrected for ring current effects. Deuterium isotope effects on  $^{13}\text{C}$  and  $^{15}\text{N}$  chemical shifts and primary isotope effects are usually used to judge the strength of hydrogen bonds. Primary isotope effects are investigated in a new range of magnitudes. Isotope ratios of NH stretching frequencies,  $\nu\text{NH}/\text{ND}$ , are revisited. Hydrogen bond energies are reviewed and two-bond deuterium isotope effects on  $^{13}\text{C}$  chemical shifts are investigated as a possible means of estimating hydrogen bond energies.

**Keywords:** intramolecular hydrogen bonds; deuterium isotope effects on chemical shifts; isotope ratios; hydrogen bond energies



**Citation:** Hansen, P.E. A Spectroscopic Overview of Intramolecular Hydrogen Bonds of NH ... O,S,N Type. *Molecules* **2021**, *26*, 2409. <https://doi.org/10.3390/molecules26092409>

Academic Editor: Mirosław Jabłoński

Received: 4 January 2021

Accepted: 18 April 2021

Published: 21 April 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.




**Copyright:** © 2021 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction


NH ... O,S,N (in the following called NH ... X for simplicity) intramolecular hydrogen bonds are very important building blocks in biomolecules, in self-organizing materials, in drugs, in switching molecules and in chemistry as such. Examples are given in this review. As the title indicates, this review is dealing with intramolecular hydrogen bonding. Recent reviews cover this subject [1,2]. Review [1] concentrates on Schiff bases made from salicylaldehyde and TRIS. Rules are set up to predict the predominant tautomer based on linear free energy relationships. Review [2] is focused on aromatic systems such as *o*-hydroxy Schiff bases and Mannich bases and mainly deals with tautomerism. However, in order to make it feasible within the limits of a review of this type, tautomeric systems as such are not dealt with. Nevertheless, NH ... X systems are of course one of the two forms of a tautomeric system involving NH and may as such provide useful information in the study of tautomeric systems. Older reviews covering hydrogen bonding and tautomeric systems can also be found [3–5]. Energetics are also treated and in that relation the question of hydrogen bond strength will be touched upon. The title may be misleadingly broad. By spectroscopic techniques NMR and infrared spectroscopy are primarily meant, as they are central in these studies. Within NMR,  $^1\text{H}$ -,  $^{13}\text{C}$ - and  $^{15}\text{N}$ -chemical shifts, isotope effects on chemical shifts, one-bond NH and long-range coupling constants are included, whereas for IR spectroscopy mainly NH stretching frequencies are explored. Theoretical calculations are included in cases when they supplement experimental results although they are not the focal point for this review. A goal is to give some guidance to which spectroscopic tool to use in a given situation. NH ... X hydrogen bonds have been investigated less than OH...X intramolecular hydrogen bonds. For an overview of the latter, see [6].

Intramolecular hydrogen bonds can be quite different, as seen in Figures 1 and 2. An important feature is the linker between the NH donor and the hydrogen bond acceptor. If this is a double bond or part of an aromatic system, the system has been termed resonance-assisted hydrogen bonding (RAHB) [6] and this clearly influences the type and the strength of the hydrogen bond. In other cases, e.g., proteins, intramolecular hydrogen bonds are not

very different from intermolecular ones, except for the fact that the protein may be keeping the donor and the acceptor close to each other. This type of hydrogen bond is clearly very important in proteins, both in defining  $\alpha$ -helices,  $\beta$ -sheets and turns. In proteins many hydrogen bonds may be present. Therefore methods to identify specific pairs and to characterize the individual hydrogen bonded pairs is needed. For DNA and RNA the hydrogen bonds are very similar.



**Figure 1.** Intramolecular hydrogen bond scheme of RAHB type or charge assisted type. The bond between the  $\alpha$ - and the  $\beta$ -carbon can be a double- or an aromatic bond. (A)  $R, R', R''=H$ , alkyl or aryl;  $X=H, C, O, N$  or  $S$ . (B)  $R, R', R''=H, C, O, N$ ;  $X=H$  or alkyl or aryl. (C)  $R=H$  or  $C$ ;  $X=H$  or  $C$  or  $OR$ . (D)  $R, R'$  and  $R''=H$  or alkyl or aryl;  $X=\text{lone pair}$  or  $O^-$  (nitrogen is positively charged, as it is a nitro group). (E)  $R, R'$  and  $R''=H$  or alkyl or aryl;  $X=\text{alkyl}$  or aryl. (F)  $R, R'$  and  $R''=\text{alkyl}$  or aryl;  $X + R'=\text{benzene ring}$ . (G)  $R, R'$  and  $R''=\text{alkyl}$ . (H)  $R$  and  $R'=\text{alkyl}$ ,  $X=\text{Ph}$ .



**Figure 2.** Non-RAHB system. The left hand molecule with  $X=N(CH_3)_2$  is the well known  $DMANH^+$  proton sponge. With  $X=\text{pyrrole}$  in Figure 4 hydrogen bonding to the  $\pi$ -electron system is found, [7] whereas with  $X=N(CH_3)C=OCH_3$  hydrogen bonding to nitrogen have been tested [8]. The right hand molecule,  $N,N',N''$ -tris(p-tolyl)azacalix [3](2,6)pyridine (TAPH), shows an extremely low field NH proton chemical shift of 22.1 ppm [9,10].


In the following, a number of typical hydrogen bond donors and acceptors and pairs of these are compared. It is of course not possible to mention all compounds with intramolecular NH hydrogen bonds. General trends will be given together with typical examples.

## 2. NMR

### 2.1. HN Chemical Shifts


Primary amines may pose a problem in those cases in which rotation around the C-N bond occurs. This leads to averaged NH chemical shifts or as seen later, to averaged one-bond NH coupling constants. In some cases rotation can be stopped at low temperature, as seen e.g., in bis(6-amino-1,3-dimethyluracil-5-yl)-methane derivatives (Figure 3). The

chemical shifts of the hydrogen bonded NH protons in these compounds are in the 8–9 ppm range [11].



**Figure 3.** Bis(6-amino-1,3-dimethyluracil-5-yl)-methane derivatives. R being ethyl, pyridine or *p*-dimethylaminopyridine. Taken from [11].

In compounds with aromatic rings close to the NH proton, the NH chemical shift has to be corrected for ring-current effects (for an example of ring current effects see Figure 4) in order to use this to characterize the NH ... X hydrogen bond. In A ring a current is present, whereas it is absent in B.




**Figure 4.** (a). Demonstration of possible ring current effects. In A the ring is twisted 34° out to the double bond plane, whereas in B the twist angle is 89°. Data from [12]. (b) Plot of <sup>1</sup>H chemical shifts of CH<sub>3</sub>-5 vs. NH chemical shifts for enaminones with following substituent at nitrogen phenyl substituent, *o*-methyl, *O,O*-dimethyl and 4-isopropyl. Data from [13–15].

The plot of Figure 4b demonstrates the low frequency shift of both the NH and the CH<sub>3</sub>-5 chemical shift as the phenyl group in the ortho-substituted phenyl ring is twisted.


In  $\beta$ -enaminones (Figure 1A), a hydroxyl group at the phenyl ring next to the carbonyl group clearly competes with the NH group and the NH chemical shift drops to 11.89 ppm [12], whereas an OH group in the ortho-position of a phenyl group at the nitrogen also leads to drop [16], but in this case it is a combination of a twist of the phenyl ring and a field effect caused by the OH group. In case of the *o*-hydroxyphenyl derivative, Rodríguez et al. [17] also report the finding of the keto-form at low concentration in the  $^1\text{H}$ -NMR spectrum but not in the  $^{13}\text{C}$ -NMR spectrum. *o*-Hydroxyaromatic Schiff bases T are usually either on the OH-form or being tautomeric [18]. However, recently a large number of Schiff bases based on salicylaldehyde and TRIS have been shown to be on the enamine form in the solid state. This is true for the following salicylaldehydes with substituents as follows: 5-nitro, 5-methylcarboxylate, 4-fluor, 4-choro, 4-bromo, 4-methoxy, 4-amino and 5-phenylazo [1]. A few other examples of compounds entirely on the NH form are Schiff bases of 1,3,5-triacyl-2,4,6-trihydroxybenzene [19–21], 1,3,5-triformyl-2,4,6-trihydroxybenzene [19], of gossypol [22,23] or more recently of primarily on the NH form (2-(anilinemethylidenen)cyclohexane-1,3-dione) [24].

A classic comparison is that of hydrogen bonding involving a ketone or an ester is seen in Figure 5A,B. Another comparison can be found in Ref. [25].




**Figure 5.** Comparison of different acceptors (A,B) and demonstration of steric compression (C,D). The numbers are NH chemical shifts in ppm. A and B from [15,26]. For the corresponding N-methyl derivatives, the chemical shifts are 10.90 ppm and 8.55 ppm. (C) is from [12] and (D) from [27].

It is obvious that substitution at nitrogen plays a role. Furthermore, steric compression is also an important feature as seen by comparing the 3-methyl derivative C with the corresponding non-substituted compound D. The introduction of the nitro groups leads to a slightly more acid NH group and to a stronger hydrogen bond (A vs. C). By comparison of the following compounds Figure 6, it is also clear that ring-size plays a role.




**Figure 6.** Illustration of the importance of ring size. Taken from [28]. C shows the effect of a six-membered ring. Taken from [29]. The numbers are the NH chemical shifts in ppm.

Another comparison is made in Figure 7, in which the NH is hydrogen bonded to a carbonyl group or to an amide group. The acceptor amide as acceptor clearly leads to the weaker hydrogen bond.




**Figure 7.** Comparison of ketone and amide groups as acceptors. Data from [30]. The numbers are the NH chemical shifts in ppm.

Amides and thioamides as donors are compared in Figure 8 [31].



**Figure 8.** Comparison of amides and thioamides as donors. The numbers are the NH chemical shifts in ppm. For the amide and thioamide the methyl derivatives have chemical shifts of 9.69 and 12.20 ppm. Taken from [31]. The amine is from [24].


It is seen from Figure 6 that the thioamide is the strongest donor followed by the amine and in third place the amide. Hydrogen bonding to a S=O acceptor is demonstrated in Figure 9.



**Figure 9.** Hydrogen bonding with different motifs. The numbers are NH chemical shifts in ppm. The sulfoxide is from [32]. The indole derivative is from [33]. Other similar motifs is seen in this reference. The benzo[d]thiazol is from [28].


The low chemical shift of the benzo[d]thiazol is due to the lack of conjugation between the donor and the acceptor. In Figure 10 are comparisons again done between different acceptors. NH chemical shifts may for hydrogen bonded hydrazo compounds reach values as high as 15.8 ppm when the NH is hydrogen-bonded to a pyridine nitrogen. In the minor isomer, in which the NH is hydrogen bonded to an ethyl ester group, the chemical shift is 12.8 ppm [34]. In the other pair it is obvious that the stronger hydrogen bond is to a  $\text{CH}_3\text{C}=\text{O}$  rather than to a  $\text{PhC}=\text{O}$ . The ability of aromatic nitrogens to form hydrogen bonds is also demonstrated in (Z)-5-((phenylamino)methylene)quinoxaline-6-(5H)-one, 13.15 ppm in

DMSO- $d_6$ . This value drops to 11.15 ppm in (Z)-4-((phenylamino)methylene)thiadiazol-5-(4H)-one, which has a sulphur instead of a  $\text{CH}_2=\text{CH}_2$  unit and hydrogen bonding nitrogen is now part of a five-membered ring [35]. This isomer is with hydrogen bonding to nitrogen is the minor form. The authors discuss hydrogen bonding in terms of quasi-aromaticity.



**Figure 10.** Comparison of rotamers. The numbers are NH chemical shifts in ppm. The chemical shifts are from [36]. The NH chemical shifts for the corresponding benzene derivative (benzene instead of naphthalene) are 14.6 ppm instead of 15.8 ppm. [34] If the ring is a 8-benzoquinoline the chemical shift is 15.36 ppm [37].

Pyrroles can also be hydrogen bond donors as seen in a series of compounds (Figure 11). The NH chemical shifts vary from 10.16 to 13.07 ppm [38]. In a similar case but with an OH group as the acceptor, and two pyrroles present, one hydrogen bonded the other not, the chemical shift drops to 9.39 ppm [39].




**Figure 11.** Bifurcated hydrogen bonds. Taken from [40]. The numbers are the NH chemical shifts in ppm.

Bifurcated intramolecular hydrogen bonds were found in azoymethylidene derivatives of 2-indanone (Figure 11) [40]. Similar kind of molecules have been used to establish a correlation between NH chemical shifts and hydrogen bond energy (see Section 3). It can be seen how the nature of the donor influences the chemical shifts slightly. In case of C the corresponding compound with only one intramolecular hydrogen bond has a chemical shift of 13.73 ppm illustrating the effect of bifurcation. The benzene ring seems to have little effects as the compound corresponding to A simply with the cyclopentanone unit also has a chemical shift of 13.20 ppm. However, by inserting a cyclohexanone ring the chemical shift drops slightly [38]. A different kind of bifurcation can be found in 5-(4-substituted phenylazo)-1-carboxymethyl-3-cyano-6-hydroxy-4-methyl-2-pyridones [41].


Not so common motifs are seen in Figure 12. The question is if the rather high chemical shifts in the N-oxide are caused by a strong hydrogen bond or an electric field effect from the  $\text{N}^+-\text{O}^-$  bond. A fact is that the deuterium isotope effects on  $\text{C}=\text{O}$  and  $\text{CH}_3$  carbon chemical are rather small [42] (for a general discussion of isotope effects see Section 2.4.1). The thio-Schiff base in Figure 12 is drawn as a neutral molecule and as a zwitterionic structure. The latter contributes quite considerably. The NH chemical shifts vary from 18.06 ppm for  $\text{R}$  and  $\text{R}' = \text{CH}_3$  to 19.26 ppm for  $\text{R} = \text{PhN}(\text{CH}_3)_2$  and  $\text{R}' = \text{CH}_3$  [43]

or 17.33 ppm for  $\text{CH}_3=\text{PhCH}_3$ ,  $\text{R}'=\text{CH}_3$  or 18.2 ppm for  $\text{R}=\text{PhOCH}_3$  and  $\text{R}'=\text{CH}_3$  [44]. Similar values were also obtained for derivatives in which  $\text{R}'$  is H [45].




**Figure 12.** The amino N-oxide demonstrates hydrogen bonding to a charged acceptor. Taken from [42] For the thio-Schiff base resonance forms are demonstrated. Taken from [43]. The number is the NH chemical shift in ppm.

In Figure 13 is seen a comparison of an aldehyde or an imine as acceptor and a donor being either an amide or a sulfamide. The imine is the best acceptor and as the amide is a better donor than the sulfamide.



**Figure 13.** Comparison of different amides as acceptors. In A  $\text{Y}=\text{O}$  the NH chemical shift is 10.97 ppm, whereas when  $\text{Y}=\text{N-butyl}$  it is 18.83 ppm. In B the NH chemical shift is 10.50 vs. 12.74 ppm. Taken from [46].

Hydrogen bonding is clearly weaker when the hydrogen bonding is to a  $\text{sp}^3$  hybridized oxygen and the hydrogen system is a five membered ring as demonstrated in the benzoxazine in Figure 14. Although the hydrogen bond is not so strong it is concentration independent [47].




**Figure 14.** Hydrogen bonding to single bonded oxygen. (a). Benzoxazine with intramolecular hydrogen bonding from [47]. (b).  $\text{N}'$ -benzylidenbenzohydrazide from [48]. (c). Protonated enamine molecular switch from [49]. (d). 1,4-dihydropyridine derivatives from [50]. The numbers are the NH chemical shifts in ppm.



In the case of *N'*-benzylidenbenzohydrazide, as seen in Figure 14b, a weak hydrogen bond to the methoxy group is formed. This is clearly stronger than when a classic amide is the hydrogen bond donor. In Figure 14c an amine is hydrogen bonded to an ester oxygen [49].

Charged species show large NH chemical shifts. The protonated DMAN's are tautomeric, but as long as they are symmetric this will not influence the NH chemical shifts. For a review of these see [51]. Very recently an amide type has been investigated. The R substituent has a small effect [8]. Recently motifs A and C have been combined [52].

It is obvious from the chemical shifts seen in Figure 15 and in Figure 2, that the charged systems have high NH chemical shifts. These systems are typically tautomeric and show strong intramolecular hydrogen bonds.



**Figure 15.** Hydrogen bonding of DMAN types (A–C). Counter ions are left out, but the numbers vary slightly with the counter ion. (C,D) are hydrogen bonding to a negative acceptor. Numbers are NH chemical shifts. (A) from [53], (B,C) from [54]. (D) from [55]. (E) from [56].

Motifs involving urea can be found in a review by Osmialowski [57]. Urea is versatile, as it can act both as an amide type donor and acceptor. The intramolecular nature of the hydrogen bonds were among other things established by measuring the temperature dependence. Temperature dependence was also used to distinguish between NH and OH hydrogen bonds [16]. However, this technique is by no means a reliable tool [58]. Oxamides and thioamides NH temperature coefficients have also been investigated to distinguish intra from inter molecular hydrogen bonding [59].

A number of non-RAHB cases are seen in Figure 14. In addition, proteins often offer many intramolecular hydrogen bonds (for use of coupling constants see Section 2.2). To use NH chemical shifts these should be corrected. This subject has been treated in dipeptides by Scheiner [60].

The results of Figures 4–14 can be summarized in the following way: thioamides seem to be better than hydrazo groups as donors. They are slightly better than aromatic amines, which again are better than aliphatic amines, amides and sulfamides in that order. The pyrroles are not so easy to fit into this scheme. Even when the hydrogen bond is part of a seven membered ring, they are clearly forming strong hydrogen bonds. As acceptors thiones are better than pyridines and other nitrogen containing rings, imines are better

than ketones, which are better than amides and esters. Sulfoxides are rather poor. Single bonded oxygens are even poorer. For charged systems the number of cases is limited, but seems to follow the neutral ones. However, the chemical shifts are much higher both in cases with the donor being positive charged or the acceptor being negatively charged as compared to the neutral cases.

## 2.2. Coupling Constants


Two types of couplings are immediately useful,  $^1J(\text{N,H})$  and for derivatives of aldehydes,  $^3J(\text{NH,CH})$ .  $^1J(\text{N,H})$ , one-bond hydrogen nitrogen couplings show often a numerical value of around 90 Hz. This coupling is of course negative. Dudek and Dudek showed a small difference between hydrogen bonded and non-hydrogen bonded cases [61].  $^1J(\text{N,H})$  couplings have also been calculated by DFT methods. A recent study optimized for secondary amines the functional and basis set as follows: B3LYP/6-311++G\*\* for structure optimization in chloroform (PCM approach) and APFD/6-311++G\*\*(mixed) for calculation of  $^1J(^{15}\text{N,H})$  coupling constants. A very good agreement with experimental values was found. The shorter the bond the larger the coupling constant [62]. A number of useful trends were found to complement the not so many experimental data. Using a simpler basis set, B3LYP-6-31G, it was found that one has to distinguish between primary and secondary amines. For the primary amine cases dissolved in a hydrogen bonding solvent like dimethylsulfoxide, a sulfoxide molecule has to be hydrogen bonded to the “free” NH in order to obtain good results [3].

The  $^3J(\text{NH,CH})$  coupling is for a non-tautomeric case close to 12 Hz [61]. The observation of a coupling of this magnitude or  $^1J(\text{N,H})$  of around 90 Hz is a clear indication that one is actually dealing with a  $\text{NH} \dots \text{X}$  hydrogen bond and not with an  $\text{OH} \dots \text{X}$  one or a tautomeric system. The access to reliable calculations of  $^1J(\text{N,H})$  enables one to calculate values for tautomeric systems, but also to estimate the influence of substituents.

## 2.3. Non-RAHB Cases. Couplings across Hydrogen Bonds


The  $\text{NH} \dots \text{X}$  bond is central both to proteins, DNA and RNA. A breakthrough was the observation of couplings across hydrogen bonds in RNA [63]. The presence of large  $J$  couplings (6–7 Hz) between the hydrogen bond (H-bond) donating and accepting  $^{15}\text{N}$  nuclei in Watson–Crick base pairs in double-stranded RNA was found [64]. For proteins, both in  $\alpha$ -helices and in  $\beta$ -sheets they are ubiquitous. These hydrogen bonds are generally not very strong. However, in the stronger cases very interesting coupling constants across hydrogen bonds between  $^{15}\text{N}$  and  $^{13}\text{C}=\text{O}$  (also referred to as  $\text{C}'$ ) have been observed [65,66] enabling pairing of hydrogen bond donors and acceptors. Correlations with bond lengths  $^3J_{\text{NC}'} = -59000 \exp(-4R_{\text{NO}}) \pm 0.09 \text{ Hz}$ , or  $R_{\text{NO}} = 2.75 - 0.25 \ln(-^3J_{\text{NC}'}) \pm 0.06 \text{ \AA}$  have been established [67]. Normally such coupling can only be observed in proteins below 10 kD. However, with perdeuteration  $^3J_{\text{NC}'}$  scalar couplings across hydrogen bonds could be observed in the uniformly  $^2\text{H}/^{13}\text{C}/^{15}\text{N}$ -enriched 30 kDa ribosome inactivating protein MAP30 [68].

A study of lysine interactions in ubiquitin with carbonyl backbone revealed that the  $\text{NH}_3^+$  groups of Lys29 and Lys33 exhibit measurable  $^3J_{\text{N}\zeta\text{C}'}$  couplings arising from hydrogen bonds with backbone carbonyl groups of Glu16 and Thr14, respectively. For an example see Figure 16.  $^3J_{\text{N}\zeta\text{C}'}$ -coupling constants could also be measured, these together with relaxation studies showed that the  $\text{NH}_3^+$  groups are involved in a transient and highly dynamic interaction [69].



**Figure 16.** Coupling from a carbonyl carbon to a side-chain lysine N via the hydrogen bond. Taken from [69] with permission from The American Chemical Society.

A plot of  $^1J(N,H)$  vs.  $d_{NH}$  for DNA and RNA demonstrated that the N1 ... N3 hydrogen bonds are stronger in dsRNA A:U than in dsDNA A:T bases pairs [70]. Both two-bond  $^1H$ - $^{31}P$  and three bond  $^{15}N$ - $^{31}P$  couplings have also been seen from a histidine to the phosphate group of DNA in a zink finger (see Figure 17) [71].



**Figure 17.** Hydrogen bond scalar coupling involving a phosphate and a histidine. Taken from [71] with permission from the American Chemical Society.

A very large N ... N coupling of 40 Hz through a hydrogen bond is seen in the compound in Figure 15E. The N...N distance is calculated as 2.54 Å. The coupling is the largest of this kind so far reported in a symmetric system [56].


Couplings across hydrogen bonds have also been calculated and summarized by Del Bene [72]. The couplings are dominated by the Fermi contribution and depends on the distance between the heavy atoms. Relationships are noted between hydrogen bond type, X–Y distances, NMR spin–spin coupling constants, and infrared proton-stretching frequencies. This also nicely reflects the experimental findings.

#### 2.4. Isotope Effects on Chemical Shifts

Three different types of deuterium isotope effects on chemical shift are useful,  $^n\Delta C(ND)$ ,  $^1\Delta N(D)$ ,  $^n\Delta H(ND)$  and in principle  $^n\Delta^{17}O(ND)$  in the study of intramolecular hydrogen bonds [73]. Isotope effects in the present review are defined as:  $^n\Delta = \delta X(H) - \delta X(D)$ .


#### 2.4.1. RAHB Cases (a “Double Bond” Connecting C $\alpha$ and C $\beta$ )

$^n\Delta C(ND)$  were early on studied in enaminones [74]. Later this study was extended [12,18,32,75,76]. An advantage of studying enaminones is that a number of these may exist both in an *E*- and a *Z*-form. The former without an intramolecular hydrogen bond, the latter with and thus giving a genuine reference compound. This kind of study clearly showed that the two-bond deuterium isotope effect on  $^{13}C$  chemical shifts,  $^2\Delta C(ND)$ , are larger in the intramolecular hydrogen bonded case (Figure 18). This was ascribed to resonance assistance. Having e.g., a substituent at the C- $\beta$  carbon can introduce steric strain. This will lead to a larger two-bond deuterium isotope effect as seen by comparing number from Figures 14 and 16 (see later) and to a stronger hydrogen bond. An interesting feature in such systems is also the observation of isotope effects at the carbon involved in the intramolecular hydrogen bond and even the carbon attached to the carbonyl group and beyond (see Figure 14) making this a tool for establishing pairs of hydrogen bonds in systems with several possibilities.




**Figure 18.** Deuterium isotope effects on  $^{13}C$  chemical shifts in ppm. Taken from Ref. [12].

Isotope effects have often been plotted vs. XH chemical shifts. [75] In the present case, two-bond deuterium isotope effects (TBDIE) are plotted vs. NH chemical shifts (Figure 19). It is clear that *E*- and *Z*-derivatives can be distinguished. The correlation covers enaminones, nitro- and sulphonyl derivatives [32].



**Figure 19.** Plot of two-bond deuterium isotope effects on  $^{13}C$  chemical shifts vs. NH chemical shifts for enamines. Open squares (*Z*)-enaminones, closed squares (*E*)-enaminones, + enamino esters, \* (*Z*)-nitroenamines, crosses (*Z*)-sulphonylenamines and diamonds (*E*)-sulphonylenamines,  $\phi$  indicates N-phenyl groups. Taken from [32] with permission from Wiley.

One-bond deuterium isotope effects,  $^1\Delta N(D)$ , depend strongly on hydrogen bonding (the geometry) and related to that RAHB (see Figure 20). [32]




**Figure 20.** One-bond deuterium isotope effects on  $^{15}\text{N}$  chemical shifts and two-bond deuterium isotope effects at carbon in ppm. From [76].

#### 2.4.2. Non-RAHB Cases

Deuterium isotope effects on  $^{15}\text{N}$  chemical shifts have been studied in the mono anion of 2,3-dipyrrol-2-ylquinoxaline and its 6,7-dinitro derivative (see Figure 15E) [56]. The effects 1.13 ppm and 0.88 ppm (signs have been changed from the original publication) are rather large and indicate a strong hydrogen bond.


Deuterium isotope effects on  $^{15}\text{N}$  and  $^1\text{H}$  chemical shifts have been used to judge whether certain salt bridges, which are observed in the solid also exist in solution. An example is protein G, B1 domain. The isotope effect demonstrated that two salt bridges found in the X-ray structure did not exist in solution [77]. This approach was also used in Barnase [78]. Both types of isotope effects have also been treated theoretically [79].

In ubiquitin the one-bond deuterium isotope effects of hydrogen bonded NH of the back-bone is correlated to the back bone angles and the angle between the acceptor oxygen and the NH bond (Figure 21) [80].




**Figure 21.** Angles and distances for an inter-chain hydrogen bond. Taken from [80] with permission from Springer.

In DNA and RNA through hydrogen bond isotope effects on chemical shifts can be seen from H-3 to C-2 between adenine and thymine respectively uracil. The isotope effects on chemical shifts are found to be sensitive to the N1-N3 distance suggesting that the isotope effect is sensitive to hydrogen bond strength (see Figure 22) [81,82].



**Figure 22.** Example of hydrogen bonding in an adenine:thymine base pair.

Very long range isotope effects due to deuteration at NH have been observed in N-substituted 3-(cycloamino)thioproionamides [83]. These effects were ascribed to electric field effects (Figure 23). The use of nuclei with a large chemical shift range like  $^{19}\text{F}$  makes this kind of effect very useful even for weaker hydrogen bonds.




**Figure 23.** Example of electric field isotope effect through space in N-substituted 3-(cycloamino)-thioproionamides.

#### 2.4.3. Primary Isotope Effects

$^{\text{P}}\Delta^1\text{H}(\text{ND})$ , and  $^{\text{P}}\Delta^1\text{H}(\text{NT})$ , in which the NH is replaced by either deuterium or tritium may also be used to gauge intramolecular hydrogen bonding. Only a few examples are available [84].

Protonated DMAN's ( $\text{DMANH}^+$ ) (Figure 2) show tautomerism. However, for symmetric compounds this will not influence the primary isotope effects [56]. For  $\text{DMANH}^+$  itself the  $^{\text{P}}\Delta^1\text{H}(\text{ND})$ , is 0.69 ppm. Similar values were found for a 4,5- $\text{CH}_2\text{OCH}_2$  derivative as well as the one in which the methyl groups are changed to ethyl groups. However, for 2,7-substituted derivatives the values are much smaller, dichloro, dibromo, bisdimethylamino, dimethoxy and ditrimethylsilyl gave values of 0.30, 0.23, 0.34, 0.31 and 0.14 ppm. [85] The much smaller values in these derivatives were ascribed to a buttressing effect leading to a shorter N...N distance [86]. These data and more together with data from Ref. [84] are plotted in Figure 24. Interesting points falling in-between at chemical shifts 14.03 and 14.84 ppm are N-phenyl derivatives, so these chemical shifts must be corrected for ring current effects. Others are also N-phenyl derivatives, but substituted in the 2-position so the rings are twisted heavily out of the double bond plane and thereby reducing the ring-current effects. The primary isotope effects were also related to IR isotope ratios [73].



**Figure 24.** Plot of primary deuterium isotope effects vs. NH chemical shifts. Data from [56,85].

As the review deals with NH hydrogen bonds the obvious isotope to use is deuterium. Secondary deuterium isotope effects over two-bonds tell in a qualitative way about the hydrogen bond strength. The larger the isotope effect, the stronger the hydrogen bond. For a relation to hydrogen bond energy see Section 3. In case of one bond deuterium isotope effects on  $^{15}\text{N}$  two different scenarios are found. In RAHB cases the isotope effect increases with increasing hydrogen bond strength, whereas for intramolecular hydrogen bond cases with no direct connection between donor and acceptor (like in DNA) the magnitude of the isotope effect decreases as the distance between heavy atoms decreases.

The primary isotope effects deuterium isotope effects show a more irregular behavior. It would be interesting in the future to compare primary isotope effects for the systems shown above with the one bond secondary isotope effects on  $^{15}\text{N}$ .

### 3. Energy

Hydrogen bond energies for intramolecular hydrogen bonds of the RAHB type are difficult to determine experimentally. Nevertheless, experimental values are necessary in order to have a gauge for theoretical calculations. Spectroscopic data can be useful in this context. The question of calculating the hydrogen bond energy for  $\text{NH} \dots \text{X}$  bonds were treated by Reuben [87]. He suggested to calculate the hydrogen bond energy by extending the equation originally suggested by Schaefer [88] (in this case based on  $\text{OH} \dots \text{O}$  intramolecular hydrogen bond). It is important to remember that Schaefer said a tentative equation and “rather involved but approximate calculations of electric field effects of the chemical shifts of the hydroxyl proton in intermolecularly hydrogen bonded phenol predict a very nearly linear relationship between the chemical shift and the energy”.

The energy was obtained from a correlation with NH chemical shifts:

$$\Delta\delta_{\text{NH}} = -1.06 + E_{\text{H}} \quad (1)$$

$\Delta\delta_{\text{NH}}$  is in the Reuben case referred to the NH chemical shift of N-methylaniline in  $\text{CDCl}_3$ . Chiara et al. [89] used these equations to obtain hydrogen bond energies of nitro-substituted enamines of the order of 29 to 34 KJ/mole. A very comprehensive overview is given by Afonin et al. [33] but mostly for weak interactions. Afonin et al. used a slightly different correlation:


$$E_{\text{HB}}(\Delta\delta) = \Delta\delta + (0.4 \pm 0.2) \text{ energy in Kcal/mol}$$

Afonin et al. [33] used in their study the NH donor in a pyrrole ring together with  $\text{OH} \dots \text{X}$  and  $\text{CH} \dots \text{X}$  intramolecular hydrogen bond. In this case the reference compound was pyrrole with a chemical shift of 9.25 ppm. Recently other theoretical approaches have been used. Tupikina et al. used  $^1\text{H}$  chemical shifts of  $\text{NH}_2$  groups using the non-hydrogen



bonded NH as reference for aniline derivatives and looking mainly at intermolecular hydrogen bonds. Unfortunately, the only RAHB system, an *o*-amino Schiff base, falls off the correlation line obtained [90].

The hydrogen bond and out scheme [91] used for intermolecular hydrogen bonds cannot really be used for NH...X intramolecular hydrogen bonds. A scheme has been set up by Jablonski et al. [92] for NH<sub>2</sub> groups as donors and later used for APO and 3-methyl APO [93]. A simpler method is to use the “in” and 90 degrees approach in which the energy of the latter is subtracted from the hydrogen bonded one. An example is hydrazone switches. A correlation is found between the hydrogen bond energy and the long range NH coupling across the hydrogen bridge [94]. A different method is to use the electron density at the bond critical point as suggested by Rozas et al. [95]. Using this method for 3-aminopropenal Vakili et al. [93] found 26.6 KJ/mol in good agreement with those of Jablonski et al. using MP2/6-31G\*\* and MP2/6-311++G\*\* [92]. The electron density at the bond critical point is used to estimate the hydrogen bond strength in a number of strategic intramolecular hydrogen bonds of enaminones. Inspired by the Reuben approach [87] energies have been related to two-bond deuterium isotope effects at carbons [96,97]. Recently, two-bond deuterium isotope effects (TBDIE) have been correlated to hydrogen bond energies in *o*-hydroxy aromatic aldehydes in which the hydrogen bond energies were calculated by the hb and out method [98]. The use of TBDIE has the advantage that no reference is needed. The hydrogen bond energies expressed as electron density at the bond critical point are plotted vs. two-bond deuterium isotope effects on <sup>13</sup>C chemical shifts in Figure 25 for a small set of enaminones. The ring critical points were calculated using the B3LYP/6-311++G(d,p) functional [99] and the AIM program [100,101]. A reasonable correlation is obtained considering that both ketones, esters and nitro groups are acceptors and compounds are both linear and cyclic and substituents at nitrogen both aliphatic (methyl and *t*-butyl) and aromatic. It is obvious that the cyclic compounds fall on a line of their own.




**Figure 25.** Plot of electron densities at the bond critical point vs. two-bond deuterium isotope effects on <sup>13</sup>C chemical shifts in ppm. Series 1 include linear compounds with ketones and nitro groups as acceptors, Series 2 include cyclic compounds both 5- and 6-membered rings, ketones and esters. Isotope effects in ppm from [76,89,102,103]. The ring critical points were calculated using the B3LYP/6-311++G(d,p) functional and the AIM program.

Considering the correlation between the TBDIE on <sup>13</sup>C and the electron density at the bond critical point (Figure 25) it is obvious that the large isotope effect is correlated to a stronger hydrogen bond. As TBDIE are also correlated to NH chemical shifts, a series of parameters may be used to predict hydrogen bond strength.



#### 4. Hydrogen Bond Strength

Hydrogen bond strength can be judged from the hydrogen bond energy (see Section 3). The work on predicting hydrogen bond strength using the  $pK_a$  slide should be mentioned as a reference method although not based on spectroscopic methods [104]. Gorobets et al. have suggested the difference between  $^1H$  chemical shifts of the chemical shifts of primary amides as a simple index of hydrogen bond strength in primary amides as demonstrated in Figure 26 [105]. The hydrogen bond strength can best be described by experimental trends like NH stretching frequencies, the lower the NH stretching frequency the stronger the hydrogen bond, for NH chemical shifts and two-bond deuterium isotope effects on  $^{13}C$  chemical shifts or one-bond deuterium isotope effects on  $^{15}N$  chemical shifts, the larger the stronger the hydrogen bond. A list of criteria also including structural parameters can be found in [106]. Martyniak et al. [107] also find that the asymmetry of the potential curve is a measure of hydrogen bond strength.



**Figure 26.** Primary amides used to estimate hydrogen bond strength. Taken from [105].


#### 5. Assignments

##### 5.1. CD Stretching Frequencies

A blue shift of  $C\delta D_2$  stretching frequencies of prolines in the Src homology 3 domain was interpreted as a correlation with a hydrogen bond  $N_{i+1}H \dots N_i$  interaction. The blue shifts were 10 and 17  $cm^{-1}$  for Pro 165 and 21 and 28  $cm^{-1}$  for Pro183 relative to two other prolines not affected. This was further supported by model studies of methyl terminated proline dipeptides similar to those of the SRC homology 3 domain. DFT calculations and NBO analysis supported this type of hydrogen bond [108].

##### 5.2. Assignments

As NH stretching vibrations can be difficult to assign, deuteration is often a good tool. As most NH stretching frequencies are found around 3000  $cm^{-1}$  the ND stretching frequencies will be around 2100  $cm^{-1}$ , which is in a region with few other resonances. However, the NH/ND ratio is not fixed. This was originally studied by Novak [109] and later supplemented by Sobczyk et al. [110] primarily using OH stretching frequencies. Novak included mostly intermolecular hydrogen bonds whereas Sobczyk et al. added intramolecular hydrogen bonds. The broad trend is a decrease of the isotope ratio as the NH stretching frequency decreases. However, as seen in Figure 27 based on data by Chiara et al. [89,103] for nitro-substituted enamines and nitro-substituted enamino esters and supplemented with data on enamino esters the picture is not so clear cut at all. This is also seen in the review by Sobczyk et al. [73] but is to some extent masked by the inclusion of a correlation line.



**Figure 27.** Plot of the  $\nu_{NH}/\nu_{ND}$  ratio vs.  $\nu_{ND}$ . Series 1  $\beta$ -amino- $\alpha$ -nitro- $\alpha,\beta$ -unsaturated ketones, hydrogen bonding to the keto group, Series 2 Hydrogen bonding to the nitro group from [89] and Series 3 enaminoesters from [89,103].

## 6. Conclusions

In conclusion it is useful to deal with the three different types of intramolecular hydrogen bonds separately. For intramolecular hydrogen bonds of RAHB type hydrogen bond energies are difficult to obtain, resulting in very few or none experimental results are available. A useful method, although not tested to a large extent, on intramolecularly hydrogen NH ... X systems, is electron densities at the ring critical point. The latter may also be correlated to deuterium isotope effects on  $^{13}\text{C}$  chemical shifts. In view of this, empirical parameters, NH chemical shifts, deuterium isotope effects on  $^{13}\text{C}$  or  $^{15}\text{N}$  chemical shifts may be used as indicators. NH chemical shifts have to be corrected for ring current effects, if the substituent at the nitrogen is an aromatic rings. Based on these parameters a large range of both donors and acceptors are investigated and rated with their ability to form intramolecular hydrogen bonds. NH chemical shifts and TBDIE have been correlated. However, NH chemical shifts have to be corrected for possible ring current effects, solvent effects and should be measured at as low concentration as possible. The TBDIE have the advantage of being measured as a difference and therefore being dependent on the just mentioned effects. Furthermore, deuterium isotope effects over hydrogen bonds may be used to identify hydrogen bonded pairs in case of multiple possibilities. The finding that  $^1J(\text{N,H})$  is rather invariant makes this a good gauge for checking for tautomerism.

Charged systems with connecting bonds between donor and acceptor of the intramolecular hydrogen bonds, but no conjugation, show very large NH chemical shifts.

For intramolecular hydrogen bonds in proteins and nucleic acids coupling through the hydrogen bond can be measured and the magnitude increases with the shorter the heavy atom is. Also  $^1J(\text{N,H})$  couplings are useful in the characterization of intramolecular hydrogen bonds.

Theoretical calculations are very useful in calculation of energies, coupling constants, isotope effects on chemical shifts and the finding that NH stretching frequencies can be calculated routinely means that they can be used more easily to identify infra red bands due to NH stretching vibrations. The use of  $\nu_{H}/\nu_{D}$  ratios to predict NH stretching frequencies based on ND stretching frequencies probably need more investigations.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** The author would warmly like to thank R. Rutu and B.A. Saeed, Basra University, Iraq for their help in calculating electron densities at the bond critical points.

**Conflicts of Interest:** The author declares no conflict of interest.

## References

1. Martínéz, R.F.; Matamoros, E.; Cintas, P.; Palacios, J.C. Imine or Enamine? Insights and Predictive Guidelines from the electronic Effect of Substituents in H-Bonded Salicylimines. *J. Org. Chem.* **2020**, *85*, 5838–5862. [\[CrossRef\]](#)
2. Jezierska, A.; Tolstoy, P.M.; Panek, J.J.; Filarowski, A. Intramolecular Hydrogen Bonds in Selected Aromatic Compounds: Recent Developments. *Catalysts* **2019**, *9*, 909. [\[CrossRef\]](#)
3. Hansen, P.E. Methods to distinguish tautomeric cases from static ones. In *Tautomerism: Ideas, Compounds, Applications*; Antonov, L., Ed.; Wiley-VCH: Weinheim, Germany, 2016.
4. Hansen, P.E.; Spanget-Larsen, J. NMR and IR investigations of strong intramolecular hydrogen bonds. *Molecules* **2017**, *22*, 552. [\[CrossRef\]](#)
5. Sobczyk, L.; Chudoba, D.; Tolstoy, P.M.; Filarowski, A. Some Brief Notes on Theoretical and Experimental Investigations of Intramolecular Hydrogen Bonding. *Molecules* **2016**, *21*, 1657. [\[CrossRef\]](#) [\[PubMed\]](#)
6. Gilli, P.; Bertolasi, V.; Ferretti, V.; Gilli, G. Evidence for intramolecular N-H . . . O Resonance-Assisted Hydrogen Bonding in  $\beta$ -Enaminones and Related Heterodienes. A Combined Crystal-Structural, IR and NMR Spectroscopic, and Quantum-Mechanical Investigation. *J. Am. Chem. Soc.* **2000**, *122*, 10405–10417. [\[CrossRef\]](#)
7. Pozharskii, A.F.; Ozeryanskii, V.A.; Filatova, E.A.; Dyablo, O.V.; Pogosa, O.G.; Borodkin, G.S.; Filarowski, A.; Steglenko, D.V. Neutral Pyrrole nitrogen Atom as a  $\pi$ - and Mixed N, $\pi$ -Donor in Hydrogen Bonding. *J. Org. Chem.* **2019**, *84*, 726–737. [\[CrossRef\]](#)
8. Mikshiev, V.Y.; Pozharskii, A.F.; Filarowski, A.; Novikov, A.S.; Antonov, A.S.; Tolstoy, P.M.; Vovk, M.A.; Khoroshilova, O.V. How Strong is Hydrogen Bonding to Amide Nitrogen? *ChemPhysChem* **2020**, *21*, 1–9. [\[CrossRef\]](#) [\[PubMed\]](#)
9. Zhou, S.; Wang, L. Quantum effects and  $^1\text{H}$  NMR chemical shifts of a bifurcated short hydrogen bond. *J. Chem. Phys.* **2020**, *153*, 114301. [\[CrossRef\]](#)
10. Kanbara, T.; Suzuki, Y.; Yamamoto, T. New proton-sponge-like macrocyclic compound: Synergistic hydrogen bonds of aminopyridine. *Eur. J. Org. Chem.* **2006**, 3314–3316. [\[CrossRef\]](#)
11. Sigalov, M.V.; Pylaeva, S.A.; Tolstoy, P.M. Hydrogen Bonding in Bis(6-amino-1,3-dimethyluracil-5-yl)-methane Derivatives: Dynamic NMR and DFT Evaluation. *J. Phys. Chem. A* **2016**, *120*, 2737–2748. [\[CrossRef\]](#)
12. Zhiglova, D.K.; Genov, D.G.; Bolvig, S.; Hansen, P.E. Deuterium Isotope Effects on  $^{13}\text{C}$  Chemical Shifts of Enaminones. *Acta Chem. Scand.* **1997**, *51*, 1016–1023. [\[CrossRef\]](#)
13. Chen, X.; She, J.; Shang, Z.; Wu, J.; Wu, H.; Zhang, P. Synthesis of Pyrazoles, Diazepines, Enaminones, and Enamino Esters Using 12-Tungstenphosphoric Acid as a Reusable Catalyst in Water. *Synthesis* **2008**, *21*, 3478–3486.
14. Kidwai, M.; Bardway, S.; Mishra, N.K.; Bansai, V.; Kumar, A.; Mozumdar, S. A novel method for the synthesis of  $\beta$ -enaminones using Cu-nanoparticles as catalyst. *Catal. Commun.* **2009**, *10*, 1514–1519. [\[CrossRef\]](#)
15. Gholap, A.R.; Chakor, N.S.; Daniel, T.; Lahoti, R.J.; Srinivasan, K.V. A remarkable rapid regioselective synthesis of  $\beta$ -enaminones using silica chloride in a heterogeneous as well as an ionic liquid in a homogeneous medium at room temperature. *J. Mol. Catal. A Chem.* **2006**, *245*, 37–46. [\[CrossRef\]](#)
16. Lasić, V.; Jurković, M.; Jednacac, T.; Hrenar, T.; Vuković, J.P.; Novak, P. Intra- and intermolecular hydrogen bonding in acetylacetone and benzoylacetone derived enaminone derivatives. *J. Mol. Struct.* **2015**, *1079*, 243–249. [\[CrossRef\]](#)
17. Rodríguez, M.; Santillan, R.; López, Y.; Farrán, N.; Barba, V.; Nakatani, K.; Garcia-Baez, E.V.; Padilla-Martínez, J.J. N-H . . . O Assisted Structural Changes Induced on Ketoenamine Systems. *Supramol. Chem.* **2007**, *19*, 641–653. [\[CrossRef\]](#)
18. Hansen, P.E.; Rozwadowski, Z.; Dziembowska, T. NMR Studies of Hydroxy Schiff bases. *Curr. Org. Chem.* **2009**, *13*, 194–215. [\[CrossRef\]](#)
19. Sauer, M.; Yeung, C.; Chong, J.H.; Patrick, B.O.; MacLachlan, M.J. N-Salicylideneanilines: Tautomers for Formation of Hydrogen-Bonded Capsules, Clefts and Chains. *J. Org. Chem.* **2006**, *71*, 775–788. [\[CrossRef\]](#)
20. Hansen, P.E.; Filarowski, A. Characterisation of the PT-form of *o*-Hydroxy Acylaromatic Schiff bases by NMR Spectroscopy and DFT Calculations. *J. Mol. Struct.* **2004**, *707*, 69–75. [\[CrossRef\]](#)
21. Kwocz, A.; Panek, J.J.; Jezierska, A.; Hetmańczyk, Ł.; Pawlukoć, A.; Kochel, A.; Lipkowski, P.; Filarowski, A. A molecular roundabout: Triple cyclically arranged hydrogen bonds in light of experiment and theory. *New J. Chem.* **2018**, *42*, 19467–19477. [\[CrossRef\]](#)
22. Prbylski, P.; Wojciechowski, G.; Schilf, W.; Brezezinski, B.; Bartl, F. Spectroscopic study and PM5 semiempirical calculations of tautomeric forms of gossypol Schiff base with *n*-butylamine in the solid state and in the solution. *J. Mol. Struct.* **2003**, *646*, 161–168. [\[CrossRef\]](#)
23. Quang, T.T.; Phung, K.P.; Hansen, P.E. Schiff Bases of Gossypol. A NMR and DFT Study. *Magn. Reson. Chem.* **2005**, *43*, 302–308.
24. Dobosz, R.; Mućko, J.; Gawinecki, R. Using Chou's 5-step rule to Evaluate the Stability of Tautomers: Susceptibility of 2-[Phenylimino]-methyl]-cyclohexane-1,3-dione to Tautomerization Based on the Calculated Gibbs Free Energy. *Energies* **2020**, *13*, 183. [\[CrossRef\]](#)
25. Durgarao, N.; Hilemlan, B.; Cox, J.D.; Scott, H.; Hoang, P.; Robbins, A.; Bowers, K.; Tsebaot, L.; Miao, K.; Cataneda, M.; et al. Acceleration of the Eschenmoser coupling reaction by sonication: Efficient synthesis of enaminones. *RSC Adv.* **2013**, *3*, 181–188.

26. Chen, J.-X.; Zhang, C.-F.; Gao, W.-X.; Jin, H.-L.; Ding, J.-C.; Wu, H.-Y. B<sub>2</sub>O<sub>3</sub>/Al<sub>2</sub>O<sub>3</sub> as a New, Highly Efficient and Reusable Heterogeneous Catalyst for the Selective Synthesis of  $\beta$ -Enamino Ketones and Esters under Solvent-Free Conditions. *J. Braz. Chem. Soc.* **2010**, *21*, S1–S28. [\[CrossRef\]](#)
27. Jebari, M.; Pasturaud, K.; Picard, B.; Maddaluno, J.; Rezgui, F.; Chataignera, I.; Legros, J. “On water” reaction of deactivated anilines with 4-methoxy-3-buten-2-one, an effective butynone surrogate. *Org. Biomol. Chem.* **2016**, *14*, 11085–11087. [\[CrossRef\]](#)
28. Hansen, P.E.; Duus, F.; Neumann, R.; Jagodzinski, T.S. Deuterium Isotope Effects of O-Hydroxythioamides, Thiazolines and 5-Acyl-2-thiobarbituric Acids. *Pol. J. Chem.* **2000**, *74*, 409–420.
29. Bai, J.; Wang, P.; Cao, W.; Chen, X. Tautomer-selective derivatives of enolate, ketone and enamino by addition reaction of picolyl-type anions with nitriles. *J. Mol. Struct.* **2017**, *1128*, 645–652. [\[CrossRef\]](#)
30. Hansen, P.E.; Bolvig, S.; Kappe, T. Intramolecular Hydrogen Bonding and Tautomerism of Acyl Pyran-2,4-diones, 2,4,6-Triones and Pyridiones and Benzannelated Derivatives. Deuterium Isotope Effects on <sup>13</sup>C NMR Chemical Shifts. *J. Chem. Soc. Perkin Trans.* **1995**, *2*, 1901–1907. [\[CrossRef\]](#)
31. Hansen, P.E.; Duus, F.; Bolvig, S.; Jagodzinski, T.S. Intra-molecular Hydrogen Bonding of the Enol forms of  $\beta$ -Ketoamides and  $\beta$ -Ketothioamides. Deuterium Isotope Effects on <sup>13</sup>C Chemical Shifts. *J. Mol. Struct.* **1996**, *378*, 45–59. [\[CrossRef\]](#)
32. Kozerski, L.; Kaweck, R.; Hansen, P.E. Data for Characterization of the Geometrical Isomers of  $\beta$ -Sulfinylenamines. *Magn. Reson. Chem.* **1994**, *32*, 517–524. [\[CrossRef\]](#)
33. Afonin, A.V.; Vashchenko, A.V.; Sigalov, M.V. Estimating the energy of intramolecular hydrogen bonds from <sup>1</sup>H NMR and QTAIM calculations. *Org. Biomol. Chem.* **2016**, *14*, 11199–11211. [\[CrossRef\]](#) [\[PubMed\]](#)
34. Landge, S.M.; Thatchouk, E.; Benitez, D.; Lanfranchi, D.A.; Elhabari, M.; Goodard, W.A., III; Aprahamian, I. Isomerization Mechanism in Hydrazone-Based Rotary Switches: Lateral Shift, Rotation, or Tautomerization? *J. Am. Chem. Soc.* **2011**, *133*, 9812–9823. [\[CrossRef\]](#) [\[PubMed\]](#)
35. Nguyen, Y.H.; Lampkin, B.J.; Venkateh, A.; Ellern, A.; VanVeller, B. Open-Resonance-assisted Hydrogen Bonds and Competing quasiaromaticity. *J. Org. Chem.* **2018**, *83*, 9850–9857. [\[CrossRef\]](#)
36. Hristova, S.; Kamounah, F.S.; Molla, N.; Hansen, P.E.; Nedeltcheva, D.; Antonov, L. The possible tautomerism of the rotary switch 2-(2-(2-Hydroxy-4-nitrophenyl)hydrazono)-1-phenylbutane-1,3-dione. *Dyes Pigments* **2017**, *144*, 249–261. [\[CrossRef\]](#)
37. Su, X.; Aprahamian, I. Switching Around Two Axles: Controlling the Configuration and Conformation of a Hydrazone-Based Switch. *Org. Lett.* **2011**, *13*, 30–33. [\[CrossRef\]](#)
38. Sigalov, M.V.; Shainyan, B.A.; Chipanina, N.N.; Oznobikhina, L.; Strashnikova, N.; Sterkhova, I. Molecular Structure and Photoinduced Hydrogen Bonding in 2-pyrrolemethylidene Cyclohexanones. *J. Org. Chem.* **2015**, *80*, 10521–10535. [\[CrossRef\]](#) [\[PubMed\]](#)
39. Maeda, H.; Takeda, Y.; Haketa, Y.; Morimoto, Y.; Yasuda, N. Ion-pairing Assemblies of p-Electronic Anions formed by Intramolecular Hydrogen bonding. *Chem. Eur. J.* **2018**, *24*, 8910–8916. [\[CrossRef\]](#)
40. Sigalov, M.V.; Shainyan, B.A.; Chipanina, N.N.; Oznobikhina, L.P. Molecular Structure, Intramolecular Hydrogen Bonding, Solvent-Induced Isomerization, and Tautomerism in Azoylmethylidene Derivatives of 2-Indanone. *Eur. J. Org. Chem.* **2017**, 1353–1364. [\[CrossRef\]](#)
41. Ladarevic, J.; Bozic, B.; Natovic, L.; Nedeljkovic, B.B.; Minj, D. Role of the bifurcated intramolecular hydrogen bond on the physico-chemical profile of the novel azo pyridone dyes. *Dyes Pigments* **2019**, *162*, 562–572. [\[CrossRef\]](#)
42. Diembowska, T.; Rozwadowski, A.; Hansen, P.E. Intramolecular Hydrogen bonding of 8-hydroxyquinoline N-oxides, Quinaldinic Acid N-oxides and Quinaldinamide N-oxide. Deuterium Isotope Effects on <sup>13</sup>C Chemical Shifts. *J. Mol. Struct.* **1997**, *436–437*, 189–199. [\[CrossRef\]](#)
43. Saeed, B.A.; Elias, R.S.; Kamounah, F.S.; Hansen, P.E. A NMR, MP2 and DFT Study of Thiophenoxyketenimines (o-ThioSchiff bases). *Magn. Reson. Chem.* **2018**, *56*, 172–182. [\[CrossRef\]](#) [\[PubMed\]](#)
44. Krinsky, J.L.; Arnold, J.; Bergman, R.G. Platinum Group Thiophenoxyimine Complexes, Syntheses and Crystallographic/Computational Studies. *Organometallics* **2007**, *26*, 897–909. [\[CrossRef\]](#)
45. Corrigan, M.F.; Rae, I.D.; West, B.O. The Constitution of N-Substituted 2-(Iminomethyl) benzenethiols (o-mercaptobenzaldimines). *Aust. J. Chem.* **1978**, *31*, 587–594. [\[CrossRef\]](#)
46. Feng, Z.; Jia, S.; Chen, H.; You, L. Modulation of imine chemistry with intramolecular hydrogen bonding. Effects from ortho-OH to NH. *Tetrahedron* **2020**, *76*, 131128. [\[CrossRef\]](#)
47. Froimowicz, P.; Zhang, K.; Ishida, H. Intramolecular Hydrogen Bonding in Benzoxazines: When Structural design becomes functional. *Chem. Eur. J.* **2016**, *22*, 2691–2707. [\[CrossRef\]](#)
48. Arya, N.; Mishra, S.K.; Suryaprakash, M. Intramolecular hydrogen bond directed distribution of conformational populations in the derivatives of N'-benzylidenebenzohydrazide. *New. J. Chem.* **2019**, *43*, 13134–13142. [\[CrossRef\]](#)
49. Ren, Y.; Kravchenko, O.; Ramström, O. Configurational and Constitutional Dynamics of Enamine Molecular Switches. *Chem. Eur. J.* **2020**, *26*, 15654–15663. [\[CrossRef\]](#)
50. Petrova, M.; Mahamadejev, R.; Vigante, B.; Duburs, G.; Liepinsh, E. Intramolecular hydrogen bonds in 1,4-dihydropyridine derivatives. *R. Soc. Open. Sci.* **2018**, *5*, 1800088.
51. Piękoś, P.; Jezierska, A.; Panek, J.J.; Goremychkin, E.A.; Pozharskii, A.F.; Antonov, A.S.; Tolstoy, P.M.; Filarowski, A. Symmetry/Asymmetry of the NHN Hydrogen Bond in Protonated 1,8-Bis(dimethylamino)naphthalene. *Symmetry* **2020**, *12*, 1924. [\[CrossRef\]](#)



52. Vlasenko, M.P.; Ozeryanski, V.A.; Pozharskii, A.F.; Khoroshilova, O.V. Positively and negatively charged NHN hydrogen bonds in one molecule: Synergistic strengthening effect superbasicity and acetonitrile capture. *New J. Chem.* **2019**, *43*, 7557–7561. [\[CrossRef\]](#)
53. Dega-Szafran, Z.; Nowak-Wydra, B.; Szafran, M. Reinvestigation of  $^1\text{H}$  and  $^{13}\text{C}$  NMR Spectra of 1,8-Bis(dimethylamino)naphthalene Complexes with Mineral Acids. *Magn. Reson. Chem.* **1993**, *31*, 726–730. [\[CrossRef\]](#)
54. Pietrzak, M.; Grech, E.; Nowicka-Scheibe, J.; Hansen, P.E. Deuterium isotope effects on  $^{13}\text{C}$  chemical shifts of negatively charged NH...N systems. *Magn. Reson. Chem.* **2013**, *51*, 683–688.
55. Yamakoda, R.; Ishibashi, H.; Motoyoshi, Y.; Yasuda, N.; Maeda, H. Ion-pairing assemblies based on p-extended dipyrrolylquinoxalines. *Chem. Commun.* **2019**, *55*, 326–329. [\[CrossRef\]](#)
56. Pietrzak, M.; Try, A.C.; Andriollette, B.; Sessler, J.L.; Anzenbacher, P., Jr.; Limbach, H.-H. The largest  $^{15}\text{N}$ - $^{15}\text{N}$  Coupling Constant across an NHN Hydrogen bond. *Angew. Chem. Int. Ed.* **2008**, *47*, 1123–1126. [\[CrossRef\]](#)
57. Osmialowski, B. Conformational equilibrium and substituent effects in hydrogen bonded complexes. *Curr. Org. Chem.* **2018**, *22*. [\[CrossRef\]](#)
58. Sosnicki, J.G.; Hansen, P.E. Temperature coefficients of NH chemical Shifts of Thioamides in Relation to Structure. *Mol. Struct.* **2004**, *700*, 91–103. [\[CrossRef\]](#)
59. Desseyn, H.O.; Perlepes, S.P.; Clou, K.; Bleton, N.; van der Veken, B.J.; Dommisse, R.; Hansen, P.E. Theoretical, Structural, Vibrational, NMR and Thermal evidence of the Inter versus Intramolecular Hydrogen Bonding in Oxamides and Thiooxamides. *J. Phys. Chem. A* **2004**, *108*, 175–182. [\[CrossRef\]](#)
60. Scheiner, S. Assessment of the Presence and Strength of H-Bonds by Means of Corrected NMR. *Molecules* **2016**, *21*, 1426. [\[CrossRef\]](#) [\[PubMed\]](#)
61. Dudek, G.; Dudek, E.P. Spectroscopic Studies of Keto-Enol Equilibria. Part XIII.  $^{15}\text{N}$  Substituted Imines. *J. Chem. Soc. B* **1971**, 1356–1360. [\[CrossRef\]](#)
62. Hansen, P.E.; Saeed, B.A.; Rutua, R.S.; Kupka, T. One-bond  $^1J(^{15}\text{N},\text{H})$  coupling constants at  $\text{sp}^2$ -hybridized nitrogen of Schiff bases, enamines and similar compounds: A theoretical study. *Magn. Reson. Chem.* **2020**, *58*, 750–762. [\[CrossRef\]](#) [\[PubMed\]](#)
63. Dingley, A.J.; Grzesiek, S. Direct observation of hydrogen bonds in nucleic acid base pairs by internucleotide (2)J(NN) couplings. *S. J. Am. Chem. Soc.* **1998**, *120*, 8293–8297. [\[CrossRef\]](#)
64. Dingley, A.J.; Masse, J.E.; Peterson, R.D.; Barfield, M.; Feigon, J.; Grzesiek, S. Internucleotide Scalar Couplings across Hydrogen Bonds in Watson-Crick and Hoogsteen Base Pairs of a DNA Triplex. *J. Am. Chem. Soc.* **1999**, *121*, 6019–6027. [\[CrossRef\]](#)
65. Grzesiek, S.; Cordier, F.; Jaravine, V.; Barfield, M. Insights into biomolecular hydrogen bonds from hydrogen bond scalar couplings. *Prog. Nuc. Magn. Reson. Spectrosc.* **2004**, *45*, 275–300. [\[CrossRef\]](#)
66. Cornilescu, G.; Hu, J.S.; Bax, A. Identification of the hydrogen bonding network in a protein by scalar couplings. *J. Am. Chem. Soc.* **1999**, *121*, 2949–2950. [\[CrossRef\]](#)
67. Cornilescu, G.; Ramirez, B.E.; Frank, M.K.; Clore, G.M.; Gronenborn, A.M.; Bax, A. Correlation between  $3hJ_{\text{NC}'}$  and Hydrogen Bond Length in Proteins. *J. Am. Chem. Soc.* **1999**, *121*, 6275–6279. [\[CrossRef\]](#)
68. Wang, Y.-X.; Jacob, J.; Cordier, F.; Wingfield, P.; Stahj, S.J.; Lee-Huang, S.; Torchia, D.; Grzesiek, S.; Bax, A. Measurement of  $3hJ_{\text{NC}'}$  connectivities across hydrogen bonds in a 30 kDa protein. *J. Biomol. NMR* **1999**, *14*, 181–184. [\[CrossRef\]](#) [\[PubMed\]](#)
69. Zandarashvili, L.; Li, D.-W.; Wang, T.; Brüscheweiler, R.; Iwahara, J. Signature of Mobile Hydrogen Bonding of Lysine Side Chains from Long-Range  $^{15}\text{N}$ - $^{13}\text{C}$  Scalar J-Couplings and Computation. *J. Am. Chem. Soc.* **2011**, *133*, 9192–9195. [\[CrossRef\]](#)
70. Manalo, M.N.; Kong, X.; LiWang, A.  $^1\text{J}_{\text{NH}}$  Values show that N1...N3 Hydrogen Bonds are stronger in dsRNA A:U than dsDNA A:T Base pairs. *J. Am. Chem. Soc.* **2005**, *127*, 17974–17975. [\[CrossRef\]](#)
71. Chattopadhyay, A.; Esadze, A.; Roy, S.; Iwahara, J. NMR Scalar Couplings across Intermolecular Hydrogen Bonds between Zinc-Finger Histidine Side Chains and DNA Phosphate Groups. *J. Phys. Chem. B* **2016**, *120*, 10679–10685. [\[CrossRef\]](#)
72. Del Bene, J. *Two-Bond NMR Spin-Spin Coupling Constants across Hydrogen Bonds*. *Encyclopedia of Theoretical Chemistry*; Wiley: Weinheim, Germany, 2004.
73. Sobczyk, L.; Obzrud, M.; Filarowski, A. H/D Isotope Effects in Hydrogen Bonded Systems. *Molecules* **2013**, *18*, 4467–4476. [\[CrossRef\]](#) [\[PubMed\]](#)
74. Kozerski, L.; von Philipsborn, W. N-15 NMR spectroscopy. 9. Solvent induced Deuterium isotope effects in C-13 NMR and N-15 NMR spectra of enamines. *Helv. Chim. Acta* **1982**, *65*, 2077–2087. [\[CrossRef\]](#)
75. Hansen, P.E. Isotope effects on chemical shift in the study of intramolecular hydrogen bonds. *Molecules* **2015**, *20*, 2405–2424. [\[CrossRef\]](#) [\[PubMed\]](#)
76. Hansen, P.E.; Kaweck, R.; Krowczynski, A.; Kozerski, L. Deuterium Isotope Effects on  $^{13}\text{C}$  and  $^{15}\text{N}$  Nuclear Shielding in Intramolecular Hydrogen-bonded Compounds. II. Investigation of Enamine Derivatives. *Acta Chem. Scand.* **1990**, *44*, 826–832. [\[CrossRef\]](#)
77. Tomlinson, J.H.; Ullah, S.; Hansen, P.E.; Williamson, M.P. Characterisation of salt bridges to lysines in the Protein G B1 domain. *J. Am. Chem. Soc.* **2009**, *131*, 4674–4684. [\[CrossRef\]](#)
78. Williamson, M.P.; Hounslow, A.M.; Ford, J.; Fowler, K.; Hebditch, M.; Hansen, P.E. Detection of salt bridges to lysines in barnase in solution. *Chem. Commun.* **2013**, *49*, 9824–9826. [\[CrossRef\]](#)
79. Ullah, S.; Ishimoto, T.; Williamson, M.P.; Hansen, P.E. Ab initio Calculations of Deuterium Isotope Effects on Chemical Shifts of Salt bridged lysines. *J. Phys. Chem. B* **2011**, *115*, 3208–3215. [\[CrossRef\]](#)

80. Abildgaard, J.; LiWang, A.; Manalo, M.N.; Hansen, P.E. Peptide Deuterium Isotope Effects on Backbone  $^{15}\text{N}$  Chemical Shifts in Proteins. *J. Biomol. NMR* **2009**, *44*, 119–124. [\[CrossRef\]](#)
81. Kim, Y.-I.; Manalo, M.N.; Pérez, L.M.; LiWang, A. Computation and empirical trans-hydrogen bond deuterium isotope shifts suggest that n1-N3 A.U hydrogen bonds of RNA are shorter than those of A:T bonds of DNA. *J. Biomol. NMR* **2006**, *34*, 229–236. [\[CrossRef\]](#)
82. Manalo, M.N.; Pérez, L.M.; LiWang, A. Hydrogen-Bonding and  $\pi$ - $\pi$  Base-stacking Interactions are Coupled in DNA as Suggested by Calculated and Experimental Trans-Hbond Deuterium Isotope Shifts. *J. Am. Chem. Soc.* **2007**, *129*, 11298–11299. [\[CrossRef\]](#)
83. Sosnicki, J.G.; Langaard, M.; Hansen, P.E. Long-range Deuterium Isotope Effects on  $^{13}\text{C}$  chemical Shifts of intramolecularly hydrogen bonded N-substituted-3-(cycloamino) thiopropionamides or amides: A case of electric field effects. *J. Org. Chem.* **2007**, *72*, 4108–4116. [\[CrossRef\]](#)
84. Bolvig, S.; Hansen, P.E.; Morimoto, H.; Wemmer, D.; Williams, P. Primary Tritium and Deuterium Isotope Effects on Chemical Shifts of Compounds having an Intramolecular Hydrogen bond. *Magn. Reson. Chem.* **2000**, *38*, 525–535. [\[CrossRef\]](#)
85. Chmielewski, P.; Ozeryanski, V.A.; Sobczyk, L.; Pozharskii, A.F. primary  $^1\text{H}/^2\text{H}$  isotope effect in the NMR chemical shift of  $\text{HClO}_4$  salts of 1,8-bis(dimethylamino)naphthalene derivatives. *J. Phys. Org. Chem.* **2007**, *20*, 643–648. [\[CrossRef\]](#)
86. Pozharski, A.F.; Ryabtsova, O.V.; Ozeryanski, V.A.; Degtyarev, A.V.; Kazheva, O.N.; Alexandrov, G.G.; Dyanchenko, O.A. Organometallic synthesis, molecular structure, and coloration of 2,7-disubstituted 1,8-bis(dimethylamino)naphthalenes. How significant is the influence of “buttressing effect” on their basicity? *J. Org. Chem.* **2003**, *69*, 10109–10122. [\[CrossRef\]](#) [\[PubMed\]](#)
87. Reuben, J. Isotopic Multiplets in the Carbon-13 NMR Spectra of Aniline Derivatives and Nucleosides with Partially Deuterated Amino Groups: Effects of Intra- and Intermolecular Hydrogen Bonding. *J. Am. Chem. Soc.* **1987**, *109*, 316–321. [\[CrossRef\]](#)
88. Schaefer, T. A Relationship between Hydroxy Proton Chemical Shifts and Torsional Frequencies in Some ortho-Substituted Phenol Derivatives. *J. Phys. Chem.* **1975**, *79*, 1888–1890. [\[CrossRef\]](#)
89. Chiara, J.L.; Gomez-Sanchez, A.; Bellanato, J. Spectral properties and isomerism of nitroenamines. Part 4.  $\beta$ -Amino- $\alpha$ -nitro- $\alpha,\beta$ -unsaturated ketones. *J. Chem. Soc. Perkin Trans.* **1998**, *2*, 1797–1806. [\[CrossRef\]](#)
90. Tupikina, E.Y.; Sigalov, M.; Shenderovich, I.G.; Mulloyarova, V.V.; Denisov, G.S.; Tolstoy, P.M. Correlations of NHN hydrogen bond energy with geometry and  $^1\text{H}$  NMR chemical shift difference of NH protons for aniline complexes. *J. Chem. Phys.* **2019**, *150*, 114305. [\[CrossRef\]](#) [\[PubMed\]](#)
91. Cuma, M.; Scheiner, S.; Kar, T. Competition between rotamerization and proton transfer in *o*-hydroxybenzaldehyde. *J. Am. Chem. Soc.* **1998**, *120*, 10497–10503. [\[CrossRef\]](#)
92. Jablonski, M.; Kaczmarek, A.; Sadlej, S.J. Estimates of the Energy of Intramolecular Hydrogen Bonds. *J. Phys. Chem. A* **2006**, *110*, 10890–10898. [\[CrossRef\]](#)
93. Seyedkatouli, S.; Vakili, M.; Tayyari, S.F.; Hansen, P.E.; Kamounah, F. SMolecular structure, vibrational analysis, and intramolecular hydrogen bond strength (IHBs) of 3-methyl-4-amino-3-penten-2-one and the effect of N-methyl and N-phenyl substitutions on the IHBs. An Experimental and Theoretical approach. *J. Mol. Struct.* **2019**, *1184*, 233–245. [\[CrossRef\]](#)
94. Gholipur, A.; Neyband, R.S.; Farhadi, S. NMR investigation of substituent effects on strength the intramolecular hydrogen bonding interaction in X-phenylhydrazones switches: A theoretical study. *Chem. Phys. Lett.* **2017**, *676*, 6–11. [\[CrossRef\]](#)
95. Rozas, I.; Alkorta, I.; Elguero, J. Behavior of Ylides Containing N, O, and C Atoms as Hydrogen Bond Acceptors. *J. Am. Chem. Soc.* **2000**, *122*, 11154–11161. [\[CrossRef\]](#)
96. Hansen, P.E.; Tüchsen, E. Deuterium Isotope Effects on Carbonyl Chemical Shifts of BPTI. Hydrogen-bonding and Structure Determination in Proteins. *Acta Chem. Scand.* **1989**, *43*, 710–712. [\[CrossRef\]](#)
97. Tüchsen, E.; Hansen, P.E. Hydrogen bonding monitored by deuterium isotope effects on carbonyl  $^{13}\text{C}$  chemical shift in BPTI: Intra-residue hydrogen bonds in antiparallel  $\beta$ -sheet. *Int. J. Biol. Macromol.* **1991**, *13*, 2–8. [\[CrossRef\]](#)
98. Hansen, P.E.; Kamounah, F.S.; Saeed, B.A.; MacLachlan, M.J.; Spanget-Larsen, J. Intramolecular Hydrogen Bonds in Normal and Sterically Compressed *o*-Hydroxy Aromatic Aldehydes. Isotope Effects on Chemical Shifts and Hydrogen Bond Strength. *Molecules* **2019**, *24*, 4533. [\[CrossRef\]](#)
99. Becke, A.D. A new mixing of Hartree-Fock and local density-functional theories. *J. Chem. Phys.* **1993**, *98*, 1372–1377. [\[CrossRef\]](#)
100. Bader, R.W.F. *Atoms in Molecules: A Quantum Theory*; Oxford University Press: New York, NY, USA, 1990.
101. Kieth, T.A. *AIMAll (Version 16.10.31)*; TK Gristmill Software: Overland Park, KS, USA, 2016.
102. Hansen, P.E.; Bolvig, S.; Duus, F.; Petrova, M.V.; Kawecki, R.; Kozerski, L. Deuterium Isotope Effects on  $^{13}\text{C}$  Chemical Shifts of Intramolecularly Hydrogen-bonded Olefins. *Magn. Reson. Chem.* **1995**, *33*, 621–631. [\[CrossRef\]](#)
103. Chiara, J.L.; Gomez-Sanchez, A.; Sanchez Marcos, E. Spectral Properties and Isomerism of Nitro Enamines. Part 2. 3-Amino-2-nitrocrotonic Esters. *J. Chem. Soc. Perkin Trans.* **1990**, *2*, 385–392. [\[CrossRef\]](#)
104. Gilli, P.; Pretto, L.; Bertolasi, V.; Gilli, G. Predicting Hydrogen-Bond Strength from Acid-Base Molecular Properties. The pKa Slide Rule: Toward the Solution of a Long-Lasting Problem. *Acc. Chem. Res.* **2009**, *42*, 33–44. [\[CrossRef\]](#) [\[PubMed\]](#)
105. Gorobets, N.Y.; Yermolayev, S.A.; Gurley, T.; Gurinov, A.A.; Tolstoy, P.M.; Shenderovich, I.G.; Leadbeater, N.E. Difference between  $^1\text{H}$  NMR signals of primary amide protons as a simple spectral index of the amide intramolecular hydrogen bond strength. *J. Phys. Org. Chem.* **2012**, *25*, 287–295. [\[CrossRef\]](#)
106. Arunan, E.; Desiraju, G.R.; Klein, R.A.; Sadlej, J.; Scheiner, S.; Alkorta, I.; Clary, D.C.; Crabtree, R.H.; Dannenberg, J.J.; Hobza, P.; et al. Definition of the hydrogen bond (IUPAC Recommendations 2011). *Pure Appl. Chem.* **2011**, *83*, 1637–1641. [\[CrossRef\]](#)
107. Martyniak, A.; Majerz, I.; Filarowski, A. Peculiarities of quasi-aromatic hydrogen bonding. *RSC Adv.* **2012**, *2*, 8135–8144. [\[CrossRef\]](#)

- 
108. Adhikary, R.; Zimmermann, J.; Liu, J.; Forrest, R.P.; Janicki, T.D.; Dawson, P.E.; Corcelli, S.A.; Romesberg, F.E. Evidence of an unusual N-H . . . N Hydrogen Bond in Proteins. *J. Am. Chem. Soc.* **2014**, *136*, 13474–13477. [[CrossRef](#)]
  109. Novak, A. Hydrogen Bonding in Solids. Correlation of Spectroscopic and Crystallographic Data. *Struct. Bond.* **1974**, *18*, 177–216.
  110. Grech, E.; Malarski, Z.; Sobczyk, L. Isotope Effects in NH.N Hydrogen bonds. *Chem. Phys. Lett.* **1986**, *128*, 259–263. [[CrossRef](#)]