

## Inhibitory Effects of 17- $\alpha$ -Ethinyl-Estradiol and 17- $\beta$ -Estradiol on Transport Via the Intestinal Proton-Coupled Amino Acid Transporter (PAT1) Investigated **In Vitro and In Vivo**

Nielsen, Carsten Uhd; Pedersen, Maria; Müller, Stefanie; Kæstel, Thea; Bjerg, Maria; Ulaganathan, Nithiya; Nielsen, Salli; Carlsen, Krestine Lundgaard; Nøhr, Martha Kamp; Holm, René

*Published in:*  
Journal of Pharmaceutical Sciences

*DOI:*  
[10.1016/j.xphs.2020.08.010](https://doi.org/10.1016/j.xphs.2020.08.010)

*Publication date:*  
2021

*Document Version*  
Peer reviewed version

*Citation for published version (APA):*  
Nielsen, C. U., Pedersen, M., Müller, S., Kæstel, T., Bjerg, M., Ulaganathan, N., Nielsen, S., Carlsen, K. L., Nøhr, M. K., & Holm, R. (2021). Inhibitory Effects of 17- $\alpha$ -Ethinyl-Estradiol and 17- $\beta$ -Estradiol on Transport Via the Intestinal Proton-Coupled Amino Acid Transporter (PAT1) Investigated *In Vitro* and *In Vivo*. *Journal of Pharmaceutical Sciences*, 110(1), 354-364. <https://doi.org/10.1016/j.xphs.2020.08.010>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

### Take down policy

If you believe that this document breaches copyright please contact [rucforsk@kb.dk](mailto:rucforsk@kb.dk) providing details, and we will remove access to the work immediately and investigate your claim.

# Journal Pre-proof



Inhibitory effects of 17- $\alpha$ -ethinyl-estradiol and 17- $\beta$ -estradiol on transport via the intestinal proton-coupled amino acid transporter (PAT1) investigated *in vitro* and *in vivo*

Carsten Uhd Nielsen, Maria Pedersen, Stefanie Müller, Thea Kæstel, Maria Bjerg, Nithiya Ulaganathan, Salli Nielsen, Krestine Lundgaard Carlsen, Martha Kamp Nøhr, René Holm

PII: S0022-3549(20)30442-1

DOI: <https://doi.org/10.1016/j.xphs.2020.08.010>

Reference: XPHS 2050

To appear in: *Journal of Pharmaceutical Sciences*

Received Date: 1 July 2020

Revised Date: 12 August 2020

Accepted Date: 18 August 2020

Please cite this article as: Nielsen CU, Pedersen M, Müller S, Kæstel T, Bjerg M, Ulaganathan N, Nielsen S, Carlsen KL, Nøhr MK, Holm R, Inhibitory effects of 17- $\alpha$ -ethinyl-estradiol and 17- $\beta$ -estradiol on transport via the intestinal proton-coupled amino acid transporter (PAT1) investigated *in vitro* and *in vivo*, *Journal of Pharmaceutical Sciences* (2020), doi: <https://doi.org/10.1016/j.xphs.2020.08.010>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Inc. on behalf of the American Pharmacists Association.

**Inhibitory effects of 17- $\alpha$ -ethinyl-estradiol and 17- $\beta$ -estradiol on transport via the intestinal proton-coupled amino acid transporter (PAT1) investigated *in vitro* and *in vivo***

Carsten Uhd Nielsen<sup>a\*</sup>, Maria Pedersen<sup>a</sup>, Stefanie Müller<sup>b</sup>, Thea Kæstel<sup>b</sup>, Maria Bjerg<sup>a</sup>, Nithiya Ulaganathan<sup>a</sup>, Salli Nielsen<sup>a</sup>, Krestine Lundgaard Carlsen<sup>a</sup>, Martha Kampp Nøhr<sup>b</sup>, and René Holm<sup>c,d</sup>

<sup>a</sup>: Department of Physics, Chemistry and Pharmacy, University of Southern Denmark, Campusvej 55, DK-5230 Odense M, Denmark.

<sup>b</sup>: Department of Pharmacy, Faculty of Health and Medical Sciences, Universitetsparken 2, DK-2100 Copenhagen, Denmark

<sup>c</sup>: Drug Product Development, Janssen R&D, Johnson & Johnson, Turnhoutseweg 30, 2340 Beerse, Belgium

<sup>d</sup>: Department of Science and Environment, Roskilde University, 4000 Roskilde, Denmark

\*: Corresponding author at: Department of Physics, Chemistry and Pharmacy, University of Southern Denmark, Campusvej 55, DK-5230 Odense M, Denmark. Phone: +45 6550 9427 e-mail: [cun@sdu.dk](mailto:cun@sdu.dk) (Carsten Uhd Nielsen)

**Abstract**

The proton-coupled amino acid transporter, PAT1, is known to be responsible for intestinal absorption drug substances such as gaboxadol and vigabatrin. The aim of the present study was to investigate, if 17- $\alpha$ -ethinyl-estradiol (E-E2) and 17- $\beta$ -estradiol (E) inhibit PAT1-mediated intestinal absorption of proline and taurine *in vitro* in Caco-2 cells and *in vivo* using Sprague-Dawley rats to assess the potential for taurine-drug interactions. E and E-E2 inhibited the PAT1-mediated uptake of proline and taurine in Caco-2 cells with IC<sub>50</sub> values of 10.0 - 50.0  $\mu$ M without major effect on other solute carriers such as the taurine transporter (TauT), di/tripeptide transporter (PEPT1), and serotonin transporter (SERT1). In PAT1-expressing oocytes E and E-E2 were non-translocated inhibitors. In Caco-2 cells, E and E-E2 lowered the maximal uptake capacity of PAT1 in a non-competitive manner. Likewise, the transepithelial permeability of proline and taurine was reduced in presence of E and E-E2. In male Sprague Dawley rats pre-dosed with E-E2 a decreased maximal plasma concentration (C<sub>max</sub>) of taurine and increased the time (t<sub>max</sub>) to reach this was indicated, suggesting the possibility for an *in vivo* effect on the absorption of PAT1 substrates. In conclusion, 17- $\alpha$ -ethinyl-estradiol and 17- $\beta$ -estradiol were identified as non-translocated and non-competitive inhibitors of PAT1.

## Introduction

Solute carriers (SLCs) are a group of membrane transport proteins encoded by 396 different human genes<sup>1</sup>. The SLCs are expressed in a variety of tissues, where they facilitate the membrane transport of nutrients, vitamins, and minerals<sup>2,3</sup>. Moreover, they are important for transporting drug substances across cell membranes, and thereby facilitating the oral absorption of some drug substances. One such solute carrier is the proton-coupled amino acid transporter, PAT1, that is expressed in the luminal membrane of the intestinal epithelium and Caco-2 cells<sup>4</sup>, in intracellular vesicles of mouse renal tubular cells<sup>5</sup>, in lysosomes of rat neurons<sup>6,7</sup>, and in the nucleus of rat A7r5 smooth muscle cells<sup>8</sup>. Human PAT1 and rat Pat1 proteins consist of 476 and 475 amino acids, respectively, and have 86% amino acid identity according to the current reference sequences. PAT1 transports neutral zwitterionic amino acids such as proline, alanine, glycine, and GABA<sup>9-11</sup>, as well as drug substances such as gaboxadol, vigabatrin and  $\delta$ -aminolevulinic acid<sup>12-18</sup> with similar affinities as Pat1. Substrate transport via Pat1 is coupled to proton transport and is rheogenic resulting in transport activity under slightly acidic extracellular condition and limited activity under conditions lacking a transmembrane proton gradient<sup>19</sup>. It has been suggested that the oral absorption of gaboxadol and vigabatrin in dogs and rats driven by the transport activity of Pat1<sup>12,14,15</sup>, and that the absorption takes place in the small intestine, where PAT1 is predominantly expressed<sup>15</sup>. Taurine is also a substrate of PAT1<sup>16,20,21</sup>, and Anderson *et al.* demonstrated *in vitro*, in intestinal Caco-2 cells, that both PAT1 and the sodium/chloride-coupled taurine transporter, TauT, contributed to cellular uptake of taurine. In addition, two studies have shown that at taurine concentrations > 100  $\mu$ M, the apical uptake of taurine under slightly acidic conditions is mediated by PAT1 rather than TauT<sup>16,20</sup>. *In vivo*, it has been shown that PAT1 mediates the oral absorption of taurine in Sprague Dawley rats<sup>16</sup>, and it was suggested that radiolabeled taurine can be used as a probe (victim substrate) for investigating physiological and pharmacological inhibitory (perpetrator) effects on PAT1-mediated intestinal absorption<sup>16</sup>. In regulatory guidelines, one concern when drug substances are transported by a membrane transport protein is the possibility for drug-drug interactions in populations taken several medications<sup>22</sup>, or that substrate exposure and/or  $C_{max}$  is decreased by food or drugs<sup>23</sup>. Such interactions can arise, when either substrates or inhibitors compete for binding to the transport protein. In the FDA guidance on "In Vitro Drug Interaction Studies" two transporters P-glycoprotein (P-gp, ABCB1) and breast cancer resistance protein (BCRP, ABCG2) and seven solute carriers; organic anion transporter 1 (OAT1, SLC22A6) and 3 (OAT3, SLC22A8), organic cation transporter 2 (OCT2, SLC22A1), multidrug and toxin extrusion transporter 1 (MATE1, SLC47A1) and 2-K (MATE2-K, SLC47A2) and organic anion transporting polypeptide 1B1

(OATP1B1, SLCO1B1) and 1B3 (OATP1B3, SLCO1B3) are mentioned, but there is no guidance for quantitatively evaluating interactions for intestinal solute carriers<sup>22</sup>. Generally, one substrate may competitively inhibit the transport of another substrate, but a number of inhibitors for PAT1 have also been identified such as serotonin, L-tryptophan and 5-hydroxy-tryptophan (IC<sub>50</sub> of 5.9, 4.7 and 0.9 mM)<sup>24</sup>, dipeptides such as Gly-Tyr, Gly-Pro, and Gly-Phe (IC<sub>50</sub> of approx. 30-50 mM)<sup>25</sup>, and the anti-depressant sertraline (IC<sub>50</sub> of 177-241 µM)<sup>26</sup>, opening the possibility of both food components and drug substances having the potential to alter PAT1-mediated (victim) bioavailability. In oocytes injected with *PAT1 cRNA* it has been proposed that 17- $\alpha$ -estradiol and 17- $\beta$ -estradiol decrease glycine induced inward currents at concentrations of 100 µM<sup>27</sup>. Shan *et al.* suggested that these estrogens bind to PAT1 and thereby change the conformation of the PAT1 protein from an open to a closed state, thereby decreasing transport activity<sup>27</sup>. Estradiol is an endogenous hormone, but it is also used as an oral hormone supplement to treat conditions related to menopause<sup>28</sup> or prostate cancer<sup>29</sup>. Depending on the disease, estradiol is given at oral doses between 1-10 mg daily, which results in a relatively low bioavailability of 2-10 %<sup>30</sup>. A structurally related hormone is 17- $\alpha$ -ethinyl-estradiol, which is used as an oral contraceptive given at daily doses of 30 µg in combination with levonorgestrel. 17- $\alpha$ -ethinyl-estradiol is more resistant to metabolism than estradiol and has an oral bioavailability ranging from 43-74 % (see references in<sup>31</sup>). Based on the observation that estradiol inhibits inward-directed PAT1 mediated H<sup>+</sup>-currents in oocytes<sup>27</sup>, we hypothesized that 17- $\alpha$ -ethinyl-estradiol could also interact with PAT1. The aim of the present study was thus to investigate, if 17- $\beta$ -estradiol and 17- $\alpha$ -ethinyl-estradiol inhibit PAT1-mediated proline and taurine transport *in vitro*, and if there could be a potential oral absorption interaction with 17- $\beta$ -estradiol or 17- $\alpha$ -ethinyl-estradiol taking place *in vivo* using Sprague Dawley rats as a preclinical model and taurine as the victim model drug substance.

## Material and methods

### Materials

Caco-2 cells were obtained from Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ) (Leibniz Institute, Braunschweig, Germany). Transwell inserts were from Corning Life Sciences (New York City, NY, USA) and all other cell culture plastic products from Sigma-Aldrich (Merck KGaA, Darmstadt, Germany). Dulbecco's Modified Eagle's Medium (DMEM), taurine (Tau), L-proline (Pro), chenodeoxycholic acid (CDCA), cholesterol, dexamethasone (Dx), dimethyl sulfoxide >99.9 % (DMSO), diarylpropionitrile (DPN), 17- $\beta$ -

estradiol (estradiol, E2), 17- $\alpha$ -ethinyl-estradiol (ethinyl-estradiol, E-E2), fulvestrant, glycocholic acid (GCA), genistein, L-glutamine (Gln), glycyl-L-proline (Gly-Pro), non-essential amino acids (NEAA), 2-(4-morpholino)ethanesulfonic acid (MES), penicillin/streptomycin, L-tryptophan (Trp), (E/Z)-endoxifen hydrochloride hydrate, staurosporine, phorbol 12-myristate 13-acetate (PMA), sodium chloride, potassium chloride, sodium bicarbonate, calcium nitrate, magnesium sulphate heptahydrate, calcium chloride dihydrate and magnesium chloride was purchased from Sigma-Aldrich (Merck KGaA, Darmstadt, Germany). Bovine serum albumin (BSA), sodium bicarbonate solution 7.5 %, fetal bovine serum (FBS) and Hank's Balanced Salt Solution (HBSS) with calcium and magnesium without sodium bicarbonate and phenol red were obtained from Thermo Fisher Scientific (Waltham, MA, USA). 4-(2-hydroxyethyl)-1-piperazineethansulfonic acid (HEPES) and Triton x-100 was obtained from AppliChem GmbH (Darmstadt, Germany). Ethanol >99.7 % (EtOH) was obtained from VWR (Radnor, PA, USA), and ultrapure water was obtained in house from Water Purification systems Milli-Q Gradient, Millipore (Merck, Darmstadt, Germany). L-[2,3,4,5- $^3\text{H}$ ]-proline (75 Ci mmol $^{-1}$ , radiochemical purity >97%), 5-[1,2- $^3\text{H}$ [N]]-hydroxytryptamine (5-HT) creatinine sulphate (28.3 Ci mmol $^{-1}$ , radiochemical purity >95%), D-[1- $^{14}\text{C}$ ]-Mannitol (57.2 mCi mmol $^{-1}$ , radiochemical purity >97%), [2,2- $^3\text{H}$ (N)]-Taurine (19.1 Ci mmol $^{-1}$ , radiochemical purity >97%) and Ultima Gold scintillation liquid were purchased from Perkin Elmer (Waltham, MA, USA). [ $^{14}\text{C}$ ]-Gly-Sar (56 mCi mmol $^{-1}$ , radiochemical purity >95%) and DL-[4- $^3\text{H}$ ]-propranolol hydrochloride (15.2 Ci mmol $^{-1}$ , radiochemical purity >97%) was obtained from GE Healthcare (Little Chalfont, UK). Thapsigargin was a kind gift from Søren Brøgger Christensen, University of Copenhagen. All other chemicals were of reagent or laboratory grade.

## Methods

### Cell culture

Caco-2 cells were seeded on either Transwell® polycarbonate membrane inserts (1.12 cm $^2$ , 0.4  $\mu\text{m}$  pore size) or 24 wells cell culture plates (1.90 cm $^2$ ) at a density of  $8.9 \cdot 10^4$  cells cm $^{-2}$ . Experiments were performed either 6 days or 13-15 days post-seeding on 24 wells plates or insert, respectively. The cells were maintained at 37 °C in Dulbecco's modified Eagle medium (DMEM) supplied with 10 % fetal bovine serum (FBS), penicillin (10 U ml $^{-1}$ ), streptomycin (10  $\mu\text{g}$  ml $^{-1}$ ), L-glutamine (2 mM), and non-essential amino acids (1 %) in an atmosphere of 5 % CO $_2$  and a 90 % relative humidity. The culture medium was changed every 2-3 days.

### ***Xenopus Laevis* Oocytes and Two-Electrode Voltage Clamp Measurements**

*SLC36A1* cRNA was synthesized by *in vitro* transcription of linearized pGEM-HE-*SLC36A1* as previously described by Frolund *et al.*<sup>32</sup> using the mMESAGE mMACHINE T7 mRNA-capping kit according to the protocol supplied by Invitrogen (Thermo Fisher Scientific, Waltham, MA, USA). Stage VI oocytes from *Xenopus laevis* frogs (collagenase treated) were injected with 18.4 nL of diluted *SLC36A1* cRNA (0.74  $\mu\text{g } \mu\text{L}^{-1}$ ) or nuclease-free water using an UMP3 Microsyringe Injector with an Micro4 Controller (World Precision Instruments Ltd, Sarasota, FL, USA) and incubated at 18 °C. The oocytes were stored in Modified Barth's Saline (in mM: NaCl, 88; KCl, 1.0; NaHCO<sub>3</sub>, 2.4; Ca(NO<sub>3</sub>)<sub>2</sub>, 0.30; MgSO<sub>4</sub>, 0.82; CaCl<sub>2</sub>, 0.40; HEPES, 15; penicillin (10 U ml<sup>-1</sup>), streptomycin (10  $\mu\text{g ml}^{-1}$ ), pH 7.6 and 183 mOsmol kg<sup>-1</sup>. Two-electrode voltage clamp (TEVC) measurements were performed using a Xenoplax Workstation and a VC<sup>38</sup> perfusion system from ALA Scientific Instruments and a TURBO TEC-10CX amplifier with an INT-20X breakout box from NPI electronic (Tamm, Germany). CellWorks Software version 6.2.1 were used for data acquisition and solution handling. The PC-10 puller from Narishige Group (Tokyo, Japan) was used to produce micropipettes from glass capillaries of fire-polished borosilicate glass and used for recordings with a resistance range between 2.5-4.5 M $\Omega$ . TEVC were performed 8 days post-injection, and oocytes were continuously perfused with Ringer's solution (in mM: NaCl, 115; KCl, 2.5; CaCl<sub>2</sub>, 1.8; MgCl<sub>2</sub>, 0.1; MES, 10), pH 6.0 and 242 mOsmol kg<sup>-1</sup> at a holding potential of -60 mV. Compounds were dissolved in Ringer's solution and pH was adjusted to 6.0. Solutions containing either 20 mM proline, 50  $\mu\text{M}$  E2 or 50  $\mu\text{M}$  E-E2 alone or in combination were added by full bath application for 30 s between 120 s periods of perfusion with buffer, while the change in membrane current was monitored continuously. Water-injected oocytes served as negative controls. All experiments were performed on 6 oocytes.

### **Buffer solutions for Caco-2 cell experiments**

The donor solutions were prepared in Hank's Balanced Saline Solution (HBSS) consisted of, in mM: CaCl<sub>2</sub>, 1.26; MgCl<sub>2</sub>, 0.49; MgSO<sub>4</sub>, 0.41; KCl, 5.33; KH<sub>2</sub>PO<sub>4</sub>, 0.44; NaCl, 138; Na<sub>2</sub>HPO<sub>4</sub>, 0.34; D-glucose, 5.56; NaHCO<sub>3</sub>, 4.17 and contained 10 mM MES at pH 6.0 (HBSS 6.0) or 10 mM HEPES pH 7.4 (HBSS 7.4). The pH of the buffers was adjusted with 0.1 M NaOH/HCl using a pH-meter (Mettler Toledo, Columbus, OH, USA). CDCA, Dx, E2, E-E2, fulvestrant, GCA, and genistein were prepared as stock solutions in 99 % ethanol. Thapsigargin, PMA, and staurosporine were prepared as stock solutions in DMSO. For cell experiment these stock solutions were diluted in buffer, where the final DMSO or ethanol concentrations did not exceed 1 %.



### Uptake experiments in Caco-2 cells

The uptake of taurine via TauT or PAT1, proline via PAT1, Gly-Sar via the proton-coupled di/tri-peptide transporter, (PEPT1, SLC15A1) 5-HT via the sodium-dependent serotonin transporter, (SERT1, SLC6A4) or propranolol via passive diffusion was investigated in HBSS buffer at either pH 6.0 or pH 7.4 for 5 min. Previous studies have shown that uptake of PAT1 and PEPT1 substrates in Caco-2 cells is linear for 5 min<sup>33-35</sup>. Caco-2 cells grown on 24 well plates (1.9 cm<sup>2</sup>) were used for the uptake study. All uptake experiments were performed using 1  $\mu\text{Ci mL}^{-1}$  radiolabeled compound alone or in buffer solution supplemented with a non-labeled concentration of substrate and/or E2 or E-E2. Before starting the experiment, all solutions were pre-heated to a temperature of 37 °C. The cells were incubated with pre-warmed 10 mM HEPES in HBSS (pH 7.4) for at least 10 min on a mini shaker (220 rpm) from Troemner (Thorofare, NJ, USA) at 37 °C to equilibrate the cells prior to the experiment. Buffer was removed by gentle vacuum suction and 500  $\mu\text{L}$  solution containing radiolabeled compound was applied on the apical side. After the 5 min incubation, the isotope-containing buffer solution was removed, and the wells were washed three times with 500  $\mu\text{L}$  ice-cold 10 mM HEPES in HBSS (pH 7.4). The cells were detached using 200  $\mu\text{L}$  of 0.1 % Triton X-100 and transferred to scintillation vials and 2 mL of scintillation liquid was added. 20  $\mu\text{L}$  donor samples were taken from each solution containing radiolabeled compound. The samples were vortexed for 10 s, and the amount of radiolabeled compound (as disintegration per minute, DPM) was measured for 10 min using liquid scintillation counters (LSC) (Tri-Carb 2900TR or TriCarb 4910TR, Perkin Elmer, Waltham, MA, USA).

### Transport of substances across Caco-2 cell monolayers

Caco-2 cells grown on Transwell<sup>®</sup> inserts (1.12 cm<sup>2</sup>, 0.4  $\mu\text{m}$ ) were used for the transport study. The transepithelial electrical resistance (TEER) was measured at room temperature (20 °C) in a tissue resistance measurement chamber (Endohm-12) with an Epithelial Voltohmmeter (EVOM 2) from World Precision Instruments (Sarasota, FL, USA). The initial transepithelial electrical resistance (TEER) was measured to assure that the barrier properties of the monolayer were intact, and all monolayers used had values > 400  $\Omega\text{ cm}^{-2}$ . Transport was measured from the apical (A) to the basolateral (B) side. Before starting the experiment, all solutions were pre-heated to a temperature of 37 °C. The cells were incubated with pre-warmed 10 mM HEPES in HBSS (pH 7.4) for at least 10 min on a mini shaker (220 rpm) from Troemner (Thorofare, NJ, USA) at 37 °C to equilibrate the cells prior to the experiment. Buffer was removed by gentle vacuum suction, and 500  $\mu\text{L}$  solution containing radiolabelled compound (1  $\mu\text{Ci mL}^{-1}$ ) was applied on the apical side. Following this, 1000  $\mu\text{L}$

10 mM HEPES in HBSS (pH 7.4) was applied to the basolateral side. Samples, 100  $\mu$ L, were taken from the receiver (basolateral) solution at 15, 30, 60, 90, and 120 min after starting the experiment and replaced with 100  $\mu$ L fresh 10 mM HEPES in HBSS (pH 7.4). 20  $\mu$ L donor samples were taken from each solution containing radiolabelled compound at 0 and 120 min. To all samples, 2 mL scintillation liquid was added and vortexed for 10 s. The amount of radiolabelled compound (as disintegration per minute, DPM) was measured for 10 min using liquid scintillation counter (LSC) (Tri-Carb 2900TR, Perkin Elmer, Waltham, MA, USA). The integrity of the Caco-2 cell monolayer was evaluated in each transport study by measuring the flux of  $^{14}\text{C}$  or  $^3\text{H}$ -mannitol. None of the treatments affected mannitol permeabilities significantly, with apparent mannitol permeabilities always being lower than  $5.2 \cdot 10^{-7} \pm 2.2 \cdot 10^{-7} \text{ cm s}^{-1}$ .

### **Estradiol and ethinyl-estradiol formulations for the pharmacokinetic (PK) study**

E2 and E-E2 were dissolved in 96 % EtOH to a concentration of 2.5 mM, and this stock solution was further diluted in 96% EtOH and given in doses of 0, 0.25, 1, 5, 10 and 25  $\mu\text{mol kg}^{-1}$  for E2 and 0, 0.25, 1, 5 and 25  $\mu\text{mol kg}^{-1}$  for E-E2. Taurine was dissolved in Elga water and dosed at 30  $\text{mg kg}^{-1}$ . The osmolalities were measured using a Wescor Vapro model 5520 vapour pressure osmometer (Wescor Inc, Logan, UT, USA) and tonicity was adjusted with sodium chloride to 274-295  $\text{mOsmol kg}^{-1}$ . pH values were measured using PHM220 LAB pH meter by Radiometer analytical SAS (Hach, Loveland, CO, USA) and was adjusted with NaOH or HCl to  $7.4 \pm 0.05$ . Immediately before dosing, [ $^3\text{H}$ ]-taurine was added to each formulation containing taurine in a concentration of 40  $\mu\text{Ci kg}^{-1}$ .

### ***In vivo* taurine PK studies after pre-administration with E2 or E-E2**

Male Sprague-Dawley rats, weighing approximately 350 g (287-398 g) on the day of the experiments, were purchased from Charles River Deutschland (Sulzfeld, Germany). The animals were acclimatised for a minimum of five days in groups of two and maintained on standard feed with free access to water. Prior to dosing, the animals were fasted for 16-20 h with free access to water. The protocol used for the animal studies was approved by the institutional animal ethics committee in accordance with Danish law regulating animal experiments and in compliance with EC Directive 2010/63/EU and the NIH guidelines on animal welfare.

E2 or E-E2 and taurine were dosed separately. Initially, E2 or E-E2 were pre-dosed at 0.3  $\text{mL kg}^{-1}$  and four and a half minute later the taurine formulations were administered at a dose of 30  $\text{mg kg}^{-1}$  (at 10  $\text{mL kg}^{-1}$ ). Animals not receiving E2 or E-E2 were pre-dosed with 0.3  $\text{mL kg}^{-1}$  96 % EtOH. All formulations were dosed

orally by gavage. Each group dosed with E2 or E-E2 consisted of four male Sprague-Dawley rats and one group of eight rats (control group) were initially dosed with 96 % ethanol only then with 30 mg kg<sup>-1</sup> taurine.

### Plasma sampling and sample analysis

The blood samples (100-200 µL), collected after administration, were drawn from the tail vein and collected into EDTA coated tubes (Microwette 500 K3E, Sarstedt, Nümbrecht, Germany) at 5-360 min after administration (5, 15, 30, 45, 60, 120, 180, 240, 360 min). Plasma was harvested immediately by 10 min of centrifugation at 4 °C, 3600 × g (Multifuge 1 S-R, Heraeus, Hanau, Germany) and stored in polypropylene tubes (Pony Vial, PerkinElmer, Waltham, MA, USA) at -20 °C until analysed. At the end of the experiment, animals were sacrificed by spinal dislocation using a guillotine. Plasma samples were left to defrost and mixed with 4 mL of liquid scintillation cocktail. The samples were counted for 3 min on a liquid scintillation counter (Tri-Carb 2900TR, Perkin Elmer, Waltham, MA, USA).

### Data analysis

The inhibition of total uptake of proline and taurine at pH 6.0 in Caco-2 cells was described by

$$V_o = \text{Bottom} + \frac{\text{Top} - \text{Bottom}}{1 + \frac{[I]}{IC_{50}}}$$

where  $V_o$  is the initial uptake rate of taurine or proline in the presence of various concentrations of E2 or E-E2 [I], Top is maximal uptake rate of taurine or proline in the absence of inhibition, and bottom is the uptake rate at maximal degree of inhibition. The total uptake of proline and taurine at pH 6.0 in Caco-2 cells was described by Michaelis-Menten like kinetics:

$$V_o = \frac{V_{\max} \cdot [S]}{K_m + [S]}$$

where  $V_o$  is the initial uptake rate of taurine or proline at a given concentration, [S], in mM in the donor solution,  $V_{\max}$  is the maximal uptake rate, and  $K_m$  is the Michaelis constant.

Transport data was described by Fick's first law;

$$J = \frac{Q}{A \cdot t} = P_{\text{app}} \cdot C_0$$

Where  $J$  is the flux and  $Q$  is the steady-state amount of transported compound over a given absorptive area ( $A$ ), per time ( $t$ ) and  $C_0$  is the starting concentration in the donor chamber. From the steady state flux the apparent permeability coefficient ( $P_{app}$ ) was calculated.

The pharmacokinetic parameters of orally administered taurine in male Sprague-Dawley rats were estimated non-compartmentally using Phoenix<sup>®</sup> 6.3 version (Pharsight Corporation, A Certara Company, USA).  $C_{max}$  and  $t_{max}$  were found as mean values of the plasma concentration profiles within each group. Area under the plasma concentration versus time profiles (AUC) were determined with non-compartmental analysis by using the linear trapezoidal rule from time zero to  $C_{max}$  and by log linear methods from  $C_{max}$  to the last measured plasma concentration and denoted  $AUC_{0-tlast}$ . The first order elimination rate constant  $k_e$  was calculated as the slope from the terminal log plasma concentration time curve of individual animals. The total clearance was calculated as the dose divided by AUC.

### Statistical analysis

Statistical analysis was performed in GraphPad Prism version 7.02. The obtained data was analysed for statistical differences using a parametric one-way ANOVA test followed by Dunnett's multiple comparison test. The following levels of significance was used  $p < 0.05$  (\*). Values were expressed as mean  $\pm$  SEM.

### Results

#### Estradiol and ethinyl-estradiol inhibit PAT1-mediated uptake of Taurine and Proline

The uptake of solute carrier substrates such as taurine, proline, glycyl-sarcosine (Gly-Sar) and 5-hydroxytryptamine was investigated in Caco-2 cells in the presence of 100  $\mu$ M E2 and E-E2 (Figure 1). The uptake of both taurine and proline was higher under slightly acidic condition compared to neutral pH in the uptake buffer (Fig. 1 A and B), consistent with proton-coupled transport via PAT1. 100  $\mu$ M E2 and E-E2 inhibited the uptake of proline and taurine at pH 6.0, but not at pH 7.4. Likewise, the uptake of 5-hydroxytryptamine was slightly, yet significantly, decreased by 100  $\mu$ M E2 and E-E2 (Fig. 1C). The uptake of Gly-Sar via hPEPT1 was unaffected by 100  $\mu$ M E2 and E-E2 but inhibited by 10 mM Gly-Pro (Fig. 1D). The uptake of propranolol was unaffected by 100  $\mu$ M E2 (Fig. 1E).

### **Estradiol and ethinyl-estradiol concentration-dependently inhibit PAT1-mediated uptake of taurine and proline**

The uptake of taurine and proline was investigated in Caco-2 cells in the presence of increasing concentrations of E and E-E2 (Fig. 2). Under slightly acidic extracellular conditions E2 and E-E2 inhibited proline uptake with  $IC_{50}$ -values of  $10.0 \pm 3.2 \mu\text{M}$  and  $50.0 \pm 14.3 \mu\text{M}$ , respectively (Fig. 2A). The uptake of taurine was inhibited by E and E-E2 with  $IC_{50}$ -values of  $8.7 \pm 5.1 \mu\text{M}$  and  $22.9 \pm 15.5 \mu\text{M}$ , respectively (Fig. 2B). In contrast, the uptake of taurine at neutral extracellular pH was not affected by E2 or E-E2 in the concentration range investigated (Fig. 2C).

### **Estradiol and Ethinyl-estradiol inhibit proline induced currents in PAT1 expressing oocytes**

To investigate the effect of E and E-E2 on PAT1-mediated transport in a system only expressing PAT1, a series of TEVC recordings were conducted on PAT1 expressing oocytes (Fig. 3). Proline induced inward directed currents in PAT1 expressing oocytes, whereas no change in baseline current was observed in water-injected oocytes (upper trace in Figure 3). The proline induced signal was attenuated, when E2 and E-E2 were present in the proline-containing solution being perfused over the oocyte. Buffers with E2 and E-E2 alone did not induce an inward current, showing that they were non-translocated inhibitors of PAT1.

### **Estradiol and ethinyl-estradiol competitively inhibit PAT1-mediated cellular uptake of proline and taurine.**

The concentration-dependent uptake of proline and taurine was measured in Caco-2 cells under slightly acid conditions (Fig. 4). The uptake of proline was saturable and in the presence of E2, the  $K_m$ -value remained statistically similar, while the  $V_{max}$  value decreased significantly with increasing E2 concentrations (Fig. 4A and Table 1). A similar pattern was observed for taurine uptake in the presence of E2 and E-E2 (Fig. 4B and Table 1). This inhibition pattern suggested a non-competitive interaction and based on this, the inhibitory constants were estimated. The  $K_i$  value for E2 inhibition of proline uptake was  $46.4 \pm 0.6 \mu\text{M}$  and  $16.1 \pm 2.1 \mu\text{M}$  for taurine uptake. For E-E2, the  $K_i$  value for inhibition of taurine uptake was  $30.3 \pm 2.7 \mu\text{M}$ .

### **Time-dependency of estradiol and ethinyl-estradiol treatment on PAT1-mediated uptake.**

The results were obtained by concomitantly applying hormone and substrate to Caco-2 cells. Therefore, the time dependency of the ability of E2 and E-E2 to decrease PAT1-mediated uptake was investigated. A one-

hour preincubation with 100  $\mu$ M E2 or E-E2 showed that the 5 min effect observed was not transient as it persisted by a prolonged incubation with E2 or E-E2 (Fig 5A). Since E2 and E-E2 are hydrophobic substances, the ability to wash out the effect of the compounds was investigated. E2 was added to the cells for 5 min followed by a buffer wash-out period before measuring the uptake of proline. This showed that approximately 15 min after removal of E2 the inhibitory effect was removed, as the uptake at 15 min was not significantly different from the control uptake (Fig. 5B).

### **PAT1-mediated uptake is not inhibited by other steroid-containing structures or modulators of estradiol signaling**

To investigate if the effect observed with E2 and E-E2 could be related to hydrophobic interactions with the Caco-2 cells, a series of steroid skeleton containing structures were investigated at a similar concentration as used for estradiol and ethinyl-estradiol (Fig. 6A). From this it was evident that, at the concentrations used, only E2 and E-E2 inhibit PAT1-mediated uptake in Caco-2 cells.

Estradiol signaling pathway modulators were also investigated (Fig. 6B). Although PI-103 reduced the uptake of proline, none of the investigated compounds were able to reverse the reduction of PAT1-mediated transport.

### **Absorptive transport of taurine and proline across Caco-2 cell monolayers**

The absorptive (apical to basolateral) transport of taurine in the presence of 100  $\mu$ M E2 and E-E2 was determined across Caco-2 cell monolayers in the presence or absence of a proton gradient across the cell layer (Fig. 7 A and B). In the absence of a proton-gradient and with a TauT-facilitating taurine concentration neither E2 nor E-E2 changed the permeability of taurine. When the apical pH was adjusted to 6.0 and the taurine concentration to 100  $\mu$ M, the presence of 100  $\mu$ M E2 and 100  $\mu$ M E-E2 significantly reduced taurine permeability. Likewise, the permeability of 1 mM proline was reduced, while the permeability of Gly-Sar, 5-HT and propranolol was statistically unaffected by the presence of E2 and E-E2. These results showed that transepithelial transport of taurine and proline was decreased by E2 and E-E2 and that the effect of E2 and E-E2 was relatively specific to PAT1. This indicated that a decreased PAT1 transport activity in the apical membrane of Caco-2 cells also translated into a decreased absorptive transport, whereby PAT1 mediated cellular uptake was rate determining for *in vitro* absorption across Caco-2 cells.

### **Pharmacokinetic interaction study of taurine with estradiol and ethinyl-estradiol**

Since E2 or E-E2 decreased PAT1-mediated transepithelial permeability of proline and taurine, it was investigated if both estrogens had an effect on the pharmacokinetic profile of orally administered taurine to assess the potential for a “drug”-drug interaction. Male Sprague-Dawley rats were orally pre-dosed with E2 or E-E2 in 96 % ethanol in various doses and the resulting plasma concentrations versus time profiles are shown in Figure 8. The pharmacokinetic parameters ( $t_{max}$ ,  $C_{max}$ ,  $AUC_{0-tlast}$  and  $k_e$ ) obtained from a non-compartmental analysis are summarized in Table 2 and 3. The pharmacokinetic profile of taurine after pre-administration of E2 were generally below the control profile, however no significant differences were found when comparing  $t_{max}$ ,  $AUC_{0-tlast}$ ,  $C_{max}$  or  $k_e$  to that of the control. In the case of E-E2, the pharmacokinetic profiles of the different treatments showed lower plasma concentrations than the control group, and there appeared to be a concentration-dependent decrease in  $C_{max}$  and a prolongation of  $t_{max}$ . However, when evaluating the pharmacokinetic parameters using an ANOVA analysis, no significant differences could be shown. Likewise, no significant differences between  $AUC_{0-tlast}$  of the control group only administered taurine and the groups administered E-E2 in various doses could be shown. However, when comparing all the sample groups, a significant difference ( $p < 0.05$ ) was observed in  $AUC_{0-tlast}$  between the group receiving  $5 \mu\text{mol kg}^{-1}$  E-E2 and the group receiving  $25 \mu\text{mol kg}^{-1}$  E-E2.

## Discussion

It is clear from the present study that E2 and E-E2 interacted with PAT1, and thereby reduced substrate transport *in vitro*. Since E2 and E-E2 did not induce currents in PAT1 expressing oocytes, they were not substrates of the carrier, something that would not be expected based on the structural difference between an amino acid and a steroid. The PAT1-mediated transport of taurine and proline was decreased in a non-competitive manner, and collectively this suggested that E2 and E-E2 were inhibitors not binding to the amino acid substrate binding site of the PAT1 protein. To our knowledge, E2 and E-E2 are the most potent PAT1 inhibitors reported in the literature with affinities approximately 3-20 times higher than sertraline<sup>26</sup>. Moreover, the effect of E2 appeared within 5 min and could be rapidly washed out over a 15 min period, which suggested a direct and non-genomic interaction. In other short-term experiments, E2 has been shown to inhibit transport via other membrane transport proteins: In striatal synaptosomes from ovariectomized Sprague Dawley rats, E2 decreased dopamine uptake in a competitive manner with an  $IC_{50}$  value of  $7.2 \pm 0.6 \mu\text{M}$ <sup>36</sup>, E2 inhibited L-type  $Ca^{2+}$  channels in A7r5 cells with an  $IC_{50}$  value of  $14.24 \pm 0.05 \mu\text{M}$ <sup>37</sup>, and non-competitively inhibited mouse 5-HT<sub>3</sub> receptors with an  $IC_{50}$  value of  $33.2 \mu\text{M}$ <sup>38</sup>. The  $IC_{50}$  values for direct non-genomic inhi-

bition by E2 and E-E2 on PAT1 transport obtained in the present study are thus quite comparable with previous findings for other membrane proteins.

E2 can exert several cellular responses related to membrane, cytosolic or nuclear effects as reviewed by Nadal *et al*<sup>39</sup>. Therefore, in Caco-2 cells E2 and E-E2 could also have other short-term effects besides PAT1 binding, which could affect the transport through regulation. However, considering that the present study did not include a treatment for more than two hours, effects of E2 and E-E2 related to altered gene transcription seemed unlikely. There are membrane targets, including estrogen and receptor(s) related to the classic nuclear estrogen receptors (ER), non-classic ERs, and ligand- and voltage-activated ion channels or cytosolic targets that could trigger signaling events by intracellular messengers, resulting in the activation of different kinases, e.g. by ER $\alpha$  and ER $\beta$ , which activate different signaling pathways<sup>39</sup>. In the present study estrogen receptor (ER) activation, phosphorylation, intracellular Ca<sup>2+</sup> signaling, and PI3K activation was investigated. These signaling events were studied with the ER antagonists endoxifen and fulvestrant, the PKC inhibitor staurosporine, the PI3K inhibitor PI-103, and the SERCA pump inhibitor thapsigargin all at concentrations known to affect the pathways. However, none of the pharmacological treatments employed altered E2-induced effects, supporting a direct E2 and E-E2 effect on PAT1 instead of an effect caused by an intracellular signaling pathway. In epithelial cells, transport via PAT1 is proton-coupled, and the hormones could affect the proton-coupling, by interacting with the sodium-proton exchanger NHE3. It has been shown that both PAT1 and the di/tri-peptide transporter PEPT1 (SLC15A1) transport is dependent on NHE3 to maintain the transmembrane proton gradient<sup>40-42</sup>. Moreover, it has been shown that E2 is able to up-regulate the activity of NHE3 in rat colonic epithelial cells, although this effect was observed after long-term treatment with E2<sup>43</sup>. However, E2 and E-E2 did not affect PEPT1-mediated transport after short-term exposure suggesting that the proton gradient was not altered by E2 or E-E2. Moreover, transport via TauT and SERT1, as well as the passively transported propranolol, was not altered by the highest investigated concentration of E2 and E-E2, and related steroid-structure containing compounds did not inhibit PAT1-mediated transport neither. The only slightly contradiction to this was that E2 slightly decrease the uptake of 5-HT, yet this effect was not found when measuring transepithelial transport of 5-HT. Collectively, the ability of E2 and E-E2 to inhibit PAT1-mediated transport appeared to be a specific interaction and not a result of unspecific cellular effects or through E2 signalling pathways. It remains unknown what the physiological reason for the ability of E2 to modulate PAT1-mediated transport is and under which conditions E2 concentration are increased to the  $\mu\text{M}$  range, as circulating E2 concentration are in the pM range<sup>44</sup>.



The likelihood of a potential *in vivo* effect can be predicted by considering the  $IC_{50}$  value of inhibition relative to the initial concentration of inhibitor in the human intestine. There are no regulatory guidance for assessing the impact of intestinal solute carriers, but for P-gp and BCRP the drug-drug interaction (DDI) index is calculated as  $I_{gut}/IC_{50}$ , where  $I_{gut}$  is the dose ( $\mu\text{mol}$ ) calculated from 1 or 10 mg E2 or 0.03 mg E-E2 divided by 250 mL water ( $I_{gut}$  14.7 – 147  $\mu\text{M}$  for E2 and  $I_{gut}$  0.4  $\mu\text{M}$  for E-E2), and a value  $\geq 10$  suggests a potential *in vivo* interaction<sup>22</sup>. Neglecting solubility issue, this translates into DDI values of 1.5-14.7 for E2 using proline as a substrate and 1.5-16.9 using taurine as a substrate, and approximately 0.01 for E-E2 using proline as a substrate. Thus, in human the interaction of E2 with PAT1 would clearly warrant further *in vivo* investigations, while interactions between proline or taurine and E-E2 would appear less likely. To mechanistically investigate if E2 and E-E2 could decrease PAT1-mediated absorption *in vivo*, an interaction study was conducted in Sprague-Dawley rats. Taurine was chosen as victim substrate for this investigation, since a previously study suggested PAT1 mediated absorption in the relevant doses in Sprague Dawley rats<sup>16</sup>. The doses of E2 and E-E2 administered orally to the rats covered a broad range of 0.25–25  $\mu\text{mol kg}^{-1}$  (approximately 0.07–7  $\text{mg kg}^{-1}$ ), where the human doses would be approximately 0.14  $\text{mg kg}^{-1}$  or 0.0004  $\text{mg kg}^{-1}$  for E2 and E-E2, respectively. The E-E2 doses used in the study are thus much higher than used in human, and assuming a rat intestinal volume of 3 mL, initial rat  $I_{gut}$  of E-E2 and E2 would be 2.5 – 250  $\mu\text{M}$ . These concentrations are in the range of the obtained  $IC_{50}$  values for E2 and E-E2. E-E2 decreased the exposure of taurine in a concentration-dependent manner with decreasing AUC and  $C_{max}$  and increasing  $t_{max}$ , although this did not meet statistical significance. After E2 administration, all taurine AUC and  $C_{max}$  values were lower than in the absence of E2, yet, the ability of E2 to alter taurine pharmacokinetics were less evident than would have been expected from the *in vitro* investigations. There are several potential factors that could explain why the clear inhibitory properties of E2 and E-E2 observed *in vitro* were less prominent *in vivo*. E2 is highly metabolized and has a low bioavailability, which could prevent the E2 concentration from being high enough to inhibit Pat1 as it was given as a 4.5 min pre-dose. Also, the hormones were dosed in 96 % ethanol due to the limited solubility of E2 and E-E2 in aqueous medium, where the solubility is around 3.9  $\text{mg L}^{-1}$  and 9.2  $\text{mg L}^{-1}$ , respectively<sup>45</sup>. As ethanol is rapidly absorbed from the gastrointestinal system<sup>46</sup>, there is a possibility that E and E-E2 could precipitate in the gastrointestinal tract as ethanol was absorbed, thereby lowering the amount in solution. Another potential difference between the *in vitro* and *in vivo* experiments is that while the *in vitro* studies were performed on human PAT1, the *in vivo* experiments were performed in rat expressing the rat ortholog Pat1, raising the possibility of a species difference in binding of E-E2 to PAT1 versus Pat1. However,

the findings presented here suggest that the potential interaction between E2 and PAT1 drug substances such as gaboxadol, vigabatrin, and  $\delta$ -aminolevulinic acid should not be overlooked but investigated further, preferably in human.

In conclusion, the present study is the first to show that 17- $\alpha$ -ethinyl-estradiol decreased PAT1-mediated cellular uptake and reduced transepithelial absorptive permeability of PAT1 substrates *in vitro* and that this to a certain degree also leads to an effect on the pharmacokinetic profile of taurine *in vivo* after oral administration.

### Acknowledgement

The staff at the animal facility at H. Lundbeck A/S for help with performing the animal study. Parts of the experimental work was performed during CUNs employment at University of Copenhagen supervising TK and SM. RH was employed at H. Lundbeck A/S, Denmark in that period.

**Compliance with ethical standards.** All studies conducted here comply with ethical standards

**Conflict of interest.** The authors declare that they have no conflict of interest

**Ethical approval.** All procedures performed in studies involving animals were in accordance with the ethical standards of the institutional animal ethics committee in accordance with Danish law regulating animal experiments and in compliance with EC Directive 2010/63/EU and the NIH guidelines on animal welfare

### Figure Legends:

**Figure 1. Uptake in Caco-2 cells measured in the absence or presence of 100  $\mu$ M estradiol (E2) or E-estradiol (E-E2).** **A.** The uptake rate of 13.3 nM [ $^3$ H]-proline (1.0  $\mu$ Ci ml $^{-1}$ ) at pH 6.0 or 7.4 (n=4-7) **B.** The uptake rate of 52 nM [ $^3$ H]-taurine (1.0  $\mu$ Ci ml $^{-1}$ ) with 100  $\mu$ M non-labelled taurine at pH 6.0 or 7.4 (n=3-4). **C.** The uptake rate of 35.3 nM [ $^3$ H]-5-HT (1.0  $\mu$ Ci ml $^{-1}$ ) at pH 6.0 (n=3-6) **D.** The uptake rate of 17.9  $\mu$ M [ $^{14}$ C]-Gly-Sar (1.0  $\mu$ Ci ml $^{-1}$ ) at pH 6.0 (n=3-5) **E.** The uptake rate of 65.8 nM [ $^3$ H]-propranolol (1.0  $\mu$ Ci ml $^{-1}$ ) at pH 6.0 (n=3). Each value represents the mean  $\pm$  SEM. One-way ANOVA test using Dunnett's multiple comparison post-test using  $p < 0.05$  (\*) was for statistical testing.

**Figure 2.** Concentrations-dependent inhibition of substrate uptake in Caco-2 cells with 1-316.2  $\mu\text{M}$  estradiol or E-estradiol. **A.** Inhibition of 13.3 nM [ $^3\text{H}$ ]-proline uptake rate ( $1.0 \mu\text{Ci ml}^{-1}$ ) at pH 6.0. **B.** Inhibition of 99 nM [ $^3\text{H}$ ]-taurine uptake ( $1.0 \mu\text{Ci ml}^{-1}$ ) with 100  $\mu\text{M}$  non-labelled taurine at pH 6.0. **C.** Inhibition of 99 nM [ $^3\text{H}$ ]-taurine uptake ( $1.0 \mu\text{Ci ml}^{-1}$ ) at pH 7.4. The  $\text{IC}_{50}$  value was determined by fitting data to a sigmoidal dose-response curve for inhibition, GraphPad Prism version 7.02. Each value represents the mean  $\pm$  SEM ( $n=4$ ).

**Figure 3.** Two-electrode voltage clamp (TEVC) measurements in PAT1 expressing *Xenopus Laevis* Oocytes. Estradiol (E2) or E-estradiol (E-E2) inhibition of proline induced current in PAT1 expressing oocytes at a holding potential of -60 mV and continuously perfused with Ringer's solution pH 6.0. 20 mM proline alone or with either 50  $\mu\text{M}$  estradiol or E-estradiol were applied (as indicated by the horizontal bars), and the change in current was monitored. Upper trace, water injected oocyte; lower trace, oocyte injected with *SLC36A1 cRNA*. The traces are representative of experiments performed on 6 different oocytes.

**Figure 4.** Concentrations-dependent substrate uptake in Caco-2 cells in the absence or presence of estradiol (E2) or E-estradiol (E-E2) at pH 6.0. **A.** The uptake rate of 13.3 nM [ $^3\text{H}$ ]-proline ( $1.0 \mu\text{Ci ml}^{-1}$ ) with 0-20 mM L-proline (o), in the presence of 10  $\mu\text{M}$  E2 (●), 50  $\mu\text{M}$  E2 (■) or 100  $\mu\text{M}$  E2 (▲) ( $n=3-5$ ). **B.** The uptake rate of 52 nM [ $^3\text{H}$ ]-taurine ( $1.0 \mu\text{Ci ml}^{-1}$ ) with 0-50 mM taurine (o), in the presence of 50  $\mu\text{M}$  E2 ( $\Delta$ ) or 50  $\mu\text{M}$  E-E2 (■) ( $n=4$ ). Each value represents the mean  $\pm$  SEM. The solid lines show the fit of the resulting data to Michaelis-Menten like kinetics.

**Figure 5.** Time-dependent treatment of estradiol (E2) or E-estradiol (E-E2) in Caco-2 cells. **A.** The uptake rate of 13.3 nM [ $^3\text{H}$ ]-proline ( $1.0 \mu\text{Ci ml}^{-1}$ ) at pH 6.0 with and without pre-incubation of 100  $\mu\text{M}$  estradiol (E2) or E-estradiol (E-E2) at pH 7.4 for 1 hour before the proline uptake study. **B.** The uptake rate of 13.3 nM [ $^3\text{H}$ ]-proline ( $1.0 \mu\text{Ci ml}^{-1}$ ) at pH 6.0 with and without pre-incubation of 100  $\mu\text{M}$  estradiol (E2), pH 7.4 for 5 minutes, followed by incubation with buffer at the indicated time points, before the 5 min proline uptake study. Each value represents the mean  $\pm$  SEM ( $n=3$ ). Statistical significance between the control (pre-incubation and uptake without E2) and the other pre-incubation conditions was tested using a one-way ANOVA test followed by a Dunnett's multiple comparisons test.

**Figure 6.** Uptake in Caco-2 cells measured in the absence or presence of other steroid-containing structures or modulators of estradiol signaling. **A.** The uptake rate of 13.3 nM [ $^3\text{H}$ ]-proline ( $1.0 \mu\text{Ci mL}^{-1}$ ) in the absence or presence of 100  $\mu\text{M}$  estradiol (E2), E-estradiol (E-E2), chenodeoxycholic acid (CDCA), cholesterol, dexamethasone (Dx), glycocholic acid (GCA), and taurocholic acid (TCA) at pH 6.0. **B.** The uptake rate of 13.3 nM [ $^3\text{H}$ ]-proline ( $1.0 \mu\text{Ci mL}^{-1}$ ) in the absence or presence of 100 nM endoxifen, 100  $\mu\text{M}$  fulvestrant, 300 nM staurosporine, 1  $\mu\text{M}$  thapsigargin and 1  $\mu\text{M}$  PI-103 at pH 6.0. The cells were pre-incubated with the compounds in 10 mM HEPES in HBSS (pH 7.4) for 2 hours. Each value represents the mean  $\pm$  SEM ( $n=3$ ). Statistical significance between the control and the tested compounds within the three conditions was tested using a one-way ANOVA test followed by a Dunnett's multiple comparisons test.

**Figure 7.** Apparent absorptive transepithelial permeability ( $P_{\text{app}}$ ) across Caco-2 cell monolayers of various SLC substrates in the absence or presence of 100  $\mu\text{M}$  estradiol (E2) or E-estradiol (E-E2). Transport was measured from the apical (A) to the basolateral (B) side. Donor solutions contained  $1.0 \mu\text{Ci mL}^{-1}$  radiolabeled compound. **A.** Taurine permeability at pH 6.0/7.4 (A/B) with a donor concentration of 100  $\mu\text{M}$  taurine. **B.** Taurine permeability at pH 7.4/7.4 (A/B) with a donor concentration of 99 nM taurine. **C.** Proline permeability at pH 6.0/7.4 (A/B) with a donor concentration of 1 mM proline. **D.** Gly-Sar permeability at pH 6.0/7.4 (A/B) with a donor concentration of 17.9  $\mu\text{M}$  Gly-Sar. **E.** 5-HT permeability at pH 6.0/7.4 (A/B) with a donor concentration of 35.3 nM 5-HT. **F.** Propranolol permeability at pH 6.0/7.4 (A/B) with a donor concentration of 23.9 nM propranolol. Each value represents the mean  $\pm$  SEM ( $n=3-4$ ). Statistical analysis was done with two-way ANOVA with Dunnett post-test and the significance level \* ( $P<0.05$ ) compared to the control in each group.

**Figure 8.** Taurine plasma concentration versus time profiles after oral administration of estradiol (E2) (**A**) or E-estradiol (E-E2) (**B**) in various concentrations (shown in legend) followed by oral administration of [ $^3\text{H}$ ]-taurine ( $40.0 \mu\text{Ci kg}^{-1}$ ) and taurine ( $30 \text{ mg kg}^{-1}$ ). Each value represents the mean  $\pm$  SEM of measurements from 4 rats per dosing group, except for the control group consisting of data from 8 animals.

**References**

1. Human genome organization. Solute Carriers, <https://www.genenames.org/data/genegroup/#!/group/752> assessed 0630-2020.
2. Hediger MA, Clemençon B, Burrier RE, Bruford EA 2013. The ABCs of membrane transporters in health and disease (SLC series): introduction. *Mol Aspects Med* 34(2-3):95-107.
3. Hediger MA, Romero MF, Peng JB, Rolfs A, Takanaga H, Bruford EA 2004. The ABCs of solute carriers: physiological, pathological and therapeutic implications of human membrane transport proteins Introduction. *Pflugers Arch* 447(5):465-468.
4. Chen Z, Fei YJ, Anderson CM, Wake KA, Miyauchi S, Huang W, Thwaites DT, Ganapathy V 2003. Structure, function and immunolocalization of a proton-coupled amino acid transporter (hPAT1) in the human intestinal cell line Caco-2. *J Physiol* 546(Pt 2):349-361.
5. Vanslambrouck JM, Broer A, Thavyogarah T, Holst J, Bailey CG, Broer S, Rasko JE 2010. Renal imino acid and glycine transport system ontogeny and involvement in developmental iminoglycinuria. *Biochem J* 428(3):397-407.
6. Agulhon C, Rostaing P, Ravassard P, Sagne C, Triller A, Giros B 2003. Lysosomal amino acid transporter LYAAT-1 in the rat central nervous system: an in situ hybridization and immunohistochemical study. *J Comp Neurol* 462(1):71-89.
7. Sagne C, Agulhon C, Ravassard P, Darmon M, Hamon M, El Mestikawy S, Gasnier B, Giros B 2001. Identification and characterization of a lysosomal transporter for small neutral amino acids. *Proc Natl Acad Sci U S A* 98(13):7206-7211.
8. Jensen A, Figueiredo-Larsen M, Holm R, Broberg ML, Brodin B, Nielsen CU 2014. PAT1 (SLC36A1) shows nuclear localization and affects growth of smooth muscle cells from rats. *Am J Physiol Endocrinol Metab* 306(1):E65-74.
9. Thwaites DT, McEwan GT, Brown CD, Hirst BH, Simmons NL 1994. L-alanine absorption in human intestinal Caco-2 cells driven by the proton electrochemical gradient. *J Membr Biol* 140(2):143-151.
10. Thwaites DT, McEwan GT, Cook MJ, Hirst BH, Simmons NL 1993. H(+)-coupled (Na(+)-independent) proline transport in human intestinal (Caco-2) epithelial cell monolayers. *FEBS Lett* 333(1-2):78-82.

11. Thwaites DT, Basterfield L, McCleave PM, Carter SM, Simmons NL 2000. Gamma-Aminobutyric acid (GABA) transport across human intestinal epithelial (Caco-2) cell monolayers. *Br J Pharmacol* 129(3):457-464.
12. Larsen M, Holm R, Jensen KG, Brodin B, Nielsen CU 2009. Intestinal gaboxadol absorption via PAT1 (SLC36A1): modified absorption in vivo following co-administration of L-tryptophan. *Br J Pharmacol* 157(8):1380-1389.
13. Nohr MK, Frolund S, Holm R, Nielsen CU 2014. Pharmacokinetic aspects of the anti-epileptic drug substance vigabatrin: focus on transporter interactions. *Ther Deliv* 5(8):927-942.
14. Nohr MK, Juul RV, Thale ZI, Holm R, Kreilgaard M, Nielsen CU 2015. Is oral absorption of vigabatrin carrier-mediated? *Eur J Pharm Sci* 69:10-18.
15. Broberg M, Holm R, Tonsberg H, Frolund S, Ewon KB, Nielsen A, Brodin B, Jensen A, Kall MA, Christensen KV, Nielsen CU 2012. Function and expression of the proton-coupled amino acid transporter PAT1 along the rat gastrointestinal tract: implications for intestinal absorption of gaboxadol. *Br J Pharmacol* 167(3):654-665.
16. Nielsen CU, Bjerg M, Ulaganathan N, Holm R 2017. Oral and intravenous pharmacokinetics of taurine in sprague-dawley rats: the influence of dose and the possible involvement of the proton-coupled amino acid transporter, PAT1, in oral taurine absorption. *Physiol Rep* 5(19).
17. Anderson CM, Jevons M, Thangaraju M, Edwards N, Conlon NJ, Woods S, Ganapathy V, Thwaites DT 2010. Transport of the photodynamic therapy agent 5-aminolevulinic acid by distinct H<sup>+</sup>-coupled nutrient carriers coexpressed in the small intestine. *J Pharmacol Exp Ther* 332(1):220-228.
18. Abbot EL, Grenade DS, Kennedy DJ, Gatfield KM, Thwaites DT 2006. Vigabatrin transport across the human intestinal epithelial (Caco-2) brush-border membrane is via the H<sup>+</sup>-coupled amino-acid transporter hPAT1. *Br J Pharmacol* 147(3):298-306.
19. Foltz M, Mertl M, Dietz V, Boll M, Kottra G, Daniel H 2005. Kinetics of bidirectional H<sup>+</sup> and substrate transport by the proton-dependent amino acid symporter PAT1. *Biochem J* 386(Pt 3):607-616.
20. Anderson CM, Howard A, Walters JR, Ganapathy V, Thwaites DT 2009. Taurine uptake across the human intestinal brush-border membrane is via two transporters: H<sup>+</sup>-coupled PAT1 (SLC36A1) and Na<sup>+</sup>- and Cl<sup>-</sup>-dependent TauT (SLC6A6). *J Physiol* 587(Pt 4):731-744.

21. Thwaites DT, McEwan GT, Simmons NL 1995. The role of the proton electrochemical gradient in the transepithelial absorption of amino acids by human intestinal Caco-2 cell monolayers. *J Membr Biol* 145(3):245-256.
22. In Vitro Drug Interaction Studies — Cytochrome P450 Enzyme- and Transporter-Mediated Drug Interactions Guidance for Industry. 2020, CDER, U.S. Department of Health and Human Services Food and Drug Administration. p 1-43.
23. Guideline on the investigation of drug interactions. 2012, CPMP/EWP/560/95/Rev. 1 Corr. 2, Committee for Human Medicinal Products (CHMP), European Medicines Agency.
24. Metzner L, Kottra G, Neubert K, Daniel H, Brandsch M 2005. Serotonin, L-tryptophan, and tryptamine are effective inhibitors of the amino acid transport system PAT1. *FASEB J* 19(11):1468-1473.
25. Frolund S, Langthaler L, Kall MA, Holm R, Nielsen CU 2012. Intestinal drug transport via the proton-coupled amino acid transporter PAT1 (SLC36A1) is inhibited by Gly-X(aa) dipeptides. *Mol Pharm* 9(9):2761-2769.
26. Nielsen CU, Frolund S, Abdulhadi S, Sari H, Langthaler L, Nohr MK, Kall MA, Brodin B, Holm R 2013. Sertraline inhibits the transport of PAT1 substrates in vivo and in vitro. *Br J Pharmacol* 170(5):1041-1052.
27. Shan L, Yang Y, Wang J, Zuo J, Dong X, Li C, Li D 2012. Estradiol inhibits the activity of proton-coupled amino acid transporter PAT1 expressed in *Xenopus oocytes*. *Eur J Pharmacol* 695(1-3):34-39.
28. Levin VA, Jiang X, Kagan R 2018. Estrogen therapy for osteoporosis in the modern era. *Osteoporos Int* 29(5):1049-1055.
29. Di Zazzo E, Galasso G, Giovannelli P, Di Donato M, Castoria G 2018. Estrogens and Their Receptors in Prostate Cancer: Therapeutic Implications. *Front Oncol* 8:2.
30. O'Connell MB 1995. Pharmacokinetic and pharmacologic variation between different estrogen products. *J Clin Pharmacol* 35(9S):18S-24S.
31. Fotherby K 1996. Bioavailability of orally administered sex steroids used in oral contraception and hormone replacement therapy. *Contraception* 54(2):59-69.

32. Frolund S, Holm R, Brodin B, Nielsen CU 2010. The proton-coupled amino acid transporter, SLC36A1 (hPAT1), transports Gly-Gly, Gly-Sar and other Gly-Gly mimetics. *Br J Pharmacol* 161(3):589-600.
33. Larsen M, Larsen BB, Frolund B, Nielsen CU 2008. Transport of amino acids and GABA analogues via the human proton-coupled amino acid transporter, hPAT1: characterization of conditions for affinity and transport experiments in Caco-2 cells. *Eur J Pharm Sci* 35(1-2):86-95.
34. Nohr MK, Holm R, Thale ZI, Nielsen CU 2014. Intestinal absorption of the antiepileptic drug substance vigabatrin in Gottingen mini-pigs is unaffected by co-administration of amino acids. *Int J Pharm* 466(1-2):18-20.
35. Nielsen CU, Andersen R, Brodin B, Frokjaer S, Taub ME, Steffansen B 2001. Dipeptide model prodrugs for the intestinal oligopeptide transporter. Affinity for and transport via hPepT1 in the human intestinal Caco-2 cell line. *J Control Release* 76(1-2):129-138.
36. Disshon KA, Boja JW, Dluzen DE 1998. Inhibition of striatal dopamine transporter activity by 17beta-estradiol. *Eur J Pharmacol* 345(2):207-211.
37. Cairrao E, Alvarez E, Carvas JM, Santos-Silva AJ, Verde I 2012. Non-genomic vasorelaxant effects of 17beta-estradiol and progesterone in rat aorta are mediated by L-type Ca<sup>2+</sup> current inhibition. *Acta Pharmacol Sin* 33(5):615-624.
38. Oz M, Zhang L, Spivak CE 2002. Direct noncompetitive inhibition of 5-HT(3) receptor-mediated responses by forskolin and steroids. *Arch Biochem Biophys* 404(2):293-301.
39. Nadal A, Diaz M, Valverde MA 2001. The estrogen trinity: membrane, cytosolic, and nuclear effects. *News Physiol Sci* 16:251-255.
40. Kennedy DJ, Leibach FH, Ganapathy V, Thwaites DT 2002. Optimal absorptive transport of the dipeptide glycylsarcosine is dependent on functional Na<sup>+</sup>/H<sup>+</sup> exchange activity. *Pflugers Arch* 445(1):139-146.
41. Thwaites DT, Ford D, Glanville M, Simmons NL 1999. H<sup>(+)</sup>/solute-induced intracellular acidification leads to selective activation of apical Na<sup>(+)</sup>/H<sup>(+)</sup> exchange in human intestinal epithelial cells. *J Clin Invest* 104(5):629-635.
42. Anderson CM, Thwaites DT 2005. Indirect regulation of the intestinal H<sup>+</sup>-coupled amino acid transporter hPAT1 (SLC36A1). *J Cell Physiol* 204(2):604-613.



43. Chojjookhuu N, Sato Y, Nishino T, Endo D, Hishikawa Y, Koji T 2012. Estrogen-dependent regulation of sodium/hydrogen exchanger-3 (NHE3) expression via estrogen receptor beta in proximal colon of pregnant mice. *Histochem Cell Biol* 137(5):575-587.
44. Rotti K, Stevens J, Watson D, Longcope C 1975. Estriol concentrations in plasma of normal, non-pregnant women. *Steroids* 25(6):807-816.
45. Johnson ASMJAJDWaBB 2006. Aqueous Solubilities of Estrone,  $17\beta$ -Estradiol,  $17\alpha$ -Ethinylestradiol, and Bisphenol A. *Journal of Chemical & Engineering Data* 51 (3):879-881.
46. Siegers CP, Strubelt O, Back G 1972. Inhibition by caffeine of ethanol absorption in rats. *Eur J Pharmacol* 20(2):181-187.

Journal Pre-proof

**Table 1**

Kinetic parameters for uptake of proline and taurine in the presence estradiol (E2) and ethinyl-estradiol (E-E2). The marked values (\*) are significantly different from the  $V_{\max}$  obtained in the absence of E2 and E-E2 evaluated by a one-way ANOVA test followed by Dunnett's multiple comparisons test. The transporter-mediated intrinsic clearance (CL) was calculated as  $V_{\max}/K_m$ .

	$K_{m, app}$ (mM)	$V_{max, app}$ (nmol cm <sup>-2</sup> min <sup>-1</sup> )	Intrinsic CL ( $\mu$ L min <sup>-1</sup> pr. cm <sup>2</sup> )
<b>Proline</b>	1.80 ± 0.24	1.54 ± 0.05	0.856
+ 10 $\mu$ M E2	2.54 ± 0.76	1.41 ± 0.11	0.551
+ 50 $\mu$ M E2	3.67 ± 1.57	0.76 ± 0.15*	0.207
+ 100 $\mu$ M E2	3.81 ± 1.37	0.67 ± 0.07*	0.176
<b>Taurine</b>	8.45 ± 1.39	3.75 ± 0.21	0.444
+ 50 $\mu$ M E2	7.15 ± 3.22	0.87 ± 0.14*	0.122
+ 50 $\mu$ M E-E2	11.28 ± 2.03	1.55 ± 0.10*	0.137

**Table 2****Pharmacokinetic parameters of taurine after oral co-administration of estradiol to rats.**


Rats were orally administered estradiol (E2) in various pre-doses followed by oral administration of 40  $\mu\text{Ci kg}^{-1}$  [ $^3\text{H}$ ]-taurine and 30  $\text{mg kg}^{-1}$  taurine. The pharmacokinetic parameters  $C_{\text{max}}$ , AUC and  $k_e$  are expressed as mean  $\pm$  SEM, while  $t_{\text{max}}$  is expressed as the median [ $Q_1$ ;  $Q_3$ ] (25 % and 75 % percentile) of measurements from 4 rats per dosing group, except 8 rats for the control group (0  $\text{mg kg}^{-1}$  E2).



E2 dose ( $\text{mg kg}^{-1}$ )	Taurine dose ( $\text{mg kg}^{-1}$ )	$t_{\text{max}}$ (min)	$C_{\text{max}}$ ( $\mu\text{g mL}^{-1}$ )	AUC <sub>0-tlast</sub> ( $\mu\text{g min mL}^{-1}$ )	$k_e$ ( $10^{-3} \text{ min}^{-1}$ )
0	30	15.0 [15.0; 15.0]	15.6 $\pm$ 1.0	1181 $\pm$ 62	4.33 $\pm$ 0.250
0.07	30	15.0 [15.0; 15.0]	11.3 $\pm$ 2.0	878 $\pm$ 61	2.78 $\pm$ 0.220
0.27	30	15.0 [15.0; 15.0]	10.4 $\pm$ 1.8	964 $\pm$ 145	2.22 $\pm$ 0.573
1.26	30	15.0 [15.0; 15.0]	15.5 $\pm$ 1.2	1038 $\pm$ 51	3.79 $\pm$ 0.147
2.72	30	15.0 [15.0; 37.5]	12.3 $\pm$ 4.1	1086 $\pm$ 102	2.90 $\pm$ 0.783
6.78	30	15.0 [15.0; 15.0]	13.4 $\pm$ 1.3	1025 $\pm$ 89	2.56 $\pm$ 0.353


**Table 3**  
**Pharmacokinetic parameters of taurine after oral co-administration of ethinyl-estradiol to rats.**


Rats were orally administered ethinyl-estradiol (E-E2) in various doses followed by oral administration of 40  $\mu\text{Ci kg}^{-1}$  [ $^3\text{H}$ ]-taurine and 30  $\text{mg kg}^{-1}$  taurine. The pharmacokinetic parameters  $C_{\text{max}}$ , AUC and  $k_e$  are expressed as mean  $\pm$  SEM, while  $t_{\text{max}}$  is expressed as the median [Q<sub>1</sub>; Q<sub>3</sub>] (25 % and 75 % percentile) of measurements from 4 rats per dosing group, except 8 rats for the control group (0  $\text{mg kg}^{-1}$  E-E2).

E-E2 dose ( $\text{mg kg}^{-1}$ )	Taurine dose ( $\text{mg kg}^{-1}$ )	$t_{\text{max}}$ (min)	$C_{\text{max}}$ ( $\mu\text{g mL}^{-1}$ )	AUC <sub>0-tlast</sub> ( $\mu\text{g min mL}^{-1}$ )	$k_e$ ( $10^{-3} \text{ min}^{-1}$ )
0	30	15.0 [15.0; 15.0]	15.6 $\pm$ 1.0	1181 $\pm$ 62	4.33 $\pm$ 0.250
0.08	30	15.0 [15.0; 15.0]	16.0 $\pm$ 1.7	1110 $\pm$ 25	4.20 $\pm$ 0.301
0.31	30	15.0 [15.0; 15.0]	13.8 $\pm$ 2.3	1062 $\pm$ 71	2.18 $\pm$ 0.758
1.46	30	22.5 [15.0; 30.0]	11.0 $\pm$ 1.9	1249 $\pm$ 95	3.17 $\pm$ 0.778
7.40	30	22.5 [15.0; 30.0]	10.7 $\pm$ 2.6	956 $\pm$ 58	2.52 $\pm$ 0.510














Journal Pre-proof

